

Italian Emission Inventory 1990 - 2022 Informative Inventory Report 2024



RAPPORTI
400/2024

Italian Emission Inventory 1990 - 2022 Informative Inventory Report 2024

Legal Disclaimer

The Institute for Environmental Protection and Research (ISPRA), together with the 21 Regional Agencies (ARPA) and Provincial Agencies (APPA) for the protection of the environment, as of 14 January 2017 is part of the National Network System for the Protection of the Environment (SNPA), established by the Law June 28, 2016, n.132.

The Institute for Environmental Protection and Research, or persons acting on its behalf, are not responsible for the use that may be made of the information contained in this report.

ISPRA - Institute for Environmental Protection and Research
Via Vitaliano Brancati, 48 – 00144 Rome
www.isprambiente.gov.it

ISPRA, Rapporti 400/2024
ISBN 978-88-448-1213-3

Extracts from this document may be reproduced on the condition that the source is acknowledged.

Graphic design

Cover design: Alessia Marinelli – Ispra – Area Comunicazione

Cover image: graphic processing from an image generated with Stable Diffusion on the Wowzer platform by Angela Fiore e Luisa Mortola

Coordination of the online publication:

Daria Mazzella

ISPRA – Area Comunicazione

Authors

Ernesto Taurino, Antonella Bernetti, Antonio Caputo, Marco Cordella, Riccardo De Lauretis, Ilaria D'Elia (ENEA), Eleonora Di Cristofaro, Andrea Gagna, Barbara Gonella, Federica Moricci, Guido Pellis, Emanuele Peschi, Daniela Romano, Marina Vitullo

Contact:

Ernesto Taurino

telephone +39 0650072568

e-mail ernesto.taurino@isprambiente.it

ISPRA- Institute for Environmental Protection and Research
Environmental Assessment, Control and Sustainability Department
Emissions, Prevention of Atmospheric Impacts and Climate Change Area
Air Emission Inventory Unit
Via V. Brancati, 48 00144 Rome ITALY

Text available on ISPRA website at <http://www.isprambiente.gov.it>

CONTENTS

EXECUTIVE SUMMARY	10
SOMMARIO (Italiano)	11
1 INTRODUCTION	12
1.1 Background information on the convention on long-range transboundary air pollution	12
1.2 National inventory	13
1.3 Institutional arrangements	14
1.4 Inventory preparation process	15
1.5 Methods and data sources	16
1.6 Key categories	18
1.7 QA/QC and verification methods	22
1.8 General uncertainty evaluation	30
1.9 General assessment of completeness	32
2 ANALYSIS OF KEY TRENDS BY POLLUTANT	33
2.1 MAIN POLLUTANTS	33
2.1.1 Sulphur dioxide (SO _x)	33
2.1.2 Nitrogen oxides (NO _x)	35
2.1.3 Ammonia (NH ₃)	37
2.1.4 Non methane volatile organic compounds (NMVOC)	39
2.1.5 Carbon monoxide (CO)	41
2.2 PARTICULATE MATTER	44
2.2.1 PM ₁₀	44
2.2.2 PM _{2.5}	46
2.2.3 Black Carbon (BC)	48
2.3 HEAVY METALS (PB, CD, HG)	50
2.3.1 Lead (Pb)	50
2.3.2 Cadmium (Cd)	51
2.3.3 Mercury (Hg)	52
2.4 PERSISTENT ORGANIC POLLUTANTS (POPs)	54
2.4.1 Polycyclic aromatic hydrocarbons (PAH)	54
2.4.2 Dioxins	55
2.4.3 Hexachlorobenzene (HCB)	56
2.4.4 Polychlorinated biphenyl (PCB)	58
3 ENERGY (NFR SECTOR 1)	60
3.1 OVERVIEW OF THE SECTOR	60
3.2 ENERGY INDUSTRIES	65
3.2.1 Methodological issues	65
3.2.1.1 <i>Public Electricity and Heat Production</i>	65

3.2.1.2	<i>Refineries</i>	66
3.2.1.3	<i>Manufacture of Solid Fuels and Other Energy Industries</i>	67
3.2.2	Time series and key categories	68
3.2.2.1	<i>Public Electricity and Heat production</i>	69
3.2.2.2	<i>Refineries</i>	69
3.2.2.3	<i>Manufacture of Solid Fuels and Other Energy Industries</i>	69
3.2.3	QA/QC and verification	70
3.2.4	Recalculations	72
3.2.5	Planned improvements	72
3.3	MANUFACTURING INDUSTRIES AND CONSTRUCTION	73
3.3.1	Methodological issues	73
3.3.1.1	<i>Iron and steel</i>	74
3.3.1.2	<i>Non-ferrous metals</i>	75
3.3.1.3	<i>Chemicals</i>	76
3.3.1.4	<i>Pulp, paper and print</i>	76
3.3.1.5	<i>Food processing, beverages and tobacco</i>	77
3.3.1.6	<i>Non-metallic minerals</i>	77
3.3.1.7	<i>Other</i>	78
3.3.2	Time series and key categories	78
3.3.2.1	<i>Iron and steel</i>	78
3.3.2.2	<i>Non-ferrous metals</i>	79
3.3.2.3	<i>Chemicals</i>	79
3.3.2.4	<i>Pulp, paper and print</i>	79
3.3.2.5	<i>Food processing, beverages and tobacco</i>	79
3.3.2.6	<i>Non-metallic minerals</i>	79
3.3.2.7	<i>Other</i>	79
3.3.3	QA/QC and verification	80
3.3.4	Recalculations	80
3.3.5	Planned improvements	80
3.4	AVIATION (NFR SUBSECTOR 1.A.3.A)	81
3.4.1	Overview	81
3.4.2	Methodological issues	81
3.4.3	Time series and key categories	83
3.4.4	QA/QC and Uncertainty	85
3.4.5	Recalculations	86
3.4.6	Planned improvements	86
3.5	ROAD TRANSPORT (NFR SUBSECTOR 1.A.3.B)	87
3.5.1	Overview	87
3.5.2	Methodological issues	87
3.5.2.1	<i>Exhaust emissions</i>	88

3.5.2.2	<i>Evaporative emissions</i>	90
3.5.2.3	<i>Emissions from automobile tyre and brake wear</i>	90
3.5.2.4	<i>Emissions from automobile road abrasion</i>	90
3.5.3	Activity data	90
3.5.4	Time series and key categories	96
3.5.5	QA/QC and Uncertainty	102
3.5.6	Recalculation	102
3.5.7	Planned improvements	103
3.6	RAILWAYS (NFR SUBSECTOR 1.A.3.C)	104
3.7	NAVIGATION (NFR SUBSECTOR 1.A.3.D)	105
3.7.1	Overview	105
3.7.2	Methodological issues	105
3.7.3	Time series and key categories	106
3.7.4	QA/QC and Uncertainty	107
3.7.5	Recalculations	108
3.7.6	Planned improvements	108
3.8	PIPELINE COMPRESSORS (NFR SUBSECTOR 1.A.3.E)	109
3.9	CIVIL SECTOR: SMALL COMBUSTION AND OFF-ROAD VEHICLES (NFR SUBSECTOR 1.A.4 - 1.A.5)	110
3.9.1	Overview	110
3.9.2	Activity data	110
3.9.3	Methodological issues	112
3.9.3.1	<i>NO_x emissions from gas powered plants in the civil sector</i>	113
3.9.3.2	<i>Emissions from wood combustion in the civil sector</i>	114
3.9.4	Time series and key categories	116
3.9.5	QA/QC and Uncertainty	118
3.9.6	Recalculation	118
3.9.7	Planned improvements	118
3.10	FUGITIVE EMISSIONS (NFR SUBSECTOR 1.B)	119
3.10.1	Overview	119
3.10.2	Methodological issues	119
3.10.2.1	<i>Fugitive emissions from natural gas distribution (1.B.2b)</i>	120
3.10.3	Time series and key categories	122
3.10.4	QA/QC and Uncertainty	122
3.10.5	Recalculation	122
3.10.6	Planned improvements	122
4	IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2)	123
4.1	OVERVIEW OF THE SECTOR	123
4.2	METHODOLOGICAL ISSUES	123
4.2.1	Mineral products (2A)	124
4.2.2	Chemical industry (2B)	125

4.2.3	Metal production (2C)	126
4.2.4	Other production (2G – 2H – 2I – 2L)	128
4.3	TIME SERIES AND KEY CATEGORIES	130
4.3.1	Mineral products (2A)	130
4.3.2	Chemical industry (2B)	131
4.3.3	Metal production (2C)	132
4.3.4	Other production (2G – 2H – 2I – 2L)	133
4.4	QA/QC AND VERIFICATION	134
4.5	RECALCULATIONS	135
4.5.1	Mineral industry (2A)	135
4.5.2	Chemical industry (2B)	135
4.5.3	Metal industry (2C)	136
4.5.4	Other product use (fireworks and tobacco) (2G)	136
4.5.5	Other industrial processes (2H)	136
4.6	PLANNED IMPROVEMENTS	136
5	IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2)	137
5.1	OVERVIEW OF THE SECTOR	137
5.2	METHODOLOGICAL ISSUES	138
5.2.1	Domestic solvent use (2D3a)	138
5.2.2	Road paving with asphalt (2D3b)	139
5.2.3	Asphalt Roofing (2D3c)	139
5.2.4	Decorative coating (2D3d1)	140
5.2.5	Industrial coating (2D3d2)	140
5.2.6	Degreasing (2D3e)	141
5.2.7	Dry cleaning (2D3f)	141
5.2.8	Chemical products, manufacture and processing (2D3g)	142
5.2.9	Other product use (2D3i)	143
5.3	TIME SERIES AND KEY CATEGORIES	143
5.4	QA/QC AND VERIFICATION	148
5.5	RECALCULATIONS	149
5.6	PLANNED IMPROVEMENTS	150
6	AGRICULTURE (NFR SECTOR 3)	151
6.1	OVERVIEW OF THE SECTOR	151
6.2	METHODOLOGICAL ISSUES	153
6.2.1	Manure management (3B)	153
6.2.1.1	<i>Dairy cattle (3B1a)</i>	155
6.2.1.2	<i>Swine (3B3)</i>	157
6.2.1.3	<i>Poultry (3B4g)</i>	159
6.3	AGRICULTURAL SOILS (3D)	164
6.4	FIELD BURNING OF AGRICULTURAL RESIDUES (3F)	168
6.5	TIME SERIES AND KEY CATEGORIES	168

6.6	QA/QC AND VERIFICATION	174
6.7	RECALCULATIONS	174
6.8	PLANNED IMPROVEMENTS	176
7	WASTE (NFR SECTOR 5)	177
7.1	OVERVIEW OF THE SECTOR	177
7.2	METHODOLOGICAL ISSUES	179
7.2.1	Solid waste disposal on land (5A)	179
7.2.2	Biological treatment of waste (5B)	180
7.2.3	Waste Incineration (5C1a – 5C1b)	181
7.2.4	Cremation of corpses (5C1bv)	183
7.2.5	Small scale waste burning (5C2)	185
7.2.6	Wastewater treatments (5D)	187
7.2.7	Other waste (5E)	189
7.3	TIME SERIES AND KEY CATEGORIES	190
7.4	RECALCULATIONS	197
7.5	PLANNED IMPROVEMENTS	199
8	OTHER (6A)	200
8.1	TIME SERIES AND KEY CATEGORIES	200
8.2	QA/QC AND VERIFICATIONS	200
8.3	RECALCULATIONS	201
8.4	PLANNED IMPROVEMENTS	201
9	RECALCULATIONS AND IMPROVEMENTS	202
9.1	RECALCULATIONS	202
9.2	PLANNED IMPROVEMENTS	204
10	PROJECTIONS	206
10.1	THE NATIONAL FRAMEWORK	206
10.2	INPUT SCENARIOS	207
10.2.1	The energy scenario	207
10.2.2	The scenario of non-energy activities	211
10.2.3	The control strategy definition	212
10.3	THE HARMONIZATION PROCESS	214
10.4	THE EMISSION SCENARIO	217
10.5	THE NEC EMISSION TARGETS	222
11	REPORTING OF GRIDDED EMISSIONS AND LPS	224
11.1	FUGITIVE	225
11.2	TRANSPORT	225
11.3	AGRICULTURE	226
11.4	LPS DATA	228
12	REFERENCES	229
12.1	INTRODUCTION	229
12.2	ANALYSIS OF KEY TRENDS BY POLLUTANT	230
12.3	ENERGY (NFR SECTOR 1)	231

12.4	IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2)	235
12.5	IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2)	236
12.6	AGRICULTURE (NFR SECTOR 3)	237
12.7	WASTE (NFR SECTOR 5)	240
12.8	OTHER (NFR SECTOR 6)	242
12.9	RECALCULATIONS AND IMPROVEMENTS	242
12.10	PROJECTIONS	243
12.11	REPORTING OF GRIDDED EMISSIONS AND LPS	243
	APPENDIX 1 SUMMARY INFORMATION ON CONDENSABLE IN PM	245
	APPENDIX 2 UNCERTAINTY ASSESSMENT	246
	APPENDIX 3 PROJECTED ACTIVITY DATA	252

EXECUTIVE SUMMARY

The Italian Informative Inventory Report (IIR) is edited in the framework of the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP). It contains information on the Italian inventory up to the year 2022, including an explanation of methodologies, data sources, QA/QC activities and verification processes carried out during the inventory compilation, with an analysis of emission trends and a description of key categories.

The aim of the document is to facilitate understanding of the calculation of the Italian air pollutant emission data, hence providing a common mean for comparing the relative contribution of different emission sources and supporting the identification of reduction policies.

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the emission inventory submission to CLRTAP, as well as to the United Nations Framework Convention on Climate Change (UNFCCC) and oversees all the work related to inventory compilation.

In particular, in compliance with the LRTAP Convention, Italy has to submit annually data on national emissions of SO_x, NO_x, NMVOC, CO and NH₃, particulate matter and various heavy metals and POPs. The submission consists of the national emission inventory, communicated through compilation of the Nomenclature For Reporting (NFR), and the informative inventory report (IIR) to ensure the properties of transparency, consistency, comparability, completeness and accuracy.

In the period 1990-2022, emissions from almost all the pollutants described in this report show a downward trend. Reductions are especially relevant for the main pollutants (SO_x -95%; NO_x -71%; CO -72%; NMVOC -58%), for BC (-63%), cadmium (-63%), mercury (-64%), lead (-95%) and hexachlorobenzene (-91%). The major drivers for the trend are reductions in the industrial and road transport sectors, due to the implementation of various European Directives which introduced new technologies, plant emission limits, the limitation of sulphur content in liquid fuels and the shift to cleaner fuels. Emissions have also decreased for the improvement of energy efficiency as well as the promotion of renewable energy.

The energy sector is the main source of emissions in Italy with a share of more than 80%, including fugitive emissions, for many pollutants (SO_x 92%; NO_x 92%; CO 93%; PM_{2.5} 85%; BC 90%; PAH 86%, HCB 91%). The industrial processes sector is an important source of emissions specifically related to the iron and steel production, at least for particulate matter, heavy metals and POPs, whereas significant emissions of SO_x derive from carbon black and sulphuric acid production; on the other hand, the solvent and other product use sector is characterized by NMVOC emissions. The agriculture sector is the main source of NH₃ emissions in Italy with a share of 90% in national total. Finally, the waste sector, specifically waste incineration, is a relevant source for BC (9%), Cd (8%), HCB (9%) and dioxins (17%).

Emission figures of the Italian emission inventory and other related documents are publicly available at <https://emissioni.sina.isprambiente.it/inventario-nazionale/> .

SOMMARIO (Italiano)

L'*Informative Inventory Report* (IIR) è redatto nel quadro della Convenzione UNECE (*United Nations Economic Commission for Europe*) sull'inquinamento atmosferico transfrontaliero a lunga distanza (CLRTAP). Contiene informazioni sull'inventario italiano fino all'anno 2022, compresa una spiegazione delle metodologie, delle fonti dei dati, delle attività di QA/QC e dei processi di verifica effettuati durante la compilazione dell'inventario, con un'analisi dell'andamento delle emissioni e una descrizione delle principali categorie.

Lo scopo del documento è quello di facilitare la comprensione delle metodologie per la stima delle emissioni degli inquinanti atmosferici in Italia, fornendo così uno strumento comune per confrontare il contributo relativo delle diverse fonti di emissione e supportare l'individuazione di politiche di riduzione.

L'Istituto Superiore per la Protezione e la Ricerca Ambientale (ISPRA) ha la responsabilità generale della presentazione dell'inventario delle emissioni al CLRTAP, nonché alla Convenzione quadro delle Nazioni Unite sui cambiamenti climatici (UNFCCC) ed è responsabile di tutto il lavoro relativo alla compilazione dell'inventario.

In particolare, in ottemperanza alla Convenzione LRTAP, l'Italia deve presentare annualmente i dati sulle emissioni nazionali di SO_x, NO_x, NMVOC, CO e NH₃, e vari metalli pesanti e POP. La *submission* è costituita dall'inventario nazionale delle emissioni, comunicato attraverso la compilazione del Nomenclature For Reporting (NFR) e dal presente rapporto (IIR) per garantire le proprietà di trasparenza, coerenza, comparabilità, completezza e accuratezza.

Nel periodo 1990-2022, le emissioni di quasi tutti gli inquinanti descritti in questo rapporto mostrano una tendenza al ribasso. Le riduzioni sono particolarmente rilevanti per i principali inquinanti (SO_x -95%; NO_x -71%; CO -72%; NMVOC -58%), per BC (-63%), cadmio (-63%), mercurio (-64%), piombo (-95%) ed esaclorobenzene (-91%). I principali *driver* del *trend* sono costituiti dalle riduzioni nei settori industriale e del trasporto su strada, dovute all'implementazione di diverse Direttive Europee che hanno introdotto nuove tecnologie, più stringenti limiti di emissione degli impianti, limitazioni del contenuto di zolfo nei combustibili liquidi e quindi il passaggio a combustibili più puliti. Altri fattori determinanti per la riduzione delle emissioni sono stati il miglioramento dell'efficienza energetica e la promozione delle energie rinnovabili.

Il settore energetico è la principale fonte di emissioni in Italia con una quota superiore all'80%, comprese le emissioni fuggitive, per molti inquinanti (SO_x 92%; NO_x 92%; CO 93%; PM_{2.5} 85%; BC 90%; PAH 86%, HCB 91%). Il settore dei processi industriali è un'importante fonte di emissioni specificamente legate alla produzione siderurgica, almeno per particolato, metalli pesanti e POP, mentre significative emissioni di SO_x derivano dalla produzione di nerofumo e acido solforico; il settore dell'uso di solventi e altri prodotti è invece caratterizzato da emissioni di COVNM. Il settore agricolo è la principale fonte di emissioni di NH₃ in Italia con una quota del 90% sul totale nazionale. Infine, il settore dei rifiuti, in particolare l'incenerimento dei rifiuti, è una fonte rilevante di BC (9%), Cd (8%), HCB (9%) e diossine (17%).

I dati sulle emissioni dell'inventario italiano delle emissioni e altri documenti correlati sono disponibili al pubblico su <https://emissioni.sina.isprambiente.it/inventario-nazionale/> .

1 INTRODUCTION

1.1 Background information on the convention on long-range transboundary air pollution

The 1979 Geneva Convention on Long-range Transboundary Air Pollution, contributing to the development of international environmental law, is one of the fundamental international means for the protection of the human health and the environment through the intergovernmental cooperation. The fact that air pollutants could travel several thousands of kilometers before deposition and damage occurred outlined the need for international cooperation. In November 1979, in Geneva, 34 Governments and the European Community (EC) signed the Convention. The Convention on Long-range Transboundary Air Pollution was ratified by Italy in the year 1982 and entered into force in 1983. It has been extended by the following eight specific protocols:

- The 1984 Protocol on Long-term Financing of the Cooperative Programme for Monitoring and Evaluation of the Long-range Transmission of Air Pollutants in Europe (EMEP); 42 Parties. Entered into force on 28th January 1988.
- The 1985 Protocol on the Reduction of Sulphur Emissions or their Transboundary Fluxes by at least 30 per cent; 23 Parties. Entered into force on 2nd September 1987.
- The 1988 Protocol concerning the Control of Nitrogen Oxides or their Transboundary Fluxes; 31 Parties. Entered into force on 14th February 1991.
- The 1991 Protocol concerning the Control of Emissions of Volatile Organic Compounds or their Transboundary Fluxes; 22 Parties. Entered into force on 29th September 1997.
- The 1994 Protocol on Further Reduction of Sulphur Emissions; 27 Parties. Entered into force on 5th August 1998.
- The 1998 Protocol on Heavy Metals; 28 Parties. Entered into force on 29 December 2003.
- The 1998 Protocol on Persistent Organic Pollutants (POPs); 28 Parties. Entered into force on 23rd October 2003.
- The 1999 Protocol to Abate Acidification, Eutrophication and Ground-level Ozone; 23 Parties. Entered into force on 17th May 2005. (Guidance documents to Protocol adopted by decision 1999/1).
- The 2001 Stockholm Convention on Persistent Organic Pollutants (POPs), 186 Parties. Entered into force on 17 May 2004.

The following table shows the dates of signature and ratification of Convention and Protocols for Italy.

Table 1.1 Dates of signature and ratification of the UNECE Convention and Protocols

	SIGNATURE	RATIFICATION
1979 Convention	14/11/1979	15/07/1982
1984 EMEP Protocol	28/09/1984	12/01/1989
1985 Sulphur Protocol	09/07/1985	05/02/1990
1988 NO _x Protocol	01/11/1988	19/05/1992
1991 VOC Protocol	19/11/1991	30/06/1995
1994 Sulphur Protocol	14/06/1994	14/09/1998
1998 Heavy Metals Protocol	24/06/1998	
1998 POPs Protocol	24/06/1998	20/06/2006
1999 Multi-effect Protocol (reviewed in 2012)	01/12/1999	
2001 Stockholm Convention on POPs	23/01/2001	12/07/2022

The following classes of pollutants should be included in the emission inventory:

Main Pollutants

- Sulphur oxides (SO_x), in mass of SO₂;

-
- Nitrous oxides (NO_x), in mass of NO₂;
 - Non-methane volatile organic compounds (NMVOC);
 - Ammonia (NH₃);
 - Carbon monoxide (CO).

Particulate matter

- TSP, total suspended particulate;
- PM10, particulate matter less than 10 microns in diameter;
- PM2.5, particulate matter less than 2.5 microns in diameter;
- Black carbon.

Heavy Metals

- Priority Metals: Lead (Pb), Cadmium (Cd) and Mercury (Hg);
- Other metals: Arsenic (As), Chrome (Cr), Copper (Cu), Nickel (Ni), Selenium (Se) and Zinc (Zn).

Persistent organic pollutants (POPs)

- As specified in Annex II of the POPs Protocol, including Polychlorinated Biphenyls (PCBs);
- As specified in Annex III of the POPs Protocol: Dioxins (Diox), Polycyclic Aromatic Hydrocarbons (PAHs), Hexachlorobenzene (HCB).

1.2 National inventory

As a Party to the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP), Italy has to submit annually data on emissions of air pollutants in order to fulfil obligations, in compliance with the implementation of Protocols under the Convention. Parties are required to report on annual national emissions of SO_x, NO_x, NMVOC, CO and NH₃, and various heavy metals and POPs according to the Guidelines for Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution (UNECE, 2008). The same data are submitted also in the framework of the National Emission Ceiling Directive of the European Union (EU, 2016). Specifically, the submission consists of the national LRTAP emission inventory, communicated through compilation of the Nomenclature For Reporting (NFR), and the Informative Inventory Report (IIR).

The Italian informative inventory report contains information on the national inventory for the year 2022, including descriptions of methods, data sources, QA/QC activities carried out and a trend analysis. The inventory accounts for anthropogenic emissions of the following substances: sulphur oxides (SO_x), nitrogen oxides (NO_x), ammonia (NH₃), non-methane volatile organic compounds (NMVOC), carbon monoxide (CO), total suspended particulate (TSP), particulate matter, particles of size <10 µm, (PM10), particulate matter, particles of size < 2.5µm, (PM2.5), black carbon (BC), lead (Pb), cadmium (Cd), mercury (Hg), arsenic (As), chromium (Cr), copper (Cu), nickel (Ni), selenium (Se), zinc (Zn), polychlorinated biphenyls (PCBs), polycyclic aromatic hydrocarbons (PAH), dioxins (Diox), hexachlorobenzene (HCB). Other pollutants are reported as not estimated; more in details polycyclic aromatic hydrocarbons have not been estimates for each compound for all the sectors and further investigation is planned for the reporting of these emissions.

Detailed information on emission figures of primary pollutants, particulate matter, heavy metals and persistent organic pollutants as well as estimation procedures are provided in order to improve the transparency, consistency, comparability, accuracy and completeness of the inventory provided.

The national inventory is updated annually in order to reflect revisions and improvements in the methodology and the availability of new information. Changes are applied retrospectively to earlier years, which accounts for any difference in previously published data. Total emissions by pollutant from 1990 to 2022 are reported in Table 1.2.

Table 1.2 Emission time series by pollutant

		1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
SO _x	Gg	1,784	1,322	756	411	222	127	112	85	79	88
NO _x	Gg	2,125	1,988	1,516	1,291	952	745	667	596	608	620
NMVOOC	Gg	1,970	2,040	1,616	1,322	1,100	891	871	815	847	823
NH ₃	Gg	529	515	517	481	425	403	387	405	393	351
CO	Gg	6,824	7,118	4,814	3,502	3,076	2,283	2,043	1,861	2,031	1,894
As	Mg	37	28	39	28	17	9	6	5	6	6
Cd	Mg	11	11	10	9	5	4	4	4	4	4
Cr	Mg	95	79	55	62	52	47	44	38	44	44
Cu	Mg	389	440	466	500	460	457	384	312	376	394
Hg	Mg	15	14	15	13	8	7	6	6	6	6
Ni	Mg	116	112	109	114	43	33	30	28	31	34
Pb	Mg	4,302	2,021	991	328	247	230	204	180	214	205
Se	Mg	8	8	9	9	8	8	7	6	7	7
Zn	Mg	971	962	925	999	899	843	818	722	864	795
TSP	Gg	499	499	440	440	483	344	304	305	312	297
PM10	Gg	346	342	297	280	301	231	206	200	212	206
PM2.5	Gg	236	234	202	183	206	165	147	139	151	144
BC	Gg	49	47	43	40	34	24	20	18	19	18
PAH	Mg	90	92	59	64	86	71	65	60	68	63
Dioxin	g ITeq	529	511	434	361	342	310	307	280	325	304
HCB	kg	142	110	33	27	16	16	15	11	13	12
PCB	kg	154	166	157	179	133	114	117	104	122	109

The NFR files and other related documents can be found on website at the following address:

<https://emissioni.sina.isprambiente.it/inventario-nazionale/>.

1.3 Institutional arrangements

The Institute for Environmental Protection and Research (ISPRA) has the overall responsibility for the compilation of the national emission inventory and submissions to CLRTAP. The Institute is also responsible for the communication of pollutants under the NEC directive as well as, jointly with the Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), the development of emission scenarios, as established by the Legislative Decree n. 171 of 21st May 2004. Every four years, from 2017 with reference to 2015 emissions, ISPRA shall provide the disaggregation of the national inventory at provincial level as instituted by the Legislative Decree n. 81 of 30 May 2018. Moreover, ISPRA is the single entity in charge of the development and compilation of the national greenhouse gas emission inventory as indicated by the Legislative Decree n. 51 of 7th March 2008. The Ministry of Environment and energy security (MASE) is responsible for the endorsement and for the communication of the inventory to the Secretariat of the different conventions. The Italian National System currently in place is fully described in the document 'National Greenhouse Gas Inventory System in Italy' (ISPRA, 2018).

A specific unit of the Institute is responsible for the compilation of the Italian Atmospheric Emission Inventory and the Italian Greenhouse Gas Inventory in the framework of both the Convention on Climate Change and the Convention on Long Range Transboundary Air Pollution. The whole inventory is compiled by the Institute; scientific and technical institutions and consultants may help in improving information both on activity data and emission factors of specific activities. All the measures to guarantee and improve the transparency, consistency, comparability, accuracy and completeness of the inventory are undertaken.

ISPRA bears the responsibility for the general administration of the inventory, co-ordinates participation in review processes, publishes and archives the inventory results. Specifically, ISPRA is responsible for all aspects of national inventory preparation, reporting and quality management. Activities include the collection and processing of data from different data sources, the selection of appropriate emissions factors and estimation methods consistent with the EMEP/EEA guidebook, the IPCC 1996 Revised Guidelines, the IPCC Good Practice Guidance and Uncertainty management and the IPCC Good Practice

Guidance for land use, land-use change and forestry, and the IPCC 2006 Guidelines, the compilation of the inventory following the QA/QC procedures, the preparation of the Informative Inventory Report and the reporting through the Nomenclature Reporting Format, the response to review checks, the updating and data storage.

Different institutions are responsible for statistical basic data and data publication, which are primary to ISPRA for carrying out estimates. These institutions are part of the National Statistical System (Sistan), which provides national official statistics, and therefore are asked periodically to update statistics; moreover, the National Statistical System ensures the homogeneity of the methods used for official statistics data through a coordination plan, involving the entire public administration at central, regional and local levels.

The main Sistan products, which are primarily necessary for the inventory compilation, are:

- National Statistical Yearbooks, Monthly Statistical Bulletins, by ISTAT (National Institute of Statistics);
- Annual Report on the Energy and Environment, by ENEA (Agency for New Technologies, Energy and the Environment);
- National Energy Balance (annual), Petrochemical Bulletin (quarterly publication), by MASE (Ministry of Environment);
- Transport Statistics Yearbooks, by MIT (Ministry of Transportation);
- Annual Statistics on Electrical Energy in Italy, by TERNA (National Independent System Operator);
- Annual Report on Waste, by ISPRA;
- National Forestry Inventory, by MASAF (Ministry of Agriculture, Food and Forest Policies).

The national emission inventory itself is a Sistan product. Other information and data sources are used to carry out emission estimates, which are generally referred to in Table 1.3 in the following section 1.5.

1.4 Inventory preparation process

ISPRA has established fruitful cooperation with several governmental and research institutions as well as industrial associations, which helps improving information about some leading categories of the inventory. Specifically, these activities aim at the improvement of provision and collection of basic data and emission factors, through plant-specific data, and exchange of information on scientific research and new sources. Moreover, when in depth investigation is needed and estimates are affected by a high uncertainty, sectoral studies are committed to ad hoc research teams or consultants. ISPRA also coordinates with different national and regional authorities and private institutions for the cross-checking of parameters and estimates, as well as with ad hoc expert panels, in order to improve the completeness and transparency of the inventory.

The main basic data needed for the preparation of the national emission inventory are energy statistics, published by the Ministry of Environment (MASE) in the National Energy Balance (BEN), statistics on industrial and agricultural production, published by the National Institute of Statistics (ISTAT), statistics on transportation, provided by the Ministry of Transportation (MIT), and data supplied directly by the relevant professional associations. Emission factors and methodologies used in the estimation process are consistent with the EMEP/EEA Guidebook, the IPCC Guidelines and Good Practice Guidance as well as supported by national experiences and circumstances.

For the industrial sector, emission data collected through the national Pollutant Release and Transfer Register (Italian PRTR), the Large Combustion Plant (LCP) Directive and in the framework of the European Emissions Trading Scheme have yielded considerable developments in the inventory of the relevant sectors. In fact, these data, even if not always directly used, are considered as a verification of emission estimates and improve national emissions factors as well as activity data figures. In addition, final estimates are checked and verified also in view of annual environmental reports by industries. For large industrial point sources, emissions are registered individually, when communicated, based upon detailed

information such as fuel consumption. Other small plants communicate their emissions which are also considered individually. Emission estimates are drawn up for each sector. Final data are communicated to the UNECE Secretariat filling in the NFR files.

The process of the inventory preparation is carried out annually. In addition to a new year, the entire time series is checked and revised during the annual compilation of the inventory. Recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions and changes due to error corrections. The inventory may also be expanded by including categories not previously estimated if enough information on activity data and suitable emission factors have been identified and collected. Information on the major recalculations is provided in the sectoral chapter of the report.

All the reference material, estimates and calculation sheets, as well as the documentation on scientific papers and the basic data needed for the inventory compilation, are stored and archived at the Institute. After each reporting cycle, all database files, spreadsheets and electronic documents are archived as 'read-only-files' so that the documentation and estimates could be traced back during the new year inventory compilation or a review process.

Technical reports and emission figures are publicly accessible on the web at the address <https://emissioni.sina.isprambiente.it/inventario-nazionale/>.

1.5 Methods and data sources

An outline of methodologies and data sources used in the preparation of the emission inventory for each sector is provided in the following. In Table 1.3 a summary of the activity data and sources used in the inventory compilation is reported.

Table 1.3 Main activity data and sources for the Italian Emission Inventory

Activity data		Source
1 Energy		
1A1 Energy Industries	Fuel use	Energy Balance - MASE Major national electricity producers European Emissions Trading Scheme
1A2 Manufacturing Industries	Fuel use	Energy Balance - MASE Major National Industry Corporation European Emissions Trading Scheme
1A3 Transport	Fuel use Number of vehicles Aircraft landing and take-off	Energy Balance - MASE Statistical Yearbooks - National Statistical System Statistical Yearbooks - Ministry of Transportation Statistical Yearbooks -Italian Civil Aviation Authority (ENAC) Maritime and Airport local authorities
1A4 Residential-public-commercial sector	Fuel use	Energy Balance - MASE
1B Fugitive Emissions from Fuel	Amount of fuel treated	Energy Balance - MASE Statistical Yearbooks - Ministry of Transportation Major National Industry Corporation
2 Industrial Processes	Production data	National Statistical Yearbooks- National Institute of Statistics International Statistical Yearbooks-UN European Emissions Trading Scheme European Pollutant Release and Transfer Register Sectoral Industrial Associations
2D Solvent and Other Product Use	Amount of solvent use	National Environmental Publications - Sectoral Industrial Associations International Statistical Yearbooks-UN
3 Agriculture	Agricultural surfaces Production data Number of animals Fertilizer consumption	Agriculture Statistical Yearbooks - National Institute of Statistics Sectoral Agriculture Associations

	Activity data	Source
4 Land Use, Land Use Change	Forest and soil surfaces	Statistical Yearbooks - National Institute of Statistics
	Amount of biomass	
	Biomass burnt	National and Regional Forestry Inventory - Carabinieri
	Biomass growth	Universities and Research Institutes
5 Waste	Amount of waste	National Waste Cadastre - Institute for Environmental Protection and research

Methodologies are consistent with the EMEP/EEA Emission Inventory Guidebook, Revised 1996 and 2006 IPCC Guidelines, and IPCC Good Practice Guidance (EMEP/CORINAIR, 2007; EMEP/EEA, 2009; EMEP/EEA, 2013; EMEP/EEA, 2016; EMEP/EEA, 2019; EMEP/EEA, 2023; IPCC, 1997; IPCC, 2000; IPCC, 2006); national emission factors are used as well as default emission factors from international guidebooks, when national data are not available. The development of national methodologies is supported by background documents. The most complete document describing national methodologies used in the emission inventory compilation is the National Inventory Report, submitted in the framework of the UN Convention on Climate Change and the Kyoto Protocol (ISPRA, 2024 [a]). Activity data used in emission calculations and their sources are briefly described here below.

In general, for the energy sector, basic statistics for estimating emissions are fuel consumption published in the national Energy Balance by the Ministry of Environment. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. On the other hand, basic information for road transport, maritime and aviation, such as the number of vehicles, harbour statistics and aircraft landing and take-off cycles are provided in statistical yearbooks published both by the National Institute of Statistics and the Ministry of Transportation such as international organizations (i.e. Eurocontrol). Other data are communicated by different category associations. Data from the Italian Emissions Trading Scheme database (ETS) are incorporated into the national inventory whenever the sectoral coverage is complete; in fact, these figures do not always entirely cover the energy categories whereas national statistics, such as the national energy balance and the energy production and consumption statistics, provide the complete basic data needed for the Italian emission inventory. However, the analysis of data from ETS is used to develop country-specific emission factors and check activity data levels. In this context, ISPRA is also responsible for developing, operating and maintaining the national registry under Directive 2003/87/CE as instituted by the Legislative Decree 51 of March 7th 2008; the Institute performs this tasks under the supervision of the national Competent Authority for the implementation of directive 2003/87/CE, amended by Directive 2009/29/EC, jointly established by the Ministry for Environment, Land and Sea and the Ministry for Economic Development.

For the industrial sector, the annual production data are provided by national and international statistical yearbooks. Emission data collected through the national Pollutant Release and Transfer Register (Italian PRTR) are also used in the development of emission estimates or considered as a verification of emission estimates for some specific categories. Italian PRTR data are reported by operators to national and local competent authorities for quality assessment and validation. ISPRA collects facilities' reports and supports the validation activities at national and at local level. ISPRA communicates to the Ministry of Environment and to the European Commission within 30th April of the current year for data referring to the previous year. These data are used for the compilation of the inventory whenever they are complete in terms of sectoral information; in fact, industries communicate figures only if they exceed specific releases thresholds; furthermore, basic data such as fuel consumption are not required and production data are not split by product but reported as an overall value. Anyway, the national PRTR is a good basis for data checks and a way to facilitate contacts with industries which supply, under request, additional information as necessary for carrying out sectoral emission estimates. In addition, final emissions are checked and verified also considering figures reported by industries in their annual environmental reports.

Both for energy and industrial processes, emissions of large industrial point sources are registered individually; communication also takes place in the framework of the European Directive on Large Combustion Plants, based upon detailed information such as fuel consumption. Other small plants communicate their emissions which are also considered individually.

For the other sectors, i.e. for solvents, the amount of solvent use is provided by environmental publications of sector industries and specific associations as well as international statistics. For agriculture, annual production data and number of animals are provided by the National Institute of Statistics and other sectoral associations. For waste, the main activity data are provided by the Institute for Environmental Protection and Research. When basic data are not available proxy variables are considered; unpublished data are used only if supported by personal communication and confidentiality of data is respected. All the material and documents used for the inventory emission estimates are stored at the Institute for Environmental Protection and Research. The inventory is composed by spreadsheets to calculate emission estimates; activity data and emission factors as well as methodologies are referenced to their data sources. A 'reference' database has also been developed to increase the transparency of the inventory; at the moment, it is complete as far as references to greenhouse gas emissions are concerned.

1.6 Key categories

A key category analysis of the Italian inventory is carried out according to the Approach 1 method described in the EMEP/EEA Guidebook (EMEP/EEA, 2023). According to these guidelines, a key category is defined as an emission category that has a significant influence on a country's inventory in terms of the absolute level in emissions. Key categories are those which, when summed together in descending order of magnitude, add up to over 80% of the total emissions. National emissions have been disaggregated into the categories reported in the NFR; details vary according to different pollutants in order to reflect specific national circumstances. Results are reported in the following tables for the year 1990 (Table 1.4) and 2022 (Table 1.5) by pollutant. The trend analysis has also been applied considering 1990 and 2022. The results are reported in Table 1.6.

Table 1.4 Key categories for the Italian Emission Inventory in 1990

	Key categories in 1990												Total (%)	
SO_x	1A1a (43.1%)	1A1b (10.8%)	1A2c (7.2%)	1A3d ii (4.4%)	1A4b i (4.1%)	1B2a iv (3.8%)	1A2gviii (3.7%)	1A2f (3.6%)						80.6
NO_x	1A3b i (27.8%)	1A1a (19.2%)	1A3b iii (16.0%)	1A2f (5.7%)	1A4c ii (4.8%)	1A3d ii (4.5%)	1A3b ii (2.9%)							80.9
NH₃	3Da1 (21.3%)	3Da2a (18.0%)	3B1a (17.6%)	3B1b (16.7%)	3B3 (6.9%)									80.5
NM VOC	1A3b i (21.8%)	2D3d (13.7%)	1A3b iv (8.9%)	1A3b v (6.1%)	2D3a (5.9%)	1A4b i (5.0%)	2D3g (3.9%)	1A4c ii (3.5%)	2D3i (3.5%)	1B2a v (3.0%)	3B1a (2.5%)	3B1b (2.5%)		80.4
CO	1A3b i (60.4%)	1A4b i (11.5%)	1A3b iv (7.2%)	1A4c ii (4.1%)										83.3
PM10	1A4b i (19.5%)	1A1a (10.9%)	2A5b (10.9%)	1A3b i (5.6%)	1A4c ii (4.6%)	1A3b iii (3.9%)	3Dc (3.7%)	1A3b ii (3.0%)	1A2f (2.9%)	2A5a (2.7%)	1A3d ii (2.7%)	1A3b vi (2.6%)	3B4g ii (2.4%)	
	2C1 (2.1%)	2A1 (2.0%)	1A2a (1.8%)											81.3
PM2.5	1A4b i (28.2%)	1A1a (10.7%)	1A3b i (8.2%)	1A4c ii (6.8%)	1A3b iii (5.7%)	1A3b ii (4.4%)	1A3d ii (3.9%)	1A2f (3.3%)	2C1 (2.5%)	1A2a (2.1%)	1A3bvi (2.0%)	1A1b (1.7%)	2A1 (1.6%)	81.0
BC	1A3b i (21.4%)	1A4c ii (18.6%)	1A3b iii (14.0%)	1A3b ii (11.7%)	1A4b i (10.7%)	1A2g vii (4.8%)								81.2
Pb	1A3b i (77.4%)	1A3b iv (5.2%)												82.6
Cd	1A2b (26.0%)	1A2a (19.1%)	2C1 (11.9%)	1A4b i (9.1%)	1A2f (5.2%)	2G (4.5%)	1A4a i (3.4%)	2B10a (3.1%)						82.3
Hg	1B2d (22.2%)	2B10a (18.4%)	2C1 (15.0%)	1A2b (10.6%)	1A2f (8.3%)	1A1a (6.5%)								81.1
PAH	2C1 (53.8%)	1A4b i (35.4%)												89.2
Dioxin	1A4a i (19.6%)	1A2a (15.4%)	1A4b i (13.2%)	2C1 (12.7%)	5C1a (8.1%)	5E (5.9%)	5C1b i (5.9%)							80.7
HCB	3Df (83.4%)													83.4
PCB	2C1 (59.4%)	1A2a (25.6%)												85.1
	1 Energy	2 IPPU - Solvent and product use	3 Agriculture	5 Waste										
	2 IPPU - Industry													

Table 1.5 Key categories for the Italian Emission Inventory in 2022

	Key categories in 2022												Total (%)	
SO_x	1A2f (26.4%)	1B2a iv (16.0%)	1A3d ii (12.4%)	1A1a (7.9%)	1A1b (6.2%)	2B10a (5.9%)	1A4b i (5.8%)							80.5
NO_x	1A3b i (20.1%)	1A3d ii (16.4%)	1A3b iii (13.6%)	1A3b ii (6.5%)	1A4b i (6.0%)	1A2f (5.7%)	1A4a i (4.4%)	1A1a (4.4%)	1A4c ii (3.6%)					80.7
NH₃	3Da2a (18.6%)	3B1b (16.3%)	3B1a (13.9%)	3Da1 (12.5%)	3B3 (8.9%)	3Da2c (3.5%)	3B4g ii (3.4%)	6A (3.4%)						80.5
NMVOC	1A4b i (16.9%)	2D3d (15.0%)	2D3a (11.9%)	2D3g (6.3%)	1A3b v (5.9%)	3B1a (4.6%)	3B1b (4.3%)	2H2 (3.3%)	1A4a i (3.3%)	2D3i (3.3%)	1A3b iv (3.1%)	2D3h (2.2%)		80.0
CO	1A4b i (63.7%)	1A3b i (8.7%)	1A3b iv (4.3%)	1A2a (3.3%)	5C2 (3.2%)									83.2
PM10	1A4b i (44.3%)	2A5a (8.4%)	2A5b (6.4%)	1A3b vi (5.1%)	3Dc (5.0%)	1A3d ii (3.8%)	1A2f (2.5%)	1A3bvii (2.3%)	5E (2.1%)					80.0
PM2.5	1A4b i (62.5%)	1A3d ii (5.5%)	1A3b vi (3.8%)	1A2f (3.1%)	5E (3.0%)	2C1 (2.1%)	1A3b i (1.9%)							81.9
BC	1A4b i (44.7%)	1A3b i (12.3%)	1A3d ii (10.0%)	1A3bvi (7.1%)	1A3b iii (5.3%)	5E (4.4%)								83.8
Pb	1A2f (33.0%)	2C1 (30.8%)	1A3b vi (18.8%)											82.5
Cd	2C1 (22.5%)	1A2f (14.2%)	1A2a (10.5%)	1A4b i (9.1%)	2G (8.5%)	5C2 (7.0%)	1A3b i (6.6%)	1A3bvi (4.2%)						82.5
Hg	2C1 (49.2%)	1A2f (10.2%)	1A1a (8.4%)	1A2a (7.4%)	1A4bi (4.1%)	1B2d (3.6%)								82.7
PAH	1A4b i (78.7%)	2C1 (11.8%)												90.5
Dioxin	1A4b i (31.5%)	2C1 (26.6%)	1A2b (17.5%)	5E (14.4%)										90.1
HCB	1A2b (48.8%)	1A4a i (15.1%)	1A4b i (12.2%)	5C1biii (7.7%)										83.8
PCB	2C1 (71.3%)	1A4b i (14.0%)												85.2
1 Energy		2 IPPU - Solvent and product use				5 Waste								
2 IPPU - Industry		3 Agriculture				6 Other								

Table 1.6 Key categories for the Italian Emission Inventory in trend 1990-2022

	Key categories in trend												Total (%)	
SO_x	1A1a (29.0%)	1A2f (18.8%)	1B2a iv (10.1%)	1A3d ii (6.6%)	1A2c (5.7%)	1A1b (3.8%)	1A4a i (3.7%)	1B2c (2.9%)						80.6
NO_x	1A1a (22.8%)	1A3dii (18.2%)	1A3b i (11.9%)	1A4a i (5.9%)	1A3b ii (5.6%)	1A4b i (5.5%)	1A3b iii (3.7%)	1A2C (3.4%)	1A4c i (3.1%)					80.1
NH₃	3Da1 (28.1%)	3B1a (11.6%)	3Da2c (10.3%)	3B3 (6.4%)	6a (5.2%)	3B4a (5.1%)	1B2d (4.7%)	1A3b i (4.1%)	3B4gii (3.4%)	5B2 (3.1%)				82.0
NMVOC	1A3b i (27.1%)	1A4b i (15.9%)	2D3a (8.0%)	1A3b iv (7.9%)	1A4a i (4.2%)	1A4c ii (3.9%)	2D3g (3.2%)	3B1a (2.8%)	3B1b (2.5%)	2H2 (2.3%)	2D3h (1.8%)	1B2a v (1.7%)		81.4
CO	1A4b i (43.1%)	1A3b i (42.8%)												85.9
PM₁₀	1A4b i (30.5%)	1A1a (13.2%)	2A5b (5.6%)	1A4c ii (5.5%)	1A3b i (5.2%)	2A5a (4.9%)	1A3biii (4.0%)	1A3b ii (3.3%)	1A3b vi (3.1%)	1A2a (2.2%)	1A1b (1.9%)	3Dc (1.7%)		80.8
PM_{2.5}	1A4b i (39.1%)	1A1a (11.9%)	1A4c ii (7.5%)	1A3b i (7.1%)	1A3b iii (5.4%)	1A3b ii (4.5%)	1A2a (2.3%)	1A3b vi (2.1%)						80.1
BC	1A4b i (30.8%)	1A4c ii (16.2%)	1A3b i (8.2%)	1A3b ii (8.1%)	1A3b iii (7.9%)	1A3d ii (6.6%)	1A3b vi (4.5%)							82.3
Pb	1A3b i (42.3%)	1A2f (17.0%)	2C1 (16.0%)	1A3b vi (9.8%)										85.1
Cd	1A2b (28.6%)	2C1 (13.6%)	1A2f (11.6%)	1A2a (11.0%)	1A3b i (6.0%)	5C2 (5.5%)	2G (5.1%)							81.5
Hg	2C1 (34.8%)	1B2d (19.0%)	2B10a (18.8%)	1A2b (9.2%)										81.8
PAH	1A4b i (45.7%)	2C1 (44.3%)												90.0
Dioxin	1A4a i (16.7%)	1A4b i (16.4%)	1A2a (13.6%)	2C1 (12.5%)	1A2b (11.2%)	5E (7.6%)	5C1a (7.2%)							85.2
HCB	3Df (45.2%)	1A2b (25.3%)	1A4a i (7.7%)	1A4b i (6.3%)										84.6
PCB	1A2a (43.7%)	2C1 (25.2%)	1A4b i (16.9%)											85.8

1 Energy	2 IPPU - Solvent and product use	5 Waste
2 IPPU - Industry	3 Agriculture	6 Other

1.7 QA/QC and verification methods

ISPRA has elaborated an inventory QA/QC procedures manual which describes specific QC procedures to be implemented during the inventory development process, facilitates the overall QA procedures to be conducted, as far as possible, on the entire inventory and establishes quality objectives (ISPRA, 2014). Specific QA/QC procedures and different verification activities implemented thoroughly in the current inventory compilation are figured out in the annual QA/QC plans (ISPRA, 2024 [b]). Quality control checks and quality assurance procedures together with some verification activities are applied both to the national inventory as a whole and at sectoral level. Future planned improvements are prepared for each sector by the relevant inventory compiler; each expert identifies areas for sectoral improvement based on his own knowledge and in response to different inventory review processes.

In addition to routine general checks, source specific quality control procedures are applied on a case by case basis, focusing on key categories and on categories where significant methodological and data revision have taken place or new sources. Checklists are compiled annually by the inventory experts and collected by the QA/QC coordinator. These lists are also registered in the 'reference' database. General QC procedures also include data and documentation gathering. Specifically, the inventory analyst for a source category maintains a complete and separate project archive for that source category; the archive includes all the materials needed to develop the inventory for that year and is kept in a transparent manner. Quality assurance procedures regard different verification activities of the inventory. Feedback for the Italian inventory derive from communication of data to different institutions and/or at local level. Emission figures are also subjected to a process of re-examination once the inventory, the inventory related publications and the national inventory reports are posted on website, specifically www.isprambiente.gov.it.

The preparation of environmental reports where data are needed at different aggregation levels or refer to different contexts, such as environmental and economic accountings, is also a check for emission trends. At national level, for instance, emission time series are reported in the Environmental Data Yearbooks published by the Institute, in the Reports on the State of the Environment by the Ministry of Environment and, moreover, figures are communicated to the National Institute of Statistics to be published in the relevant Environmental Statistics Yearbooks as well as used in the framework of the EUROSTAT NAMEA Project. Technical reviews of emission data submitted under the CLRTAP convention are undertaken periodically for each Party. Specifically, an in-depth review of the Italian inventory was carried out in 2010 and 2013 (UNECE, 2010; UNECE, 2013). A summary of the main findings of the last review can be found in the relevant technical report at the address:

http://www.ceip.at/fileadmin/inhalte/emep/pdf/2013_s3/ITALY-Stage3ReviewReport-2013.pdf.

Moreover, under the European National Emission Ceiling Directive (NECD), an in-depth review has been conducted in 2017, 2018, 2019, 2020, 2021, 2022 and in 2023 (EEA, 2017 [a]; EEA 2018; EEA 2019; EEA 2020; EEA 2021; EEA 2022; EEA, 2023). The main resulting findings and how the recommendations were addressed are reported in the following table.

Recommendations from TERT, considering revised estimates (RE) and technical corrections (TC)

Observation	Key Category	NFR, Pollutant(s), Year(s)	Recommendation	RE or TC	Implementation
IT-2D3b-2022-0001	No	2D3b Road paving with asphalt, NMVOC, 1990-2021	For 2D3b Road paving with asphalt and NMVOC for all years, the TERT notes that there is a lack of transparency regarding the methodology used to estimate emissions. This was raised in the 2022 NECD inventory review. In response to a question raised during the review Italy stated that there was still discussion about the emission factor and did not provide a revised estimate. The TERT decided to calculate a technical correction for the 2005 and 2019-2021, which was accepted by Italy. The	TC	Implemented

			estimates demonstrate that the issue is above the threshold of significance. The TERT recommends that Italy include a revised emission calculation in the 2024 submission.		
IT-2A5a-2017-0001	No	2A5a Quarrying and mining of minerals other than coal, PM _{2.5} , PM ₁₀ , 1990-2021	For 2A5a Quarrying and mining of minerals other than coal PM _{2.5} and PM ₁₀ for all years, the TERT identified a potential under-estimate exceeding the threshold of significance. This was also raised during the 2017, 2018, 2019, 2020, 2021 and 2022 NECD inventory review. The TERT noted that Italy had partly implemented the estimate that was the outcome of a recommendation in the 2022 NECD review, but also noted that the reported PM _{2.5} emissions were higher than the PM ₁₀ emissions. In response to a question raised during the review, Italy acknowledged that there was a factor 1000 error. Italy provided a revised estimate for 1990-2021 and stated that it will be included in the 2024 submission. The TERT agreed with the revised estimate provided by Italy. The TERT recommends that Italy include the revised estimate in the next submission.	RE	Implemented
IT-5D1-2022-0001	No	5D1 Domestic wastewater handling, NH ₃ , 1990-2021	For 5D1 Domestic wastewater treatment and NH ₃ for all years, the TERT noted that NH ₃ emissions from dry toilets are reported as 'NA'. This was raised during the 2022 NECD inventory review. In response to a question raised during the 2023 review, Italy provided a revised estimate for years 1990-2021. The TERT agreed with the revised estimate provided by Italy which is consistent with data reported under the UNFCCC. The TERT recommends that Italy include a revised estimate in their next submission and provide more transparency on the activity data in the IIR of their next submission.	RE	Implemented
IT-1A2e-2023-0001	No	1A2e Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco, NH ₃ , 1990-2021	For 1A2e Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco, solid biomass, NH ₃ for all years, the TERT notes that there may be an under-estimate of emissions because the 2019 EEA/EMEP Guidebook includes a methodology for NH ₃ estimation from biomass consumption, but that 'NA' is reported for 2005 and 2019-2020 and only about 0.00002 kt NH ₃ are reported for 2021. In response to a question raised during the review, Italy explained that 99.5% of biomass consumption is biogas and that it will estimate NH ₃ emissions from fuel wood consumption in their next submission. The TERT notes that the issue is below the threshold of significance for a technical correction. The TERT recommends that Italy include NH₃ from solid biomass in the next submission.	NO	Implemented
IT-1A3ai(ii)-2023-0001	No	1A3ai(ii) International aviation cruise (civil) - Memo Item, NMVOC, 2021	For 1A3ai(ii) International aviation cruise (civil) - Memo Item and NMVOC for 2021, the TERT notes that the value "zero" is reported in NFR table. In response to a question raised during the review, Italy explained that an error occurred in the reporting and that emissions are equal to 0.365 kt. The TERT recommends that Italy correct this error for the next submission.	NO	Implemented

IT-1A3di(i)-2023-0003	No	1A3di(i) International maritime navigation - Memo Item, PM _{2.5} , SO ₂ , 2021	For 1A3di(i) International maritime navigation - Memo Item, PM _{2.5} and SO ₂ , and the year 2021, the TERT notes that there is a lack of transparency in the IIR regarding the emissions factors (EF) that are used. Furthermore, the implied EF for PM _{2.5} and SO ₂ have the highest values compared to other Member States for this category and year. In response to a question raised during the review, Italy explained that the consumption of mainly fuel oil is considered as having 3 % sulphur content, and that also PM emissions are high. However, the TERT assumes that information on page 108 of the IIR is correct as “since January 2020, IMO introduced a limit of sulphur content in marine fuel oil equal to 0.5 % (previously equal to 3.5 %) in IMO jurisdiction”. It must be noted that this is a global standard. Moreover, the effect of this regulation can be linked to PM _{2.5} and SO ₂ emissions reductions from 2019 to 2020; there is a higher reduction if compared to activity data reduction (fuel consumption). The TERT recommends that Italy include relevant assumptions (e.g. share of fuel oil and marine diesel oil) in the calculations of their next submission and explanations in the IIR.	NO	Implemented
IT-2D-2023-0001	No	2D Non energy products from fuels and solvent uses, NMVOC, 2017, 2021	For 2D3e Degreasing and NMVOC for 2017 and 2021, the TERT notes that there is a lack of transparency regarding the activity data (AD) and EFs reported in the IIR, and activity data and NMVOC emissions reported in NFR tables. The values of AD reported in the IIR on p.147 for 2017 and 2021 are the same (2,248,707 Mg product) and the value for 2021 differs to that reported in the NFR table (2,602.385 kt), also EF for 2021 in the IIR Table 5.6 p.148 (1,000,000 g/Mg solvents) differ from the IEF (700,000 g/Mg solvents) calculated from values of AD and NMVOC emission reported in NFR tables. In response to a question raised during the review, Italy explained that there was a mistake in the two tables reported in the IIR, and that emission estimates and AD in the NFR table are correct, and as a proof they sent an Excel file. The TERT notes that this issue is related to a lack of transparency regarding activity data and EFs reported in the IIR. The TERT recommends that Italy correct the errors identified in the IIR and ensures consistency between the information presented in the IIR and the data reported in the NFR in the next submission.	No	Implemented
IT-2D3a-2023-0002	Yes	2D3a Domestic solvent use including fungicides, NMVOC, 2016-2021	For 2D3a Domestic solvent use including fungicides and NMVOC for 2016-2021, the TERT notes that there is a lack of transparency regarding activity data and EFs reported in the IIR and activity data and NMVOC emissions reported in NFR tables. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that there has been an error in the tables reported in the IIR and sent corrected activity data set and used EFs. The TERT recommends that Italy correct the errors identified in the IIR and ensures that the information provided in the IIR is correct and	NO	Implemented

			consistent with the information in the NFR of the next submission.		
IT-2D3a-2023-0001	Yes	2D3a Domestic solvent use including fungicides, NMVOC, 2020	For 2D3a Domestic solvent use including fungicides and NMVOC for 2020, the TERT notes that there is a lack of transparency regarding significant recalculations that took place (e.g. -33.2 %; from 126.84 kt to 84.77 kt in 2020 for NMVOC) in the IIR for this key category. In response to a question raised during the review, Italy explained that there has been an update of the activity data for domestic solvent use, sum of cleaning and cosmetics products (the update refers to the data on the production of cosmetics), and by mistake, the description of this recalculation had not been included in the IIR. This does not relate to an over- or under-estimate of emissions. The TERT recommends that Italy identify and explain any recalculations in the IIR of the next submission.	NO	Implemented
IT-2D3d-2023-0002	Yes	2D3d Coating applications, NMVOC, 1990-2021	For 2D3d Coating applications and NMVOC for 1990-2021, the TERT notes a potential over/under-estimate due to differences in reporting activity data and emission factors in the IIR and in NFR tables. The TERT checked NMVOC emission estimations from the AD and EFs reported in the IIR for a few years and got the following differences when data in the IIR with emissions reported in the NFR tables: for 2005 the difference is -23.45 kt, for 2015 there is no difference, for 2016 the difference is -24.86 kt, for 2020 the difference is -4.74 kt, for 2021 the difference is -26.42 kt. In response to a question raised during the review, Italy explained that there has been an error in the tables reported in the IIR and sent corrected activity data sets and used EFs, documenting that the emissions reported in the NFR tables are correct. This does not relate to an over- or under-estimate of emissions. The TERT recommends that Italy correct the errors identified in the IIR and ensures that the information provided in the IIR of the next submission is correct and consistent with the data reported in the NFR tables.	NO	Implemented
T-2D3d-2023-0001	Yes	2D3d Coating applications, NMVOC, 2020	For 2D3d Coating applications and NMVOC for 2020, the TERT notes that there is a lack of transparency regarding not providing an explanation for significant recalculations that took place (e.g. -12.9 %; from 134.87kt to 117.54 kt in 2020 for NMVOC) in the IIR for this key category. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that the recalculations occurred on account of an update of activity data for vehicles production, and by mistake, the description of this recalculation has not been included in the IIR. The TERT recommends that Italy identify all recalculations and provides explanations in the IIR of the next submission.	NO	Implemented
IT-2D3i-2023-0002	Yes	2D3i Other solvent use, NMVOC, 1990-2021	For 2D3i Other solvent use and NMVOC for 1990-2021, the TERT notes that there is a lack of transparency regarding the activity data and EFs reported in the IIR, and activity data reported in the NFR tables. The TERT notes that	NO	Implemented

			Italy reports the same value of 9,000 with no activity units in the NFR table, but in the IIR on p. 147 there are time series of 7 different activities that do not sum to 9,000. The TERT also notes that the numerical value for Application of glues and adhesives, Domestic solvent use (other than paint application), Vehicles dewaxing are exactly the same in the year 2017 and 2021. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that the value 9,000 refers to the activity data of one single category (kt creosote oil consumption) and not to all the categories. Emissions are estimated under 2D3i for application of glues and adhesives, glass wool enduction, fat oil extraction, preservation of wood and vehicle dewaxing. Italy stated its intent to provide information on methods and activity data and EFs in the IIR of the next submission. The TERT recommends that Italy include information on activity data and emission factors and ensures that there are no inconsistencies between data reported in the NFR tables and the IIR of the next submission.		
IT-2G-2023-0001	Yes	2G Other product use, SO ₂ , NO _x , NH ₃ , NMVOC, PM _{2.5} , PAHs, Cd, Pb, PCDD/F, PM ₁₀ , CO, TSP, 1990-2021	For 2G Other product use and all pollutants for all years, the TERT notes that there is a lack of transparency regarding reporting of activity data and emission factors used for the emission estimations. In the NFR tables AD is reported with notation key 'NA' (Not applicable), in the IIR Italy reports that emissions occurring from use of lubricants, tobacco and fireworks but without time series for these activities. In the IIR there is no information on the use of shoes, which is included in 2G. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that these activity data cannot be summed, and that emissions from the use of shoes are included in 2D3i Other solvent use (use of glues and adhesives). Italy provided the TERT with the activity data and emissions for each activity included in 2G and stated that it would be included in the IIR in the future. The TERT recommends that Italy include this information on activity data and emission factors in the IIR of the next submission.	NO	Implemented
IT-3B-2023-0001	No	3B Manure management, NH ₃ , 1990-2021	For 3B Manure management and NH ₃ for all years, the TERT notes that there is a lack of transparency regarding the distribution between animal waste management systems (AWMS) and abatement technologies for dairy cattle and swine, because it is not clear which numbers are used for the period 2014-2021. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy clarified which numbers were used for which years and explained that they have plans to update the information on AWMS and abatement technologies. The TERT recommends that Italy clarify the information about AWMS and abatement technologies in the IIR of the next submission and update data for AWMS and for	NO	Implemented

			the extent of the use of abatement technologies when data are available.		
IT-3B1-2023-0001	No	3B1 Manure management - Cattle, NMVOC, 1990-2021	For 3B1 Manure management – Cattle and NMVOC for all years, the TERT notes that there is a lack of transparency regarding the values used for the calculations, since no information is provided in the IIR. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy provided a spreadsheet with information on the values used for the calculations for 2021. The TERT recommend that Italy include the values for gross feed intake, X_house, Frac_silage, E_NH₃_house, E_NH₃_storage and E_NH₃_application used for the calculations of NMVOC in the IIR of the next submission.	NO	Implemented
IT-3F-2023-0003	No	3F Field burning of agricultural residues, NMVOC, 1990-2021	For 3F Field burning of agricultural residues and NMVOC for all years, the TERT notes that there is a lack of transparency regarding the emission factor used. The IIR refers to the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC, 1997) as the reference for the emission factor, but in IPCC (1997) no emission factors are given for NMVOC. In response to a question raised during the review, Italy explained that the reference to the 1997 IPCC emission factor was given because NMVOC emissions were assumed to be equal to CH ₄ emissions. The TERT recommends that Italy include more information and justification for the current emission factor or update to the information available in the 2019 EMEP/EEA Guidebook in the next submission.	No	Implemented
IT-3F-2023-0002	No	3F Field burning of agricultural residues, SO ₂ , NO _x , NH ₃ , NMVOC, PM _{2.5} , PAHs, Cd, Hg, Pb, PCDD/F, PM ₁₀ , CO, BC, TSP, 1990-2021	For 3F Field burning of agricultural residues and all pollutants for all years, the TERT notes that there is a lack of transparency regarding the units used for the activity data. The activity data are referred to as “Area burned [k ha/yr]”, but the data are not the same as those reported in the CRF. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy explained that the unit of the activity data in NFR were “Area burned [k ha/yr] multiplied by “Biomass available” and they will correct the data in NFR by entering the amount of hectares burned and not the biomass available for combustion or will correct the description of activity data in the NFR. The TERT recommends that Italy enter the amount of hectares burned as activity data in the NFR and include other relevant information related to the methodology in the IIR of the next submission.	NO	Implemented
IT-5A-2022-0002	No	5A Biological treatment of waste - Solid waste disposal on land, PM _{2.5} , PM ₁₀ , 1990-2021	For 5A Biological treatment of waste - Solid waste disposal on land and PM _{2.5} and PM ₁₀ for all years, the TERT noted that Italy does not include all amounts of mineral waste handled. This was raised during the 2022 NECD inventory review and the recommendation was to include all mineral waste in the estimate. The TERT notes that the issue is below the threshold of significance for a technical correction. In response to a question raised during the review, Italy indicated that this issue will be addressed for the 2024 submission. The TERT reiterates	NO	Implemented

			the recommendation that Italy include all mineral waste handled in the estimate of PM emissions from 5A and to include a methodological description in the IIR of the next submission.		
IT-5B1-2022-0001	No	5B1 Biological treatment of waste - Composting, NH ₃ , 1990-2021	For 5B1 Biological treatment of waste - Composting and NH ₃ for all years, the TERT notes that the implied emission factor of 0.024 g/Mg of waste corresponds to the application of the Tier 2 emission factor with an abatement efficiency of 90% applied to the total amount of waste composted. This was raised during the 2022 NECD inventory review and the recommendation was to take into consideration the penetration rate of abatement technologies and improve the associated methodological description. The TERT notes that the issue is below the threshold of significance for a technical correction. In response to the question raised during the review Italy indicated that it will further investigate information about compost plants in the 1990's and, if there is no information available, it will apply EFs without abatement. The TERT reiterates the recommendation that Italy take into consideration the penetration rate of abatement technologies in the estimate and improve the corresponding methodological description in the IIR of the next submission.	NO	Not implemented, more info in the IIR
IT-5C1bi-2022-0002	No	5C1bi Industrial waste incineration, PM _{2.5} , PM ₁₀ , 2010-2020	For 5C1bi Industrial waste incineration and PM _{2.5} and PM ₁₀ for 2010-2020, the TERT noted that the EF indicated in the IIR accounts for an additional abatement efficiency (99%) to the default Tier 1 emission factor proposed in the 2019 EMEP/EEA Guidebook rather than using a higher Tier method or using the Tier 1 method as indicated in the Guidebook. This was raised during the 2022 NECD inventory review. The TERT notes that the issue is below the threshold of significance for a technical correction. In response to a question raised during the review Italy explained that it will assess the possibility of the implementation of a higher Tier methodology but, if not possible, it will implement the Tier 1 emission factor. The TERT reiterates the recommendation that Italy either develop a higher Tier method or strictly apply the Tier 1 method as described in the Guidebook.	NO	Implemented
IT-5C1biii-2023-0002	No	5C1biii Clinical waste incineration, NMVOC, 1990-2021	For 5C1biii Clinical waste incineration and NMVOC for all years, the TERT notes that there may be an over-estimate of emissions because the Tier 1 emission factor recommended for industrial waste incineration (7.4 kg/Mg) is applied instead of the emission factor recommended for clinical waste incineration (0.7 kg/Mg). In response to a question raised during the review, Italy agreed and explained that this will be updated in the next submission. The TERT notes that the issue is below the threshold of significance for a technical correction. The TERT recommends that Italy apply the correct NMVOC emission factor in their next submission.	NO	Implemented

iT-5C1biv-2023-0001	No	5C1biv Sewage sludge incineration, SO ₂ , NO _x , NMVOC, PM _{2.5} , PM ₁₀ , 2020	For 5C1biv Sewage sludge incineration and SO ₂ , PM ₁₀ , NO _x , NMVOC and PM _{2.5} emissions for 2020, the TERT notes that no emissions are reported (- is reported in the NFR table), whereas emissions have been reported for these pollutants for other years. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy stated that the activity data was zero in 2020. The TERT recommends that Italy report the notation key 'NO' for the category in 2020.	NO	Implemented
iT-5C2-2023-0001	No	5C2 Open burning of waste, SO ₂ , NO _x , NMVOC, PM _{2.5} , PM ₁₀ , 1990-2021	For 5C2 Open burning of waste and the pollutants SO ₂ , NO _x , NMVOC, PM _{2.5} , PM ₁₀ emission for all years, the TERT notes that in the IIR (page 183-185) there is a lack of transparency regarding the emission factors applied to estimate emissions. This does not relate to an over- or under-estimate of emissions. In response to a question raised during the review, Italy provided information on the emission factors, some of which are derived from the IPCC methodology, some refer to the EMEP/EEA Guidebook and the particle size distribution to CEPMEIP. Italy stated that they will evaluate the emission factors for the next submission and update the information in the IIR. The TERT recommends that Italy provide the emission factor values and their reference(s) in the IIR of the next submission.	NO	Implemented, more info in the IIR

A bilateral independent review between Italy and Spain was undertaken in the year 2012, with a focus on the revision of emission inventories and projections of both the Parties. With regard to the emission inventory the Italian team revised part of the energy sector of Spain, specifically the public power plants, petroleum refining plants, road transport and off-road categories, whereas the Spanish team revised the Industrial processes and solvent and other product use, and the LULUCF sectors of Italy. Results of these analyses are reported in a technical report. Aim of the review was to carry out a general quality assurance analysis of the inventories in terms of methodologies, EFs and references used, as well as analyzing critical cross cutting issues such as the details of the national energy balances and comparison with international data (EUROSTAT and IEA) and use of plant specific information. In addition, an official independent review of the entire Italian inventory was undertaken by the Aether consultants in 2013. Main findings and recommendations are reported in a final document, and regard mostly the transparency in the NIR, the improvement of QA/QC documentation and some pending issues in the LULUCF sector. These suggestions were considered in the implementation of the following inventories.

Comparisons between national activity data and data from international databases are usually carried out in order to find out the main differences and an explanation to them. Emission intensity indicators among countries (e.g. emissions per capita, industrial emissions per unit of added value, road transport emissions per passenger car, emissions from power generation per kWh of electricity produced, emissions from dairy cows per tonne of milk produced) can also be useful to provide a preliminary check and verification of the order of magnitude of the emissions. Additional comparisons between emission estimates from industrial sectors and those published by the industry itself in the Environmental reports are carried out annually in order to assess the quality and the uncertainty of the estimates. The quality of the inventory has also improved by the organization and participation in sector specific workshops. A specific procedure undertaken for improving the inventory regards the establishment of national expert panels (in particular, in road transport, land use change and forestry and energy sectors) which involve, on a voluntary basis, different institutions, local agencies and industrial associations cooperating for improving activity data and emission factors accuracy.

Furthermore, activities in the framework of the improvement of local inventories are carried out together with local authorities concentrating on the comparison between top down and bottom-up approaches and identifying the main critical issues. In 2021, ISPRA has finalized the provincial inventory at local scale for the years 1990, 1995, 2000, 2005, 2010, 2015 and 2019 applying a top-down approach. Methodologies and results were checked out by regional and local environmental agencies and authorities, and figures are available at ISPRA web address <https://emissioni.sina.isprambiente.it/inventari-locali/>. Methodologies used for spatial disaggregation are described in related publications (ISPRA, 2009; ISPRA, 2022). This work is also relevant to carry out regional scenarios, for the main pollutants, within the Gains Italy project implemented by ENEA and supported by ISPRA and the regional authorities. In addition to these expert panels, ISPRA participates in technical working groups within the National Statistical System. These groups, named "Circoli di qualità" ("Quality Panels"), coordinated by the National Institute of Statistics, are constituted by both producers and users of statistical information with the aim of improving and monitoring statistical information in specific sectors such as transport, industry, agriculture, forest and fishing. These activities should improve the quality and details of basic data, as well as enable a more organized and timely communication.

Other specific activities relating to improvements of the inventory and QA/QC practices regard the progress on management of information collected in the framework of different European obligations, Large Combustion Plant, E-PRTR and Emissions Trading, which is gathered together in an informative system thus highlighting the main discrepancies among data, detecting potential errors and improving the time series consistency. ISPRA collects these data from the industrial facilities and the inventory team manages the information and makes use of it in the preparation of the national inventory. The informative system is based on identification codes to trace back individual point sources in different databases and all the figures are considered in an overall approach and used in the compilation of the inventory. A proper archiving and reporting of the documentation related to the inventory compilation process is also part of the national QA/QC plan. All the material and documents used for the inventory preparation are stored at the Institute for Environmental Protection and Research. Information relating to the planning, preparation, and management of inventory activities are documented and archived. The archive is organized so that any skilled analyst could obtain relevant data sources and spreadsheets, reproduce the inventory and review all decisions about assumptions and methodologies undertaken. A master documentation catalogue is generated for each inventory year and it is possible to track changes in data and methodologies over time. Specifically, the documentation includes:

- electronic copies of each of the draft and final inventory report, electronic copies of the draft and final NFR tables;
- electronic copies of all the final, linked source category spreadsheets for the inventory estimates (including all spreadsheets that feed the emission spreadsheets);
- results of the reviews and, in general, all documentation related to the corresponding inventory year submission.

After each reporting cycle, all database files, spreadsheets and electronic documents are archived as 'read-only' mode.

A 'reference' database is also compiled every year to increase the transparency of the inventory. This database consists of a number of records that references all documentation used during the inventory compilation, for each sector and submission year, the link to the electronically available documents and the place where they are stored as well as internal documentation on QA/QC procedures.

1.8 General uncertainty evaluation

Uncertainty assessment is an important part of compiling an emissions inventory and to assess how uncertainties evolve over time. It is impossible to eliminate uncertainty completely, leading to the conclusion that for every value reported in an inventory there will exist an associated uncertainty. Knowledge of this uncertainty is an integral part of the inventory compilation effort. Uncertainty in emission estimates is a function of the uncertainty of input data i.e. activity or emission factors, used to

compile the inventory. Hence, data collection and uncertainty evaluation are strongly linked and all data contributing to the estimation of emissions should have an associated uncertainty assessment.

Until the 2023 inventory submission, Italy had not systematically assessed the uncertainty of national estimates. Nevertheless, different studies on uncertainty have been carried out (Romano et al., 2004) and a quantitative assessment of the Italian GHG inventory is performed by the Tier 1 method defined in the 2006 IPCC Guidelines (IPCC, 2006) which provides a calculation based on the error propagation equations. Details on the results of the GHG inventory uncertainty figures can be found in the National Inventory Report 2024 (ISPRA, 2024 [a]). The EMEP/EEA Guidebook provides guidance on the quantification of uncertainties in emissions inventories and is aligned with the “2019 Refinement to the 2006 IPCC Guidelines for National GHG Inventories” guidance. Guidelines define two approaches to estimating uncertainties in national greenhouse gas inventories: Approach 1, based on the error propagation equations, and Approach 2, corresponding to the application of Monte Carlo analysis. The Italian inventory team plans to calculate quantitative estimates of the uncertainties using Approach 1 for most estimated pollutants in next years. As above reported, uncertainty in emission estimates is a function of the uncertainty of input data: activity and emission factors. The uncertainty associated with activity data has been borrowed from the National Inventory Report 2024 while values associated with EFs have been taken on the basis of ranges proposed by the 2019 EMEP/EEA Guidebook and the expert judgement of the sectoral experts. In 2023 Italy reported NO_x emissions uncertainty assessment, emission categories are disaggregated into a detailed level and uncertainties are therefore estimated for these categories. In addition, Italy has decided to achieve an in-depth analysis of the most important categories for which punctual data are available starting from the NO_x emissions of the thermoelectric power plants whose evaluation of the uncertainty is reported in 3.2.2.1.

As already mentioned, the Approach 1 has been applied to the Italian inventory to estimate uncertainties for the last submitted year and for the trend. In 2021, the results of Approach 1 for NO_x suggest an uncertainty of 4.1% in the total emissions. The analysis also estimates an uncertainty of 0.9% in the trend. In the Table 1.7 a summary by sector is reported, detailed data have been reported in the Appendix 2.

Table 1.7 Summary by sector of the uncertainty associated with the NO_x inventory.

Sector	Emissions in base year	Emissions in year t	Contribution to total uncertainty by sector in year t	Contribution to total trend uncertainty by sector
	Gg NO _x	Gg NO _x	%	%
Energy	2034.66	550.70	75.05	79.94
IPPU	25.05	5.18	0.02	0.04
Agriculture	61.66	52.69	24.74	19.83
Waste	2.68	2.12	0.19	0.19
Total	2124.05	610.69	100.00	100.00
Uncertainty in total inventory			4.1	0.9

It should be noted that different levels of uncertainty pertain to different pollutants. Estimates of the main pollutants are generally of high level, but PM emissions, especially those of small particle sizes, heavy metal and POP estimates are more uncertain. For this reason, even though not quantified in terms of uncertainty, improvements are planned especially for the specified pollutants.

Nevertheless, since quantitative uncertainty assessments constitute a mean to either provide the inventory users with a quantitative assessment of the inventory quality or to direct the inventory preparation team to priority areas, the Italian Inventory team plans to evaluate uncertainties in next years providing detailed results in the Appendix 2.

In the current submission, uncertainties for SO_x have been evaluated for the last submitted year and in trend. Results have been reported in the next tables.

In 2022, the results of Approach 1 for SO_x suggest an uncertainty of 5.4% in the total emissions. The analysis also estimates an uncertainty of 0.3% in the trend. In the Table 1.8 a summary by sector is reported, detailed data have been reported in the Appendix 2.

Table 1.8 Summary by sector of the uncertainty associated with SO_x inventory.

Sector	Emissions in base year Gg SO _x	Emissions in year t Gg SO _x	Contribution to total uncertainty by sector in year t %	Contribution to total trend uncertainty by sector %
Energy	1714.39	81.19	98.61	99.78
IPPU	68.59	6.75	1.38	0.21
Agriculture	0.08	0.05	0.00	0.00
Waste	0.55	0.16	0.01	0.01
Total	1783.60	88.15	100.00	100.00
Uncertainty in total inventory			5.4	0.3

1.9 General assessment of completeness

The inventory covers all major sources, as well as all main pollutants, included in the UNECE reporting guidelines (UNECE, 2014). NFR sheets are complete as far as the details of basic information are available.

Allocation of emissions is not consistent with the guidelines only where there are no sufficient data available to split the information. For instance, emissions from category 1.A.5.a other stationary are reported and included under category 1.A.4.a i commercial and institutional emission estimates. Mobile commercial and institutional emission estimates (1.A.4.a ii) are included in 1.A.3 sector. PM and HMs emissions from 2.A.3 glass production are included in 1.A.2.f combustion category source as well as those from lead, zinc and copper production are included in 1.A.2.b category. HCB, PCB and Dioxin emissions from aluminium production are included in 1.A.2.b category as well as PM emissions from secondary aluminium production while HCB from iron and steel are included in 1.A.2.a category. NO_x, SO_x and NH₃ from 1.B.1.b, fugitive emissions from solid fuel transformation, are included in the 1.A.2.a category and HMs and POPs from 1.B.2.a iv are included in 1.A.1.b category. PM emissions from storage, handling and transport of mineral products (2.A.5.c) are included under the relevant emission categories.

There are a few emission sources not assessed yet: PAH, dioxin and PCB emissions from 1.A.3.b v gasoline evaporation, dioxin and PCB non-exhaust emissions from 1.A.3.b vi, automobile tyre and brake wear, HMs, PAH, dioxin and PCB non-exhaust emissions from 1.A.3.b vii, road abrasion, NH₃ emissions from 1.A.3.a domestic and international aviation LTO cycle, NH₃ from 1.A.3.e i, pipeline transportation, NO_x from 3.D.a iv, crop residues applied to soils, and 3.D.b, indirect emissions from managed soils. Emission factors for these categories, when available in the Guidebook (EMEP/EEA, 2019), need further assessment for the applicability to the national circumstances. PAH emissions are not detailed in the four indicator compounds for all the categories; we should still estimate them for categories 1.A.1, 1.A.2 stationary, 1.A.3.a, 1.A.3.c, 1.A.3.d ii, 1.A.4.b ii, 1.A.4.c ii, 1.A.4.c iii, 1.A.5.b, 2.C.1, and 2.C.3, because for some categories emission factors are not fully available by compound. Black carbon emissions from the categories reported in the NFR under 2.A.5 a, quarrying and mining of minerals other than coal, and 2.A.5.b, construction and demolition, are not estimated because no information on EF is still available. Emissions of PAH from asphalt blowing, 2.D.3g, are also under further investigation and reported as NE, although according to the relevant industrial association PAH emissions are negligible because all the asphalt blowing plants have abatement filter system of PM and afterburners of gas. Moreover, these plants should respect national environmental legislation not exceeding at the stack more than 0.1mg/Nm³ for total PAH.

Further investigation will be carried out about these source categories and pollutants in order to calculate and improve figures.

2 ANALYSIS OF KEY TRENDS BY POLLUTANT

2.1 MAIN POLLUTANTS

In the following sections, Italian emission series of sulphur oxides, nitrogen oxides, non-methane volatile organic compounds, carbon monoxide and ammonia are presented.

2.1.1 Sulphur dioxide (SO_x)

The national atmospheric emissions of sulphur oxides have significantly decreased in recent years, as occurred in almost all countries of the UNECE. Figure 2.1 and Table 2.1 show the emission trend from 1990 to 2022. Figure 2.1 also illustrates the share of SO_x emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.1 SO_x emissions trend, percentage share by sector and variation 1990-2022.




Table 2.1 SO_x emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	1,000.8	776.4	466.8	187.0	77.1	29.6	13.4	10.9	9.9	13.7
Non industrial combustion plants	82.3	32.5	25.0	22.7	12.1	10.3	10.1	9.8	10.0	9.2
Combustion - Industry	324.1	230.7	119.8	91.4	57.6	35.2	31.7	31.3	27.9	28.5
Production processes	136.0	115.5	38.8	46.4	35.9	24.3	20.0	16.7	19.1	20.9
Solvent and other product use	0.009	0.009	0.032	0.040	0.032	0.017	0.013	0.014	0.028	0.023

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Road transport	129.3	71.6	11.9	2.2	0.4	0.4	0.4	0.3	0.4	0.4
Other mobile sources and machinery	98.3	84.1	84.0	50.6	31.7	23.1	32.1	11.8	7.6	11.6
Waste treatment and disposal	12.9	11.6	10.0	10.7	7.0	4.4	3.8	4.0	4.1	3.9
Agriculture	0.1	0.1	0.1	0.1	0.1	0.1	0.05	0.05	0.05	0.05
Total	1,783.6	1,322.4	756.5	411.0	221.8	127.4	111.5	84.9	79.1	88.1

Figures show a general decline of SO_x emissions during the period, from 1,784 Gg in 1990 to 88 Gg in 2022. The national target of SO_x emissions, set by the National Emission Ceilings Directive at 475 Gg for 2010 (EC, 2001) was reached and continues to be respected after this year revision of the time series. The targets established for 2020 in the framework of the UNECE/CLRTAP Convention and in the framework of 2030 of the revised National Emission Ceiling Directive (EU, 2016), equal for Italy respectively to 65% and 29% of 2005 emissions, has been already reached. The decreasing trend is determined mainly by the reduction in emissions from combustion in energy (-99%) and in industry (-91%), representing in 2022 about 16%, and 32% of the total, respectively. Emissions deriving from non-industrial combustion plants and road transport show a strong decrease too (-89% and -100%, respectively), but these emissions represent about 10% and 0.5% of the total in 2022. Production processes and other mobile sources and machinery also present a significant decreasing trend, showing an influence on the total of 24% and 13% and dropping by about -85% and -88%, respectively. Waste treatment and disposal account for 4% of the national total decreasing of about -70% with respect to 1990. SO_x emissions from agriculture and from solvent and other product use are also estimated but their contribution is irrelevant. An explanation of the sectoral decreasing trend is outlined more in details in the following.

Combustion in energy and transformation industries

The trend of emissions of this sector shows a reduction in the early eighties mainly due to the use of natural gas in place of coal in the energy production and to the implementation of the Directive EEC 75/716 (EC, 1975) which introduces more restrictive constraints in the sulphur content of liquid fuels. During the years 1985-1990, there was an increase of energy consumption that, not sufficiently hampered by additional measures, led to an increase in the emissions of the sector and consequently of total SO_x levels. However, in the nineties, there was an inverse trend due to the introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorisation of plants, and the Ministerial Decree of 12th July 1990, which introduced plant emission limits. Also the European Directive 88/609/EEC (EC, 1988) concerning the limitation of specific pollutants originated from large combustion plants, transposed in Italy by the DM 8th May 1989 (Ministerial Decree of 8th May 1989) gave a contribution to the reduction of emissions in the sector.

Finally, in recent years, a further shift to natural gas in place of fuel oil and a further reduction in the use of coal fuels have contributed to a decrease in emissions.

Non industrial combustion plants

The declining of the emissions occurred mainly as a result of the increase in natural gas and LPG as alternative fuel to coal, diesel and fuel oil for heating; furthermore, several European Directives on the sulphur content in fuels were adopted. In accordance with national legislation, the sulphur content allowed in diesel fuel has decreased from 0.8% in 1980 to 0.2% in 1995 and 0.1% in 2008, while in fuel oil for heating from 3% in 1980 to 0.3% in 1998. Moreover, coal is not more allowed for residential and commercial heating.

Combustion in industry

Emissions from this sector show the same trend of reduction as the category previously analysed, as in the scope of the same rules.

Production processes

Emissions from refineries have been reduced as a result of compliance with the DM 12th July 1990 (Ministerial Decree of 12th July 1990), which introduces limit values. The reduction of emissions from chemical industry is due to the drop off of the sulphuric acid production and to the decrease of emissions in the production of carbon black.

Road transport

The reduction of emissions is mainly due to the introduction of European Directives regulating the sulphur content in liquid fuels.

Other mobile sources and machinery

As regards off roads, emissions mainly derive from maritime transport, which show a decrease due to the introduction of European Directives regulating the sulphur content in fuels.

2.1.2 Nitrogen oxides (NO_x)

The national atmospheric emissions of nitrogen oxides show a decreasing trend in the period 1990-2022, from 2,125 Gg to 624 Gg. Figure 2.2 and Table 2.2 show emission figures from 1990 to 2022. Figure 2.2 also illustrates the share of NO_x emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.2 NO_x emission trend, percentage share by sector and variation 1990-2022.




Table 2.2 NO_x emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	457.4	344.3	172.6	117.9	81.3	52.4	38.7	34.0	35.8	38.9
Non industrial combustion plants	64.2	65.5	64.8	74.9	85.5	86.2	86.1	82.9	85.3	77.6

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
Combustion - Industry	250.6	182.4	154.0	155.5	100.2	60.3	52.4	45.9	49.9	51.8
Production processes	29.9	31.0	9.2	16.0	10.7	9.5	10.5	9.3	10.3	10.4
Solvent and other product use	0.2	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1
Road transport	996.1	1039.5	786.8	629.1	432.2	346.0	282.5	221.8	251.1	251.5
Other mobile sources and machinery	261.5	258.5	262.7	235.2	190.1	137.6	145.2	144.8	120.7	143.4
Waste treatment and disposal	3.0	2.9	2.6	2.7	2.5	2.3	2.2	2.2	2.3	2.2
Agriculture	61.7	64.1	63.3	59.6	49.3	50.7	48.8	55.0	52.6	44.1
Total	2,124.5	1,988.3	1,516.2	1,291.2	952.0	745.2	666.6	596.0	608.1	620.1

Total emissions show a reduction of about 71% from 1990 to 2022, with a marked decrease between 1995 and 2000, especially in the road transport and energy combustion sectors. The target value of emissions, fixed for 2010 by the National Emission Ceilings Directive (EC, 2001) at 990 Gg has been reached and continues to be respected. In 2015, in the framework of the UNECE/CLRTAP Convention, and in particular the Multieffects Protocol, a new target has been established for Italy equal to 60% of 2005 emissions in 2020 and it has been reached. Moreover, the revised National Emission Ceiling Directive (EU, 2016), established a target for Italy equal to 35% of 2005 emissions in 2030.

The main source of emissions is road transport (about 41% in 2022), which shows a reduction of 75% between 1990 and 2022; other mobile sources and machinery in 2022 contributes to the total emissions for 23% and have reduced by 45% from 1990. Combustion in energy and in industry shows a decrease of about -92% and -79%, respectively, having a share on the total of about 6% and 8% in 2022, respectively. Among the sectors concerned, the only one which highlights an increase in emissions is non-industrial combustion plants showing an increase by 21%, accounting for 13% of the total. Details on the sectoral emission trend and respective variation are outlined in the following sections, starting from the early eighties.

Combustion in energy and transformation industries

Emissions from this sector show an upward trend until 1988 due to an increase in energy consumption, not prevented by reduction measures. From 1988 onwards, emissions present a gradual reduction due, mainly, to the introduction of the two regulatory instruments already mentioned for sulphur dioxide: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules for the authorization of facilities and the Ministerial Decree of 12th July 1990, which introduces plant emission limits. The adoption of these regulations, as the Ministerial Decree of 8th May 1989 on large combustion plants, has led to a shift in energy consumption from oil with high sulphur content to oil with lower sulphur content and to natural gas.

In recent years, the conversion to the use of natural gas to replace fuel oil has intensified, thanks to incentives granted for the improvement of energy efficiency. Furthermore, a significant reduction in the use of coal fuels for energy production has been recorded in the last years. These measures, together with those of promoting renewable energy and energy saving, have led to a further reduction of emissions in the sector. In addition, in the last years, more stringent emission limits to the new plants have been established during the authorization process with the aim to prevent air quality issues at local level.

Non industrial combustion plants

The increase in emissions is explained by the growing trend of energy consumption during the period considered. This is because in the last twenty years all the new buildings are equipped with heating system and old buildings have been modernized.

A national survey on energy consumption of households, conducted by the National Institute of Statistics (ISTAT, 2014), has supplied the amount of biomass burned to heating. Estimated values of biomass burnt are about 80% higher than previous estimates reported in the National Energy Balance (MASE, several years) and derived from regional or incomplete surveys. From 2013 this new biomass figures are reported in the National Energy Balance. In 2015 the reconstruction backwards of the time series has been finalised, with the collaboration of ISTAT and GSE (Energy Services Manager), and official data have been communicated to Eurostat. Furthermore, the continuous improvement of appliances for biomass combustion has led to a significant reduction in emissions for different pollutants (PM, COVNM, PAH) but on the other hand an increase of NO_x EFs because of the optimization of combustion process.

Combustion in industry

Emissions from this sector show a decreasing trend, motivated by the same reasons as the energy industry, having undergone the same legislation.

Road transport

The decrease is the result of two opposing trends: an increase in emissions in the early years of the historical series, with a peak in 1992, due to the increase in the fleet and in the total mileage of both passengers and goods transported by road, and a subsequent reduction in emissions. This decrease is, once more, the result of two opposing trends: on one hand, the growth of both the fleet and the mileage, on the other hand the introduction of technologies to reduce vehicle emissions, as the catalytic converter, provided by European Directives, in particular the Directives 91/441/EC (EC, 1991), 94/12/EC (EC, 1994) and 98/69/EC (EC, 1998) and subsequent modifications and integrations.

To encourage the reduction of emissions, different policies have also been implemented, including incentives to renew the public and private fleet and for the purchase of electric vehicles, promotion for the integrated expansion of rail, maritime and urban transport system, and programmes of sustainable mobility. In 2020 pandemic and relevant lockdown measures affected 2020 emission levels especially for this sector.

Other mobile sources and machinery

From 1980 emissions have a slightly rising trend until 1998 and then decrease slightly until arriving in 2017 at lower levels. Emissions in the sector are characterized predominantly by maritime transport, by machinery used in agriculture and industry.

Regarding mobile machinery used in agriculture and industry, these sectors were not governed by any legislation until the Directive 97/68/EC (EC, 1997 [a]), which provides for a reduction in NO_x limits from 1st January 1999, and Directive 2004/26/EC (EC, 2004) which provide further reduction stages with substantial effects from 2011, with a following decreasing trend particularly in recent years.

2.1.3 Ammonia (NH₃)

The national atmospheric emissions of ammonia show a slight decline in the period 1990-2022, from 529 Gg to 351 Gg. Figure 2.3 and Table 2.3 report the emission figures from 1990 to 2022. Figure 2.3 also illustrates the share of NH₃ emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

According to the National Emission Ceilings Directive, the target value of emissions for 2010 amounts to 419 Gg which was achieved. The target established for 2020 in the framework of the UNECE/CLRTAP Convention and relevant protocol is equal for Italy to 95% of 2005 emissions and has been reached. Moreover, the revised national emission Ceiling Directive (EU, 2016) introduced a ceiling equal to 84% of 2005 emissions for 2030.

Figure 2.3 NH₃ emission trend, percentage share by sector and variation 1990-2022.




Table 2.3 NH₃ emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	0.3	0.2	0.2	0.3	0.3	0.3	0.1	0.1	0.1	0.2
Non industrial combustion plants	1.1	1.1	1.0	1.0	1.8	1.6	1.3	1.2	1.4	1.3
Combustion - Industry	0.5	0.6	0.6	4.0	1.6	1.0	1.1	1.0	1.2	1.3
Production processes	0.9	0.5	0.5	0.6	0.6	0.5	0.5	0.5	0.3	0.3
Solvent and other product use	0.4	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3
Geothermal production	8.4	9.0	12.3	13.3	6.0	4.1	1.1	0.9	0.4	0.4
Road transport	0.8	5.3	20.6	15.7	9.5	6.1	5.5	4.3	5.1	5.5
Other mobile sources and machinery	0.03	0.03	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03
Waste treatment and disposal	8.1	9.8	11.6	12.0	10.9	12.2	12.1	12.6	12.6	12.4
Agriculture	499.9	478.2	459.8	423.0	382.1	364.5	353.2	372.5	359.3	317.3
Other	9.2	9.7	10.1	10.7	11.4	11.8	12.1	12.0	12.0	11.9
Total	529.5	514.8	517.1	481.0	424.5	402.6	387.4	405.5	392.7	350.8

In 2022 agriculture is the main source of emissions, with a 90% contribution out of the total NH₃ emissions; from 1990 to 2022 emissions from this sector show a decrease of about 37%. Emissions from road transport show a strong increase, but the share on the total is 1.6%. Emissions from waste treatment and disposal, accounting also only for 3.5% of the total, show an increase of about 53% because of the


increase of NH₃ emissions from anaerobic digestion at biogas facilities. Emissions from non industrial combustion plants show a relevant increase, equal to 22%, but in 2022 the contribution to total emissions is 0.4%. Emissions from combustion in energy and transformation industries as emissions from combustion in industry are not relevant accounting for 0.05% and 0.4% respectively. Emissions from production processes show a reduction of about 61%, but also this contribution is irrelevant as well as emissions from solvent and other product use. Finally, emissions from geothermal production contribute in 2022 for 0.1% of total national emissions.

Specifically, emissions from agriculture have decreased because of the reduction in the number of animals and the trend in agricultural production, and the introduction of abatement technologies due to the implementation of the EU IPPC Directive (EC, 1996). In the last years further emissions reduction result from the implementation of the European Union Rural Development Programs which provide incentives to the introduction of good practice and technologies for the environmental protection and mitigation of GHG and ammonia emissions. Emissions from road transport have increased as a result of the introduction of catalytic converter but during the last years a decrease is observed due to the introduction of more stringent limits in the new vehicles. Emissions from geothermal production have decreased because of the introduction of control and abatement systems in the production plants. Waste sector trend is driven by the increase of biogas facilities due to the incentives for energy production by renewable sources.

2.1.4 Non methane volatile organic compounds (NMVOC)

The national atmospheric emissions of NMVOC show a decreasing trend in the period 1990-2022. Figure 2.4 and Table 2.4 illustrate the emissions values from 1990 to 2022. Figure 2.4 also illustrates the share of NMVOC emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.4 NMVOC emission trend, percentage share by sector and variation 1990-2022.



The total emission trend shows a reduction of about 58% between 1990 and 2022, from 1,970 Gg to 823 Gg.

In the framework of the National Emission Ceilings Directive (EC, 2001), the target value of NMVOC for 2010 fixed at 1,159 Gg was reached. The target established in the framework of the UNECE/CLRTAP Convention for 2020, equal to 65% of 2005 emission level, has been reached taking in account that it

does not include emissions from the agriculture sector apart those from combustion of agriculture residues. In the framework of the European National Emission Ceiling Directive (EU, 2016) a target has been established for Italy equal to 54% of 2005 emissions in 2030.

Solvent and other product use is the main source of emissions, contributing to the total with 40% and showing a decrease of about -44%. The main reductions relate to the road transport sector (-88%), accounting for 11% of the total and to the sector of extraction and distribution of fossil fuels/geothermal energy (-67%), accounting only for 4%. Emissions from agriculture decrease of about 21%, accounting for 15% of the national total. Emissions from other mobile sources and machinery, accounting for 2% of the total, decrease of about 86%. Emissions from non industrial combustion plants show the largest increase (64%), accounting for 20%. Emissions from waste treatment and disposal show a decrease of about 3% while combustion in industry remain the same (0.1%); both these sources account only for about 1%.

Details on the sectoral emission trend and respective variation are outlined in the following sections.

Table 2.4 NMVOC emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	7.6	7.4	6.1	5.6	4.9	3.7	3.7	3.4	3.5	3.6
Non industrial combustion plants	102.6	112.6	112.8	121.1	215.5	188.3	171.9	164.1	180.8	167.8
Combustion - Industry	7.2	8.0	8.4	8.4	6.7	6.7	6.6	6.1	7.4	7.2
Production processes	105.2	94.7	79.0	82.0	67.0	51.4	47.0	48.2	46.6	45.8
Geothermal production	90.9	103.7	56.6	53.9	48.6	37.7	31.2	28.8	31.4	30.3
Solvent and other product use	596.5	550.7	488.2	473.8	391.2	315.1	357.7	320.7	336.5	331.3
Road transport	767.0	879.7	612.2	358.9	179.1	128.6	96.5	86.7	86.6	90.6
Other mobile sources and machinery	133.4	122.0	97.9	73.9	51.7	30.1	25.3	24.3	21.8	18.2
Waste treatment and disposal	8.8	10.3	10.6	11.1	9.9	8.5	8.2	8.6	8.5	8.5
Agriculture	151.2	151.2	144.7	133.2	125.6	121.1	122.9	124.6	123.9	120.0
Total	1,970.4	2,040.4	1,616.4	1,321.9	1,100.2	891.4	871.1	815.5	847.2	823.3

Solvent and other product use

Emissions from this sector stem from numerous activities such as painting (both domestic and industrial), degreasing and dry cleaning, manufacturing and processing of chemicals, other use of solvents and related activities including the use of household products that contain solvents, such as cosmetics, household products and toiletries.

Significant reductions occurred in the nineties by the introduction in the market of products with low solvent content in paints, and the reduction of the total amount of organic solvent used for metal degreasing and in glues and adhesives; furthermore, in many cases, local authorities have imposed abatement equipment in the industrial painting sector and forced the replacement of open loop with closed loop laundry machines even before the EU Directive 99/13/EC (EC, 1999) came into force. In 2020 due to pandemic a strong increase of household products has been registered.

Road transport

The trend of emissions in this sector is characterized by a first stage of reduction in the early eighties, which occurred despite the increase of consumption and mileage because of the gradual adjustment of the national fleet to the European legislation, ECE Regulation 15 and subsequent amendments, introducing stricter emission limits for passenger cars. Subsequently, in the early nineties, an increase in emissions is observed, with a peak in 1992, due to a high increase in gasoline consumption not efficiently opposed by the replacement of the fleet. With the introduction of Directive 91/441/EC (EC, 1991) and following legislation, which provide the use of catalytic device to reduce exhaust and evaporative emissions from cars, NMVOC emissions gradually reduced.

A different explanation of the emission trend pertains to the nineties. In fact, in this period an increase of the fleet and the mileage is observed in Italy, especially for the emergent use of mopeds for urban mobility, which, until 1999, were not subject to any national emission regulation. Thereafter, various measures were introduced in order to facilitate the reduction of NMVOC emissions, including incentives for replacement of both the fleet of passenger cars and of mopeds and motorcycles with low-emission vehicles; incentives were also provided for the use of fuels different from gasoline, such as LPG and natural gas. In addition, funds were allocated for the implementation of urban traffic plans, for the establishment of restricted traffic areas and car-free days, for checks on exhaust pipes of cars, for the implementation of voluntary agreements with manufacturers of mopeds and motorcycles in order to anticipate the timing provided by the European Directive 97/24/EC (EC, 1997 [b]) as regards the placing on the market of mopeds with low emissions.

Non industrial combustion plants

The increasing emission trend is driven by the increase of wood biomass fuel consumption for residential heating. The 2013 consumption value reported in the national energy balance results from a detailed survey conducted by the national institute of statistics in 2014 (ISTAT, 2014) and is much higher than previous estimates. In 2015 the reconstruction backwards of the time series of wood combustion has been finalized with the collaboration of ISTAT and GSE (Energy Services Manager), and official data have been communicated to Eurostat.

Other mobile sources and machinery

The reduction in emissions is explained by the reduction of gasoline consumption in the sector, largely for two-stroke engines used in agriculture and in maritime activities.

Agriculture

NMVOC emissions from agriculture, mainly depend on activity data about different livestock categories. These emissions became significant because of the implementation of the 2016 Guidebook EMEP/EEA emission factors. For the compliance with the established targets these emissions could be subtracted by the total according to the National emission Ceiling Directive (EU, 2016) due to their uncertainty.

As regards the *other sectors*, a decrease in emissions from production processes is observed, mainly in the food industries, in the chemical sector and in the processes in the refineries. The emissions concerning the extraction and distribution of fuels, even in the presence of an increase in quantity treated, have been reduced as a result of the application of the DM 16th May 1996 (Ministerial Decree 16 May 1996), concerning the adoption of devices for the recovery of vapours and of the applications of measures on deposits of gasoline provided by the DM 21st January 2000 (Ministerial Decree 21 January 2000). Emissions from the other sectors are not subject to specific regulations.

2.1.5 Carbon monoxide (CO)

The national CO emissions show a decreasing trend in the period 1990-2022, from 6,824 Gg to 1,894 Gg. The emission figures from 1990 to 2022 are shown in Figure 2.5 and Table 2.5. Figure 2.5 also illustrates the share of CO emissions by category in 1990 and 2022, as well as the total and sectoral variation from 1990 to 2022.

Figure 2.5 CO emission trend, percentage share by sector and variation 1990-2022.




Table 2.5 CO emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	58.9	54.1	54.4	53.9	34.5	39.9	38.6	39.0	34.5	30.9
Non industrial combustion plants	795.1	894.0	913.1	930.4	1,664.9	1,395.4	1,267.7	1,204.6	1,356.8	1,243.9
Combustion - Industry	305.6	410.9	314.6	326.0	233.6	95.3	112.8	88.4	99.9	94.5
Production processes	223.7	139.8	129.2	143.6	105.0	63.6	69.6	56.8	66.0	58.6
Solvent and other product use	5.1	5.1	5.7	5.3	5.1	4.4	4.0	3.9	3.9	3.9
Road transport	4,874.5	5,126.1	3,009.5	1,689.8	757.7	470.3	348.3	261.4	280.3	280.3
Other mobile sources and machinery	480.5	402.5	303.6	264.0	195.2	133.9	126.6	129.6	112.8	111.3
Waste treatment and disposal	61.7	67.7	65.9	72.5	68.5	68.6	65.1	66.6	66.0	60.6
Agriculture	18.5	18.0	18.1	16.2	11.5	11.1	10.2	10.2	10.7	9.9
Total	6,823.6	7,118.3	4,814.0	3,501.7	3,076.0	2,282.5	2,043.0	1,860.5	2,030.9	1,894.0

The decrease in emissions (-72%) is mostly due to the trend observed for the transport sector (including road, railways, air and maritime transport) which shows a total reduction from 1990 to 2022 of about 93%. Specifically, by sector, emissions from road transport and other mobile sources and machinery, accounting in 2022 respectively for 15% and 6% of the total, show a decrease from 1990 to 2022 of about 94% and 77% respectively. On the other hand, emissions from non-industrial combustion plants, representing about 66% of the total in 2022, show a strong increase between 1990 and 2022, equal to 56% due to the increase of wood combustion for residential heating. Figures show a slight decrease in emissions from waste treatment and disposal too (-2%), whose share is 3% of the total and a decrease (-46%) for agriculture which accounts for less than 1% of the total.

2.2 PARTICULATE MATTER

2.2.1 PM10

The national atmospheric emissions of PM10 show a decreasing trend in the period 1990-2022, from 346 Gg to 206 Gg. Figure 2.6 and Table 2.6 illustrate the emission trend from 1990 to 2022. Figure 2.6 also illustrates the share of PM10 emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.6 PM10 emission trend, percentage share by sector and variation 1990-2022.




Table 2.6 PM10 emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	44.8	39.6	18.4	5.9	2.8	1.2	0.7	0.6	0.6	0.7
Non industrial combustion plants	67.8	71.2	68.6	68.6	123.1	106.8	94.0	89.9	101.5	93.2
Combustion - Industry	27.6	25.1	18.6	17.9	12.4	7.7	7.3	6.7	7.6	7.9
Production processes	70.0	70.3	63.5	72.5	76.9	45.3	38.5	41.1	40.6	41.1
Extraction and distribution of fossil fuels	0.7	0.6	0.6	0.8	0.7	0.6	0.3	0.2	0.3	0.4
Solvent and other product use	5.9	6.1	6.6	7.1	6.6	5.3	6.2	6.2	6.5	5.9

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Road transport	59.9	58.9	53.4	48.4	35.6	27.3	21.4	16.8	19.7	20.6
Other mobile sources and machinery	31.6	32.1	30.8	25.3	16.8	10.4	10.6	10.8	8.1	9.3
Waste treatment and disposal	4.8	5.1	5.0	5.1	4.3	4.7	5.4	5.4	5.7	5.7
Agriculture	32.5	33.2	32.0	28.9	21.4	21.5	21.7	21.7	21.6	21.0
Total	345.6	342.2	297.5	280.3	300.6	230.9	206.2	199.6	212.2	205.7

From 1990 to 2022 the trend shows a reduction of about 40%. A considerable amount of emissions is mostly to be attributed to non-industrial combustion plant (45% in 2022) which is increasing its emissions, about 38%, due to the increase of wood combustion for residential heating. Emissions from production processes accounting for 20% of the total in 2022 decrease of about 41% between 1990 and 2022. Agriculture sector, accounting for 10% of total emissions in 2022, reduced its emissions by 36% in 2022 with respect to 1990, due to a reduction in emissions from livestock, which fall by about 47%, and a reduction in emissions from crops, which fall by about 18% due to a reduction in the area of arable land in 2022 compared to 1990. Road transport accounts for 10% of total emissions in 2022 and decrease by 66% due to the introduction of the relevant European Directives controlling and limiting PM emissions at the car exhaust pipe. In 2022 other mobile sources and machinery, accounting for 5% of the total, shows a reduction of about 71% in consideration of the implementation of the relevant European Directives on machinery. Emissions from combustion in industry account for about 4% of the total and decrease by about 72%. The largest decrease (-99%) is observed in emissions deriving from combustion in energy and transformation industries, whose contribution to total emissions is almost irrelevant in 2022 and lower than 1%. The reduction in the energy and industrial sectors is mainly due to the introduction of two regulatory instruments, already mentioned for other pollutants, the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules for the authorization of facilities and the Ministerial Decree of 12th July 1990, which introduces plant emission limits.

2.2.2 PM2.5

The trend of the national atmospheric emissions of PM2.5 is decreasing between 1990 and 2022, with a variation from 236 Gg to 144 Gg. Figure 2.7 and Table 2.7 illustrate the emission trend from 1990 to 2022. Figure 2.7 also illustrates the share of PM2.5 emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.7 PM2.5 emission trend, percentage share by sector and variation 1990-2022.




Table 2.7 PM2.5 emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	30.1	27.8	12.7	3.7	1.8	0.8	0.5	0.4	0.4	0.5
Non industrial combustion plants	66.9	70.5	67.9	67.8	121.8	105.6	92.8	88.8	100.2	92.0
Combustion - Industry	19.9	18.3	14.0	13.6	9.8	6.3	6.1	5.6	6.4	6.5
Production processes	18.2	17.7	15.2	16.6	15.4	10.0	8.7	8.4	9.1	8.8
Extraction and distribution of fossil fuels	0.07	0.06	0.06	0.08	0.07	0.06	0.03	0.02	0.03	0.04
Solvent and other product use	4.7	4.8	5.1	5.4	5.0	4.1	4.6	4.6	4.8	4.4
Road transport	53.7	51.9	45.8	40.1	27.9	19.6	14.5	11.2	12.9	13.4

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Other mobile sources and machinery	31.5	32.0	30.7	25.2	16.8	10.4	10.6	10.8	8.1	9.3
Waste treatment and disposal	4.7	5.0	4.9	5.0	4.2	4.7	5.3	5.4	5.7	5.6
Agriculture	6.1	6.0	5.9	5.2	3.8	3.8	3.8	3.8	3.8	3.6
Total	236.0	234.2	202.3	182.7	206.5	165.4	147.1	139.1	151.3	144.2

In the framework of the revision of the Multieffect protocol of the UNECE/CLRTAP Convention, a target has been established for this pollutant. Italy should reduce in 2020 their PM2.5 emissions by 10% with respect the 2005 emission level and it has been reached. Moreover, in the national emission Ceiling Directive a target has been established for 2030 equal to 60% of 2005 emissions.

Total emissions show a global reduction from 1990 to 2022 of about 39%. Specifically, emissions from road transport, accounting for 9% of total emissions, decrease of about 75%. Emissions from other mobile sources and machinery show a reduction of 71%, accounting in 2022 for 6% of total emissions. Emissions from non-industrial combustion plants and from combustion in industry account for 64% and 5% of the total respectively, but while the former shows an increase of about 38%, the latter decreases by about 67%.

Agriculture sector, accounting for 2 of total emissions in 2022, reduced its emissions by 41% in 2022 respect to 1990. Emissions from waste treatment and disposal, accounting for 4% of the total in 2022, show an increase of about 19%. The largest decrease is observed for combustion in energy and transformation industries (-98%), being the influence on the total in 2022 lower than 1%.

For the explanation of the trends see what already reported for PM10.

2.2.3 Black Carbon (BC)

Black Carbon emissions have been estimated as a fraction of PM_{2.5}. National BC atmospheric emissions are decreasing between 1990 and 2022, with a variation from 49 Gg to 18 Gg. Figure 2.8 and Table 2.8 illustrate the emission trend from 1990 to 2022. Figure 2.8 also illustrates the share of BC emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.8 BC emission trend, percentage share by sector and variation 1990-2022.




Table 2.8 BC emission trend from 1990 to 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Combustion in energy and transformation industries	1.7	1.6	0.8	0.3	0.1	0.04	0.03	0.03	0.02	0.02
Non industrial combustion plants	5.4	5.7	5.6	5.7	10.3	9.1	8.4	8.0	9.1	8.3
Combustion - Industry	0.7	0.6	0.5	0.5	0.4	0.3	0.3	0.2	0.3	0.3
Production processes	0.5	0.4	0.3	0.3	0.2	0.1	0.1	0.1	0.1	0.1
Extraction and distribution of fossil fuels	0.1	0.05	0.05	0.1	0.1	0.1	0.03	0.02	0.02	0.03
Road transport	24.7	23.0	21.0	20.5	15.1	9.7	6.2	4.6	5.0	5.2
Other mobile sources and machinery	14.1	14.5	13.6	10.7	6.0	3.6	3.3	3.2	2.6	2.6

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Waste treatment and disposal	1.4	1.4	1.4	1.5	1.4	1.5	1.6	1.6	1.7	1.6
Agriculture	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	48.6	47.4	43.3	39.8	33.8	24.4	20.0	17.8	18.8	18.2

Total emissions show a global reduction from 1990 to 2022 of about 63%. Specifically, emissions from road transport, accounting for 29% of total emissions, decrease of about 79%. Emissions from other mobile sources and machinery show a reduction of 82%, accounting in 2022 for 14% of total emissions. Emissions from non-industrial combustion plants and from combustion in industry account for 46% and 2% of the total respectively, but while the former shows an increase of about 54%, the latter decreases by about 62%. Industrial processes, accounting for less than 1% in 2022, decrease of 78%. Emissions from waste treatment and disposal, accounting for 9% of the total in 2022, show a decrease of about 16%. The largest decrease is observed for combustion in energy and transformation industries (-99%), being the influence on the total in 2022 less than 1%.

For the explanation of the trends refer to previous paragraph.

2.3 HEAVY METALS (PB, CD, HG)

This section provides an illustration of the most significant developments between 1990 and 2022 of lead, cadmium and mercury emissions.

2.3.1 Lead (Pb)

The national atmospheric emissions of lead show a strong decreasing trend (-95%) between 1990 and 2022, varying from 4,302 Mg to 205 Mg. Figure 2.9 and Table 2.9 illustrate the emission trend from 1990 to 2022. Figure 2.9 also illustrates the share of Pb emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.9 Pb emission trend, percentage share by sector and variation 1990-2022.




Table 2.9 Pb emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	4.0	4.0	3.6	3.8	3.1	2.7	1.7	1.4	1.5	1.9
Non industrial combustion plants	14.5	16.6	22.4	46.3	16.5	15.1	14.1	13.6	14.7	13.7
Combustion - Industry	263.1	234.9	153.6	141.9	104.7	95.4	78.2	68.7	80.5	77.9
Production processes	63.7	68.2	67.3	74.2	69.5	66.1	68.1	61.0	72.6	65.3
Solvent and other product use	2.4	2.4	8.4	10.3	8.4	4.3	3.4	3.7	7.2	6.1
Road transport	3,806.2	1,644.8	719.8	46.0	43.6	44.6	37.3	30.2	36.6	38.4

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Other mobile sources and machinery	142.2	44.2	13.3	1.1	1.1	1.1	1.4	0.8	0.8	1.2
Waste treatment and disposal	5.9	5.5	2.8	4.0	0.3	0.2	0.2	0.2	0.2	0.2
Agriculture	0.02	0.02	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01
Total	4,302.1	2,020.6	991.2	327.7	247.2	229.5	204.3	179.6	214.1	204.7

In 2022 emissions from combustion in industry have the most significant impact on the total (38%) and show a reduction of about 70%; this reduction is to be attributed primarily to processes with contact, which account for almost the total share of the sector. Emissions from production processes and, in particular, from processes in iron and steel industries and collieries, are not very variable increasing by about 3% and representing 32% of the total. Emissions from non-industrial combustion plants show a 5% decrease and represent, in 2022, 7% of the total. As to emissions from transport activities, because of changes occurred in the legislation regarding fuels, trends show a sharp reduction in emissions from 2002 onwards.

2.3.2 Cadmium (Cd)

The national atmospheric emissions of cadmium show a decreasing trend. Figure 2.10 and Table 2.10 illustrate the emission trend from 1990 to 2022. Figure 2.10 also illustrates the share of Cd emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.10 Cd emission trend, percentage share by sector and variation 1990-2022.




Table 2.10 Cd emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	0.2	0.2	0.2	0.2	0.1	0.1	0.1	0.1	0.1	0.1
Non industrial combustion plants	1.5	1.2	1.7	2.6	0.7	0.5	0.5	0.5	0.5	0.5
Combustion - Industry	5.6	5.6	5.0	2.5	1.6	1.1	1.2	1.2	1.3	1.2
Production processes	2.0	1.8	1.4	1.5	1.4	1.1	1.1	1.0	1.2	1.0
Solvent and other product use	0.5	0.5	0.6	0.5	0.5	0.4	0.4	0.3	0.3	0.3
Road transport	0.5	0.6	0.6	0.7	0.6	0.6	0.5	0.4	0.5	0.5
Other mobile sources and machinery	0.03	0.03	0.03	0.03	0.03	0.02	0.03	0.03	0.02	0.03
Waste treatment and disposal	0.6	0.6	0.5	0.5	0.3	0.4	0.3	0.3	0.3	0.3
Agriculture	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	11.1	10.6	10.1	8.6	5.2	4.4	4.2	3.9	4.4	4.1

Emissions show a global reduction of 63% between 1990 and 2022, from 11.1 Mg to 4.1 Mg, mainly driven by the reduction of emissions in the non-ferrous metal industry, with the installation of the relevant abatement technologies and the drop of production. Among the most significant variations, emissions from combustion in industry and from non-industrial combustion plants represent 29% and 11% of the total respectively, showing a decrease of 79% and 69% respectively. Emissions from production processes decrease by about 48% and represent 26% of the total. Emissions from waste treatment and disposal (i.e. waste incineration), accounting for 8% of the total, register a reduction of about 48% while emissions from road transport, accounting for 13% of the total levels, increased by 4% and emissions from stubble burning in agriculture account for 2% of the total and decrease of about 28%.

2.3.3 Mercury (Hg)

The national atmospheric emissions of mercury show a reduction trend in the period 1990-2022. Figure 2.11 and Table 2.11 illustrate the emission trend from 1990 to 2022. Figure 2.11 also illustrates the share of Hg emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Emission trend shows a global reduction of about 64% from 1990 to 2022, varying from 15.3 Mg to 5.6 Mg. The general trend is driven by reduction of emissions in non-ferrous production industry as well as in cement production industry, with the installation of the relevant abatement technologies. The main variations concern: emissions from combustion in industry, accounting for 23% and decreasing by 70%; emissions from production process - processes in iron and steel industries and collieries, representing 50% of the total and increasing by 12%; emissions from non-industrial combustion plants which represent 8% of the total and decrease by 27%. Emissions deriving from combustion in energy and transformation industries, accounting for 10%, show a 49% reduction. Emissions from production process - processes in inorganic chemical industries, not contributing to the total in 2022, show a reduction equal to 100% totally due to the technological changes to produce chlorine. Emissions from road transport account for 3% and are quite stable (-0.2%). Emissions from waste treatment and disposal contributing to the total for 2% and agriculture, contributing to the total only for less than 1%, show a large reduction, equal respectively to -70% and -31%. Emissions from geothermal production account for 4% of the national total and shows a reduction of 94% with respect to 1990 due to the introduction of control and abatement system at the production plants.

Figure 2.11 Hg emission trend, percentage share by sector and variation 1990-2022.




Table 2.11 Hg emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	1.1	1.1	1.1	1.1	0.9	0.7	0.5	0.5	0.5	0.6
Non industrial combustion plants	0.6	0.7	1.0	2.0	0.5	0.5	0.5	0.5	0.5	0.4
Combustion - Industry	4.2	3.9	3.4	3.4	2.5	2.1	1.7	1.6	1.7	1.3
Production processes	5.5	4.4	3.6	3.4	2.9	2.7	2.9	2.6	3.1	2.8
Geothermal production	3.4	3.6	5.0	2.1	1.3	1.0	0.4	0.3	0.2	0.2
Road transport	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2
Waste treatment and disposal	0.3	0.3	0.2	0.3	0.2	0.1	0.1	0.1	0.1	0.1
Agriculture	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Total	15.3	14.4	14.6	12.5	8.5	7.2	6.4	5.8	6.3	5.6

2.4 PERSISTENT ORGANIC POLLUTANTS (POPs)

In this section, the most significant peculiarities of polycyclic aromatic hydrocarbons and dioxins such as HCB and PCB, occurred between 1990 and 2022, will be presented.

2.4.1 Polycyclic aromatic hydrocarbons (PAH)

The national atmospheric emissions of polycyclic aromatic hydrocarbons decreased from 90 Mg to 63 Mg between 1990 and 2022. Figure 2.12 and Table 2.12 illustrate the emission trend from 1990 to 2022. Figure 2.12 also illustrates the share of PAH emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.12 PAH emission trend, percentage share by sector and variation 1990-2022.




Table 2.12 PAH emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	0.9	1.0	0.7	0.5	0.4	0.5	0.5	0.5	0.4	0.4
Non industrial combustion plants	31.9	35.2	35.7	38.9	67.3	55.9	51.2	48.6	54.8	50.4
Combustion - Industry	4.5	4.6	2.2	2.4	0.4	0.5	0.5	0.4	0.5	0.5
Production processes	48.4	47.4	16.8	17.7	14.1	9.3	9.0	7.1	8.5	7.4
Solvent and other product use	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02
Road transport	1.9	2.0	2.1	2.6	2.7	2.6	2.6	2.1	2.5	2.6
Other mobile sources and machinery	0.3	0.3	0.3	0.4	0.3	0.3	0.3	0.3	0.3	0.3

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Waste treatment and disposal	1.3	1.2	1.2	1.2	1.0	1.0	1.0	1.0	1.0	1.0
Agriculture	0.4	0.4	0.4	0.4	0.3	0.3	0.3	0.3	0.3	0.3
Total	89.7	92.1	59.4	64.0	86.5	70.5	65.4	60.3	68.4	62.8

Between 1990 and 2022, total emissions show a decrease of about 30%. Among the most significant changes, non-industrial combustion plants, prevalently residential plants, account for 80% of the total in 2022 and show a strong increase (about 58%) due to the increase in wood consumption for heating. Emissions from production processes, mainly processes in iron and steel industries, account for 12% of the total and show a decrease of 85% due to the adoption of best abatement technologies for the coke production; emissions from waste treatment and disposal, mainly open burning of agricultural wastes except stubble burning, account for 2% of the total and show a decrease of 24%. Emissions from road transport, accounting for 4% in 2022, show an increase of about 34%. The share of other subsectors is lower than 1%.

2.4.2 Dioxins

The national atmospheric emissions of dioxins show a decreasing trend between 1990 and 2022, with values varying from 529 g I Teq to 304 g I Teq. Figure 2.13 and Table 2.13 illustrate the emission trend from 1990 to 2022. Figure 2.13 also illustrates the share of dioxin emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.13 Dioxin emission trend, percentage share by sector and variation 1990-2022.




Table 2.13 Dioxin emission trend from 1990 to 2022 (g I Teq)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
g I Teq										
Combustion in energy and transformation industries	24.9	27.9	20.9	14.0	8.2	6.2	4.2	4.0	4.0	5.0
Non industrial combustion plants	173.8	164.6	151.0	87.0	134.6	112.4	101.7	96.4	108.5	99.4
Combustion - Industry	116.7	120.9	112.0	118.7	65.3	58.2	57.0	47.1	61.8	59.3
Production processes	67.2	71.7	70.7	78.6	76.2	76.8	84.5	75.7	90.8	80.9
Solvent and other product use	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Road transport	10.7	13.6	18.2	21.4	20.6	14.3	9.4	6.9	7.4	7.9
Other mobile sources and machinery	1.3	1.4	1.3	1.5	1.2	1.0	1.1	1.1	1.0	1.0
Waste treatment and disposal	134.3	110.7	59.9	40.1	35.7	41.1	48.8	48.8	51.7	50.3
Agriculture	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total	529.0	511.0	434.1	361.4	341.7	310.1	306.7	280.0	325.4	304.0

The general trend shows a decrease from 1990 to 2022 equal to 43%, with a noticeable decline between 1995 and 2004 and between 2008 and 2011 because of the implementation of abatement system in the largest Italian integrated iron and steel plant (steel production > 80% with respect to national production from integrated plants):

- Double filtering system ESP (Electrostatic Precipitator) + MEEP (Moving Electrode Electrostatic Precipitator);
- Reduction of the chlorine amount in the charge;
- Injections of urea (able to form stable compounds with metals that catalyse the formation of dioxins).

The most considerable reductions, between 1990 and 2022, are observed in waste treatment and disposal, combustion in energy and transformation industries and combustion in industry, (-63%, -80% and -49%, respectively). Specifically, the reduction is principally due to the cut of emissions from the combustion of municipal waste both with energy recovery, reported under the non-industrial sector, and without recovery, reported under the waste sector due to the introduction of regulations establishing more stringent limits of dioxin emissions from stacks. The waste sector includes accidental fires of vehicles and buildings which account in 2022 for 14% of total national emissions.

In 2022, the subsectors which have contributed most to total emissions are non-industrial combustion plants, production processes and combustion in industry accounting for 33%, 27% and 20% of the total respectively. In particular emissions from production processes show an increase of 20% in the period 1990-2022 due to the increase of the iron and steel production in electric arc furnaces.

2.4.3 Hexachlorobenzene (HCB)

The national atmospheric emissions of hexachlorobenzene show a decreasing trend in the period 1990-2022, varying from 142 kg to 12 kg due to the decrease of the use of pesticide in agriculture. Figure 2.14 and Table 2.14 illustrate the emission trend from 1990 to 2022. Figure 2.14 also illustrates the share of HCB emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.14 HCB emission trend, percentage share by sector and variation 1990-2022.




Table 2.14 HCB emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	0.0002	0.0002	0.0002	0.0004	0.001	0.001	0.001	0.001	0.001	0.001
Non industrial combustion plants	0.002	0.003	0.006	0.006	0.004	0.003	0.003	0.003	0.004	0.003
Combustion - Industry	0.008	0.008	0.009	0.009	0.007	0.007	0.007	0.005	0.007	0.007
Road transport	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000	0.000
Other mobile sources and machinery	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Waste treatment and disposal	0.013	0.014	0.010	0.008	0.000	0.001	0.001	0.001	0.001	0.001
Agriculture	0.119	0.084	0.007	0.002	0.002	0.003	0.002	0.000	0.000	0.000
Total	0.142	0.110	0.033	0.027	0.016	0.016	0.015	0.011	0.013	0.012

The use of pesticide in agriculture category is the main driver for the decreasing trend of the HCB national emissions, emissions from this category show 100% decrease between 1990 and 2022. The second sector contributing to the general trend is waste treatment and disposal, in particular waste incineration - sludge incineration. Specifically, the considerable increase of the amount of sludge burnt at a specific incinerator is the reason of the peaks observed in 2001-2003 (incineration with energy recovery). The other relevant sectors are combustion in industry and non-industrial combustion plants accounting for 54% and 27% respectively. Emissions from combustion in energy and transformation industry and emissions from non-industrial combustion plants show an increase of 187% and 82% respectively between 1990 and 2022. In

the same years for emissions from waste treatment and disposal a decrease of 92% must be noted while emissions from combustion in industry show a decrease of 14%.

2.4.4 Polychlorinated biphenyl (PCB)

The national atmospheric emissions of polychlorinated biphenyl show a decreasing trend in the period 1990-2022, about -29%, from 154 kg to 109 kg. Figure 2.15 and Table 2.15 illustrate the emission trend from 1990 to 2022. Figure 2.15 also illustrates the share of PCB emissions by category in 1990 and 2022 as well as the total and sectoral variation from 1990 to 2022.

Figure 2.15 PCB emission trend, percentage share by sector and variation 1990-2022.




Table 2.15 PCB emission trend from 1990 to 2022 (Mg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mg										
Combustion in energy and transformation industries	0.0000	0.0000	0.0000	0.0001	0.0002	0.0004	0.0004	0.0004	0.0004	0.0003
Non industrial combustion plants	0.015	0.018	0.025	0.036	0.022	0.019	0.019	0.018	0.020	0.018
Combustion - Industry	0.042	0.043	0.033	0.035	0.017	0.014	0.013	0.010	0.012	0.011
Production processes	0.092	0.100	0.096	0.106	0.093	0.079	0.083	0.073	0.088	0.078
Road transport	0.000003	0.000003	0.000004	0.000005	0.000004	0.000003	0.000002	0.000001	0.000001	0.000002
Other mobile sources and machinery	0.0004	0.0004	0.0004	0.0005	0.0004	0.0003	0.0003	0.0003	0.0003	0.0003

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Mg									
Waste treatment and disposal	0.005	0.005	0.002	0.002	0.001	0.001	0.001	0.001	0.001	0.001
Total	0.154	0.166	0.157	0.179	0.133	0.114	0.117	0.104	0.122	0.109

Among the most significant variations, emissions from combustion in industry and from production processes represent 10% and 71% of the total respectively, showing the former a decrease of -74% and the latter of 15%. The noticeable decline between 2008 and 2009 was due to the implementation of abatement systems in the largest Italian steel plant. The other relevant sector is non industrial combustion plants accounting for 17% and relevantly increasing (26%) between 1990 and 2022. The share of other sectors is about or lower than 1%.

3 ENERGY (NFR SECTOR 1)

3.1 OVERVIEW OF THE SECTOR

For the pollutants and sources discussed in this section, emissions result from the combustion of fuel. All the pollutants reported under the UNECE/CLRTAP are estimated. Stationary and mobile categories are covered for:

- Electricity production (power plants and Industrial producers);
- Refineries (Combustion);
- Iron and steel industries (Combustion)
- Chemical and petrochemical industries (Combustion);
- Construction industries (roof tiles, bricks);
- Other industries (metal works factories, food, textiles, others);
- Road Transport;
- Coastal Shipping;
- Railways;
- Aircraft;
- Domestic heating;
- Commercial heating;
- Public Service;
- Fishing and Agriculture.

Fugitive emissions are also reported under the energy sector as well as emissions from geothermal production.

The national emission inventory is prepared using energy consumption information available from national statistics and an estimate of the actual use of the fuels. The latter information is available at sectoral level in a different number of publications and different details, such as fuel consumption, distance travelled or some other statistical data related to emissions. For most of the combustion source categories, emissions are estimated from fuel consumption data reported in the National Energy Balance (BEN) as supplied by the Ministry for the Environment (MASE, several years (a)) and reported to the international energy organization, and from emission factors appropriate to the type of combustion and the pollutant. Currently, a review process of the national energy balance is underway with Eurostat. In a first step, those responsible for the National Energy Balance implemented methodological changes for 2022 and subsequently applied these changes to 2021 to make it consistent. The review process, carried out in agreement with the relevant international institutions, regards definitions of "main activity producer" and "autoproducer" and the consequent adoption by Italy of these definitions in place of those previously used and linked to Italian legislation. This change in methodology has caused a fairly evident break in the series for some fuels and it is necessary further work to improve the time series consistency.

The estimate from fuel consumption emission factors refers to stationary combustion in boilers and heaters. The other categories are estimated by more complex methods discussed in the relevant sections. The fuel consumption of "Other industries" is estimated so that the total fuel consumption of these sources is consistent with the national energy balance.

Electricity generation by companies primarily for their own use is auto-generation, and the relevant emissions should be reported under the industry concerned. However, national energy statistics report emissions from electricity generation as a separate category. The Italian inventory makes an overall calculation and then attempts to report as far as possible according to the guidelines:

- auto-generators are reported in the relevant industrial sectors of section "1.A.2 Manufacturing Industries and Construction";
- refineries auto-generation is included in section 1.A.1b;
- iron and steel auto-generation is included in section 1.A.1c;
- incinerators auto-generation of energy and heat is included in section 1.A.4a.

These reports are based on estimates of fuel used for steam generation connected with electricity production supplied by the National Independent System Operator (TERNA, several years).

Emissions from the energy production plants in integrated iron and steel plants and emissions from coke ovens are included in 1.A.1c category. Emissions from waste incineration facilities with energy recovery are reported under category 1.A.4a i (Combustion activity, commercial/institutional sector), whereas emissions from other types of waste incineration facilities are reported under category 5.C (Waste incineration). In particular, for 2022, almost 99% of the total amount of urban waste incinerated is treated in plants with energy recovery system. The energy recovered by these plants is mainly used for district heating of commercial buildings or used to satisfy the internal energy demand of the plants. Different emission factors for municipal, industrial and oils, hospital waste, and sewage sludge are applied, as reported in the waste chapter. Waste amount is then converted in energy content applying the relevant factor as resulting from data provided by TERNA, which in 2022 is equal to 11.3 GJ/t of waste.

Landfill gas is generally recovered and used for heating and power in commercial facilities, the resulting emissions are reported under 1.A.4.a. Biogas recovered from the anaerobic digester of animal waste is used for utilities in the agriculture sector and relative emissions are reported under 1.A.4.c.

Under 1.A.2 g vii industrial off road machinery are reported; the methodology used to estimate emissions from a range of portable or mobile equipment powered by reciprocating diesel engines is summarized. Industrial off-road include construction equipment such as bulldozers, loaders, graders, scrapers, rollers and excavators and other industrial machines as portable generators, compressors and cement mixers. Estimates are calculated taking in account especially the population of the different classes, annual usage, average power rating, load factor and technology distribution (EURO) according to the Guidebook (EMEP/EEA, 2019). COPERT II has been used for years 1994 and 1995 to estimate emissions and average emission factors for vehicles and diesel fuel consumption. Population data have been estimated on the basis on a survey of machinery sales. Machinery lifetime was estimated on the European averages reported in EMEP/CORINAIR, 2007, the annual usage data were taken either from industry or published data by EEA. The emission factors used came from EMEP/EEA and COPERT. The load factors were taken from COPERT. It was possible to calculate fuel consumptions for each class based on fuel consumption factors given in EMEP/CORINAIR, 2007. Comparison with known fuel consumption for certain groups of classes suggested that the population method overestimated fuel consumption by factors of 1.2 -1.5 for industrial vehicles. Time series is reconstructed in relation to the diesel fuel use in industry reported in the national energy balance as gasoil final consumption. Emission factors for NO_x, CO, NMVOC and PM have been updated taking into account the reduction factors established in the European Directive 97/68/EC, the timing of application of the new limits and the tax of penetration of the new industrial vehicles in the total fleet. Emission reduction factor reported in the European Directive 2004/26/EC Directive have been applied and introduced in the emission estimates.

The emission factors used are based on national sources, or else on values specified in the EMEP/EEA guidebook and/or IPCC guidelines which are appropriate for Italy. Emission factors used for energy and manufacturing industries and non-industrial combustion, specifically categories 1A1, 1A2, 1A4, and their references are available on the ISPRA website at <https://emissioni.sina.isprambiente.it/inventario-nazionale/> as well as emission factors for road transport (1A3b) .

In 2022 the energy sector accounts for more than 50% of total emissions for all the estimated pollutants, except for NMVOC, which accounts for 39%, Hg for 49%, PCB for 28% and ammonia for 3%. In particular, emissions from the energy sector are 93% of CO, 92% of SO_x and NO_x, 91% of HCB, 90% of BC, 86% of PAH and 85% of PM2.5 national total emissions.

In 2022, the following categories are key categories for different pollutants: public electricity and heat production (1A1a), petroleum refining (1A1b), stationary combustion in iron and steel industries (1A2a), stationary combustion in non-ferrous metal industries (1A2b), stationary combustion in non-metallic mineral industries (1A2f), road transport categories (1A3b), national navigation (1A3d ii), stationary combustion plants in commercial/institutional (1A4a i) and residential (1A4b i), off-road vehicles in agriculture, forestry and fishing (1A4c ii), fugitive emissions from refining and storage (1B2a iv) and other fugitive emissions from energy production (1B2d).

The same categories are key categories for 1990 and for the trend analysis. In addition, for 1990, stationary combustion in chemical industries (1A2c) for SO_x, mobile combustion in manufacturing industries and construction (1A2g vii) for BC, stationary combustion in other industries (1A2g viii) for SO_x, fugitive emissions from distribution of oil products (1B2a v) for NMVOC emissions are also key categories.

The following sections present an outline of the main key categories in the energy sector. Table 3.1 highlights the key categories identified in the sector.

The energy sector is the main source of emissions in Italy with a share of more than 80% for different pollutants; specifically, for the main pollutants, in 2022 the sector accounts for:

- 93% in national total CO emissions;
- 92% in national total SO_x emissions;
- 92% in national total NO_x emissions;
- 91% in national total HCB emissions;
- 90% in national total BC emissions;
- 86% in national total PAH emissions;
- 85% in national total PM_{2.5} emissions.

Moreover, the sector is also an important source for heavy metals; specifically in 2022, energy sector is responsible for 58% of total Cd emissions, 49% for Hg and 65% for lead emissions.

There are limited differences as compared to the sectoral share in 1990, except for heavy metals, in particular lead whose contribution in 1990 was 98% of total emissions, 33% higher than in 2022, for PAH whose contribution in 1990 was 44%, 42% lower than in 2022 and for HCB emissions with 7% in 1990 against 91% in 2022.

One of the most important sources of emissions in the sector and key category, in 2022, is represented by road transport (1A3b), at least for the main pollutants: NO_x (40.6%), BC (28.6%), Pb (18.8%), CO (14.8%), Cd (13.0%), NMVOC (11.0%) and particulate matter (PM₁₀ 10.0%, PM_{2.5} 9.3%). There has been a strong reduction in lead emissions from 1990 to 2022 in road transport due to replacement of lead gasoline. An in depth analysis of the road transport category and its emission trends is reported in paragraph 3.5.

Manufacturing industries and construction (1A2) is a main source of heavy metals and POPs, accounting for about 38.1% of lead total emissions, 29.2% for cadmium, 22.7% for mercury, 54.0% for HCB and 19.5% for dioxin. The sector is key category also for PM₁₀ and PM_{2.5} (3.8% and 4.5%) as well as SO_x, NO_x and CO, about 32.3%, 8.4% and 5.0% of total emissions. The main sectors are iron and steel sector, which is key for CO, Cd and Hg, the non-ferrous metal sector, key for HCB and Dioxin, and non-metallic mineral sector that is key category for SO_x, NO_x, PM₁₀, PM_{2.5}, Pb, Cd and Hg.

Public electricity and heat production (1A1a) is a key source of SO_x emissions in 2022 with a share of 7.9%, Hg (8.4%) and NO_x emissions (4.4%) while SO_x emissions from petroleum refining (1A1b) are key category in 2022.

National navigation (1A3d ii) is key category for SO_x (12.4%), NO_x (16.4%), PM₁₀ (3.8%), PM_{2.5} (5.5%), and BC (10.0%). The weight of this category on the total emissions has increased for SO_x and NO_x during the period because of a sectoral delay in the introduction of relevant normative to reduce air emissions.

A sector increasing its level of emissions is the non-industrial combustion (1A4): NO_x and NMVOC, emissions of this category account in 2022 for 17.1% and 21.1% of national total, respectively; SO_x

emission account for 10.5%; CO emissions account for 67.5%; Cd emissions account for 11.5%; PM10 and PM2.5 emissions account for 45.7% and 64.4% respectively while BC emissions account for 48.3%; dioxin is 33.0%, PAH is 80.4%, PCB is 17.1% and HCB is 31.2% of national totals. These emissions are prevalently due to biomass combustion, in winter, and they are also becoming critical for air quality issues. An in depth analysis of this category is reported in the paragraph 3.12.

Fugitive emissions in refinery from fossil fuel distribution and storage (1B2a iv) is key category in 2022 for SO_x emissions (16.0%). Total SO_x fugitive emissions from distribution of fossil fuels account for 20.3% of the total. Fugitive emissions from geothermal production are key category for Hg in 2022 accounting for 3.6% of the national total.

Table 3.1 Key categories in the energy sector in 2022

	1A1a	1A1b	1A1c	1A2	1A3a i	1A3a ii	1A3b i	1A3b ii	1A3b iii	1A3b iv	1A3b v	1A3b vi	1A3b vii	1A3c	1A3d ii	1A3e i	1A4a i	1A4bi	1A4bii	1A4c	1A5b	1B1a	1B1b	1B2
SO _x	7.9	6.2	1.5	32.3	0.3	0.2	0.3	0.1	0.1	0.0				0.0	12.4	0.0	4.7	5.8	0.0	0.1	0.2			20.3
NO _x	4.4	1.5	0.3	8.4	0.7	0.4	20.1	6.5	13.6	0.4				0.1	16.4	0.1	4.4	6.0	0.0	6.6	0.5			0.9
NH ₃	0.0	0.0	0.0	0.4			1.4	0.1	0.1	0.0				0.0	0.0		0.0	0.4	0.0	0.0	0.0			0.1
NM VOC	0.3	0.1	0.0	0.9	0.0	0.0	1.6	0.1	0.3	3.1	5.9			0.0	1.2	0.0	3.3	16.9	0.1	0.9	0.1		0.1	4.0
CO	1.1	0.2	0.4	5.0	0.1	0.1	8.7	0.4	1.4	4.3				0.0	2.3	0.0	1.4	63.7	0.0	2.4	1.1			0.0
PM ₁₀	0.2	0.1	0.0	3.8	0.0	0.0	1.3	0.3	0.7	0.2		5.1	2.3	0.0	3.8	0.0	0.7	44.3	0.0	0.7	0.2	0.2	0.0	0.0
PM _{2.5}	0.2	0.1	0.0	4.5	0.0	0.0	1.9	0.5	1.0	0.4		3.8	1.8	0.0	5.5	0.0	1.0	62.5	0.0	1.0	0.3	0.0	0.0	0.0
BC	0.1	0.0	0.0	1.5	0.1	0.0	12.3	2.9	5.3	0.5		7.1	0.6	0.0	10.0	0.0	0.8	44.7	0.0	2.9	1.2	0.2	0.1	0.0
Pb	0.8	0.1	0.0	38.1	0.3	0.2	0.0	0.0	0.0	0.0		18.8			0.1		2.3	4.3		0.0	0.0		0.2	0.0
Cd	1.5	0.4	0.0	29.2	0.0	0.0	6.6	0.8	0.7	0.7		4.2		0.0	0.5		1.9	9.1	0.0	0.5	0.0		2.0	0.0
Hg	8.4	1.6	0.2	22.7			2.2	0.4	0.7	0.1							3.6	4.1		0.3			0.9	3.6
PAH	0.6	0.1	0.0	0.8	0.0	0.0	2.7	0.5	0.8	0.0		0.1		0.0	0.1		1.0	78.7	0.0	0.7	0.0			0.0
Dioxin	0.8	0.9		19.5			2.0	0.2	0.3	0.1					0.1		1.0	31.5		0.4				
HCB	5.1			54.0			0.0	0.0	0.0	0.0					1.1		15.1	12.2		3.9				
PCB	0.3			10.2			0.0	0.0	0.0	0.0					0.1		2.9	14.0		0.3				

Note: key categories are shaded in blue

3.2 ENERGY INDUSTRIES

This chapter deals with emissions from categories 1A1a Public electricity and heat production, 1A1b Petroleum refining and 1A1c Manufacture of solid fuels and other energy industries.

3.2.1 Methodological issues

Methodologies used for estimating emissions from this sector are based on and conform to the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007; EMEP/EEA, 2013; EMEP/EEA, 2016; EMEP/EEA, 2019; EMEP/EEA), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000). A detailed description on the methods and national specific circumstances as well as reference material of the energy sector is documented in the national inventory report of the Italian greenhouse gas inventory (ISPRA, 2024[a]). The National Energy Balance, published by the Ministry of Environment (MASE), is the main source of information to estimate emissions from the energy sector as it reports fuel consumption for different sectors at national level. Additional information for electricity production is provided by the major national electricity producers and by the major national industry corporation. Other data are communicated by different category associations.

The use of PRTR data has been intensified in recent years, as also described in the following paragraphs. In this regard it should be noted that in response to the review process a survey has been conducted to verify if emission data submitted by operators are calculated subtracting the confidence interval. The issue has been discussed also with the colleagues from the Ministry of Environment (MASE) in charge of the implementation at national level for the IED legislation. In principle it is to be noted that the validated average values (with the confidence interval subtracted from the measured data) are the data used to verify the compliance of the operators to prescriptions included in the permits issued to the same operators and not for the calculation of the total annual emissions submitted in the framework of the relevant European Union Directives and Regulations. In addition, the implementation at national level of the IED requires Italian operators with emissions reported on the basis of Continuous Monitoring System (CMS) data to refer to confidence intervals which are not those included in the IED: in fact, the confidence intervals must result from quality assurance procedure and the implementation of UNI EN 14181:2005 and QAL2 procedure. This national circumstance implies that the validated average values used by the Italian operators are more realistic compared to those calculated using the procedure laid out in the IED. Moreover, the use of CMS at the stacks is largely implemented at facilities with installations exceeding 50 MWth input. In order to assess consistency along the timeseries, data reported by the largest Italian operator in the Energy production (about 25% of energy production) show that no issues concerning consistency can be raised. For consistency issue we believe that official data, as air emission values, communicated by the operators in the EU official frameworks, as the LCP Directive, PRTR registry and IPPC Directive should be considered as they were reported and without any further adjustment (apart from QA/QC procedures).

Notation key NO for activity is used indicating that a fuel is not consumed at all while NA is reported in the column where is requested to specify a different indicator than fuel consumption.

3.2.1.1 Public Electricity and Heat Production

For 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM10 emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM 2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level.

With regard to heavy metals country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies in 2001 while for PCB, emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have

been used following the recommendation of the review process (EEA, 2019). Emission factors for the PAH, Dioxin and HCB for Italy are from a study of TNO at European level (Berdowski et al, 1997). A comparison with data from PRTR has been made but in the case of HMs and POPs the information provided in that framework is generally poor and regard only few plants probably because of the ceiling which is high. For most of the pollutants the available information is not representative of the total. Dioxin, PCB and HCB emissions are generally missing. Gradually, in the next submissions, Italy plans to improve the quality of HMs and POPs estimate on the basis of the figures reported in the PRTR updating the EFs where they are representative of the national total or at least checking the order of magnitude of the values already used when the figures from PRTR are not sufficient to be considered representative of the total. In particular for power plants and coal fuelled main plants some HMs under the PRTR are available.

In the following Table 3.2 the main EFs for Public Electricity and Heat production sector are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM10 EFs too are country specific while PM2.5 EFs have been derived from PM10 EFs and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH EFs come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.2 Main emission factors for Public Electricity and Heat Production - 2022

	SO _x	NO _x	NMVOC	CO	Cd	Hg	Pb	PM2.5	Dioxins	PAH
	g/GJ				mg/GJ			g/GJ		
Steam coal	27.1	34.9	1.5	12.0	0.1	0.9	4.6	0.42	0.0038	0.0000009
Lignite	382.2	34.9	1.5	12.0	0.1	0.9	4.6	0.39	0.0096	0.0000024
Natural gas	0.1	17.3	2.5	20.0	0.0	0.3	0.6	0.01	-	-
Gasoil	13.9	50.0	1.5	12.0	0.1	0.8	2.5	0.05	0.0234	0.0009
Low sulphur fuel oil	27.9	26.7	3.0	15.0	0.1	0.8	2.5	0.77	0.0244	0.0009
High sulphur fuel oil	27.9	26.7	3.0	15.0	0.1	0.8	2.5	0.77	0.0244	0.0009
Other fuel oils	13.2	26.7	3.0	15.0	0.1	0.8	2.5	0.77	0.0244	0.0009
Biomass	1.8	48.8	1.5	12.0				2.10	0.0096	0.0037

3.2.1.2 Refineries

As stated above, for 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM10 emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level. In particular for 1A1b category, the implied emission factor refers both to the production of energy and heat and to the other combustion activities in refineries.

With regard to heavy metals country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies in 2001 while for PCB, emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have been used following the recommendation of the review process (EEA, 2019). A comparison with data from PRTR has been made but in the case of HMs and POPs the information provided in that framework is generally poor and regard only few plants probably because of the ceiling which is high. For most of the pollutants the available information is not representative of the total. Dioxin, PCB and HCB emissions are generally missing. For the next submissions Italy plans to improve the quality of HMs and POPs estimate on the basis of the figures reported in the PRTR updating the EFs where they are representative of the

national total or at least checking the order of magnitude of the values already used when the figures from PRTR are not sufficient to be considered representative of the total. For refineries a quite complete reporting is available for selenium and zinc. In the first case the average EFs is very close to the default used while for zinc the resulting EFs from EPRTTR is one order bigger than the default used up to now.

In the following Table 3.3 the main EFs for the Refinery sector are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM10 EFs too are country specific while PM2.5 EFs have been derived from PM10 EFs and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH EFs come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.3 Main emission factors for Refineries - 2022

	SO _x	NO _x	NMVOC	CO	Cd	Hg	Pb	PM2.5	Dioxins	PAH
	g/GJ				mg/GJ			g/GJ		
Combustion plants in refineries										
Refinery gas	0.4	27	2.5	13	-	-	-	5.00	-	-
Kerosene	55	27	3	60	0.14	0.77	2.52	2.50	0.023	0.00036
Petcoke	55	27	1.5	9	0.14	0.77	2.52	4.42	0.029	0.00036
Heavy residual + synthesis gasses	55	27	3	8	0.14	0.77	2.52	7.66	0.024	0.00036
High sulphur fuel oil	55	27	3	10	0.14	0.77	2.52	7.66	0.024	0.00036
Natural gas	0.2	8	2.5	13	0.03	0.31	0.60	0.20	-	-
LPG	0.2	27	2	10	-	-	-	0.20	0.022	-
Refinery furnaces										
LPG	0.2	41	2	10	-	-	-	0.20	0.022	-
Refinery gas	0.4	41	2.5	13.0	-	-	-	5.00	-	-
Kerosene	55	41	3	60	0.14	0.77	2.52	2.50	0.023	0.00036
Gasoline	23	41	3	60	0.14	0.77	2.52	1.25	0.023	0.00036
Gasoil	47	41	3	10	0.14	0.77	2.52	1.25	0.023	0.00036
High sulphur fuel oil	55	41	3	10	0.14	0.77	2.52	7.66	0.024	0.00036
Low sulphur fuel oil	55	41	3	10	0.14	0.77	2.52	7.66	0.024	0.00036
Petcoke	55	41	1.5	9	0.14	0.77	2.52	4.42	0.029	0.00036
No energy fuel	55	41	3	10	0.14	0.77	2.52	7.66	0.024	0.00036

3.2.1.3 Manufacture of Solid Fuels and Other Energy Industries

As already mentioned, for 1A1 categories, a Tier 3 is used and SO_x, NO_x and PM10 emissions are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTTR Regulation. The average implied emission factors at fuel level result from the analysis of the information provided and available at plant level, including technologies for energy production and emissions abatement. These IEF at fuel level have been used to estimate emissions for those plants where some pollutants have not been declared and to verify emissions declared. PM 2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level.

In particular, for coke production according to the review (EEA, 2019) PAH emission factor have been disaggregated into those deriving from the combustion process and the fugitive ones and estimated with the emission factors in the Guidebook (EMEP/EEA, 2019).

In the following Table 3.4 the main EFs for the Manufacture of Solid Fuels and Other Energy Industries are reported. The data shown in the blue cells are country specific emission factors as they are derived from specific measurement campaigns and direct communications by the operators. PM10 emission factors too are country specific while PM2.5 have been derived from PM10 and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. Finally, Dioxins and PAH emission factors come from a study of TNO at European level (Berdowski et al, 1997).

Table 3.4 Main emission factors for Manufacture of Solid Fuels and Other Energy Industries – 2022

	SO _x	NO _x	NM VOC	CO	Cd	Hg	Pb	PM2.5	Dioxins	PAH
	g/GJ				mg/GJ			g/GJ		
Combustion in coke oven										
Steam coal	27	40	1.5	12	0.15	0.85	4.63	3.50	0.004	0.000001
Natural gas	0.1	8	2.5	20	0.03	0.31	0.60	0.10	-	-
Oxygen furnaces gas	37	70	2.5	13	0.03	0.31	0.60	0.10	-	-
Coke oven gas	37	70	2.5	13	0.03	0.31	0.60	0.10	-	-
Blast furnace gas	37	70	2.5	13	0.03	0.31	0.60	0.10	-	-
High sulphur fuel oil	58	23	3	15	0.14	0.77	2.52	0.10	0.024	0.000890
Coke oven furnaces g/Mg										
Coke production	418	505	126	3861				5		0.008

3.2.2 Time series and key categories

The trend of emissions from energy industries has been influenced by the implementation of the legislative framework and by the evolution and replacement of fuels and these effects can be noticed in particular from the analysis of the NO_x and SO_x trends. The adoption of new regulations, in the late 80s, has led to a shift in energy consumption from oil with high sulphur content to oil with lower sulphur content and to natural gas.

In recent years, the conversion to the use of natural gas to replace fuel oil has intensified, thanks to incentives granted for the improvement of energy efficiency. Furthermore, a significant reduction in the use of coal fuels for energy production has been recorded in the last years. These measures, together with those of promoting renewable energy and energy saving, have led to a further reduction of emissions in the sector. In addition, in the last years, more stringent emission limits to the new plants have been established during the authorization process with the aim to prevent air quality issues at local level.

Until the 2023 inventory submission, Italy had not systematically assessed the uncertainty of national estimates. In 2023 Italy reports NO_x emissions uncertainty assessment (see paragraph 1.8 and Appendix 2), emission categories are disaggregated into a detailed level and uncertainties are therefore estimated for these categories. In addition, Italy has decided to achieve an in-depth analysis of the most important categories for which punctual data are available starting from the NO_x emissions of the thermoelectric power plants (1A1 and 1A2). On the basis of plant by plant consumptions and emissions data reported in the framework of E-PRTR and LCP registers, it has been possible to estimate uncertainty by fuel for natural gas (108 plants), coal (7 plants) and biomass (18 plants). The methodology consists into calculate, for each plant whose activity data and emissions are available, the weighted deviation based on fuel consumptions

and emissions. Calculating sigma as the square root of the normalized sum of deviations and considering the average emission factor and the number of plants it is possible to obtain the uncertainty for the considered plants. The consequent results are: for natural gas uncertainty equal to 1.2%, for biomass the uncertainty is equal to 5.3% and for coal equal to 9%.

3.2.2.1 *Public Electricity and Heat production*

This paragraph refers to the main electricity producers that produce electricity for the national grid. From 1998 onwards, the expansion of the industrial cogeneration of electricity and the split of the national monopoly have transformed many industrial producers into “independent producers”, regularly supplying the national grid. These producers account in 2022 for 89.3% of all electricity produced with combustion processes in Italy (TERNA, several years).

In 2022, Public Electricity and Heat production is a key category for SO_x, NO_x and Hg.

Public electricity and heat production (1A1a) is a key source of SO_x emissions in 2022 with a share of 7.9%, Hg (8.4) and NO_x emissions (4.4%). A strong reduction of SO_x, NO_x and PM10 emissions is observed for this category along the time series (as well as for 1A2 sector). The introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorization of plants, and the Ministerial Decree of 12th July 1990, which introduced plant level limits to emissions of PM10, NO_x and SO_x for new plants and required old plants to conform to the limit by 1997, explained the emission reduction in the nineties. The shift from fuel oil to natural gas combined with the increase of energy efficiency of the plants and the introduction of PM10 abatement technologies have been implemented to comply with the emission limit values. From 2000 lower limits to emissions at the stacks have been introduced, in the framework of environmental integrated authorizations, for the authorization of new plants and the implementation of the old ones, especially for those facilities located in areas with air quality critical values. For this reason, the plants have increased the use of natural gas heat and power combined technology. In 2022 in Italy there are still 6 coal plants, and 1 fuel oil plant out of around 200 power plants included in this source category. With exception of few biomass plants and some gasoil stationary engines in the small islands, the other plants are natural gas combined cycle thermoelectric power plant.

3.2.2.2 *Refineries*

This subsector covers the energy emissions from the national refineries (15 plants in 2022), including the energy used to generate electricity for internal use and exported to the national grid by power plants that directly use off-gases or other residues of the refineries. These power plants are generally owned by other companies but are located inside the refinery premises or just sideways. In 2022 the power plants included in this source category have generated 6.6% of all electricity produced with combustion processes in Italy.

Petroleum refining (1A1b) is a key category for SO_x in 2022. In addition, for 1990, petroleum refining (1A1b) is a key category for SO_x and PM2.5. Emissions are estimated on the basis of emission and consumption data provided by refineries in the framework of LCP, ETS European Directives and EPRTTR Regulation and refer both to the production of energy and heat and to the other combustion activities in the plants. Emission trends are driven by the same legislation quoted for 1A1a category, where specific rules and ceiling were set up for refineries.

3.2.2.3 *Manufacture of Solid Fuels and Other Energy Industries*

In this section, emissions from power plants, which use coal gases, are also reported. In particular, data refer to the electricity generated in the iron and steel plant sites (using coal gases and other fuels). In 2022 the power plants included in this source category have generated about 1.7% of all electricity produced with combustion processes in Italy.

Manufacture of Solid Fuels and Other Energy Industries (1A1c) is not a key category in 2022. Emissions show a decreasing trend linked to a reduction in activity data (only 2 plants remaining in 2022) and to the implementation of abatement technologies and the plants revamping. About abatement technologies, the largest integrated plant in Italy (and in Europe), in the last ten years, has carried out several interventions on the coking plant in the framework of the IPPC permit, applying BAT in several cases.


3.2.3 QA/QC and verification

A complete description of methodological and activity data improvements is documented every year in a QA/QC plan (ISPRA, 2024[b]).

The analysis of data collected from point sources allowed to distribute emissions at local level, for 2019 and previous years, as submitted under the CLTRAP. To illustrate an example, NO_x emissions from point sources are reported in for the year 2019. Point sources include public electricity and heat production plants and stationary combustion plants in manufacturing industries and construction.


The figure highlights that the most critical industrial areas are distributed in few regions. The visualization of the following maps created for the disaggregation of 2019, 2015 and 2010 demonstrates how information and data collection have been improved. More information in the chapter 10 and in the relevant publications (ISPRA, 2009; ISPRA, 2022).

Figure 3.1 NO_x emissions from point sources in 2019 (t)




Similarly, in Figure 3.2 NO_x emissions from point sources in 2015 are reported.

Figure 3.2 NO_x emissions from point sources in 2015 (t)



In Figure 3.3, NO_x emissions communicated by 229 facilities (power plants, refineries, cement plants and iron and steel integrated plants), in the framework of the national E-PRTR register and LCP Directive, have been processed and geographically located. The territorial distribution shows similar results to those reported in the previous figure highlighting the industrial areas still in activity in 2010.

Figure 3.3 NO_x emissions from point sources in 2010 (t)



3.2.4 Recalculations

In the 2024 submission recalculations have been performed for energy industries, in particular PM and BC emission factors from coke production have been updated for 2021. Minor recalculations (<0.1%) occur in 2010 and 2013-2020 because an update of heavy residual hydrocarbons in refineries.

3.2.5 Planned improvements

Specific improvements are detailed in the 2024 QA/QC plan (ISPRA, 2024[b]).

For the energy sector, a major progress regards the management of the information system where data collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. Moreover, the complete use of the energy data provided by the Ministry of Environment to the Joint Questionnaire IEA/OECD/EUROSTAT is planned in substitution of the national energy balances used till now; liquid, gaseous and solid fuel are now aligned for the whole time series and renewable fuels and biomass too have been used as possible.

Currently, a review process of the national energy balance is underway with Eurostat. In a first step, those responsible for the National Energy Balance implemented methodological changes for 2022 and subsequently applied these changes to 2021 to make it consistent. The review process, carried out in agreement with the relevant international institutions, regards definitions of “main activity producer” and “autoproducer” and the consequent adoption by Italy of these definitions in place of those previously used and linked to Italian legislation. This change in methodology has caused a fairly evident break in the series for some fuels and it is necessary further work to improve the time series consistency. In this framework, in 2024, it is planned the integration, where possible, of fuel consumption data reported under the ETS in the National Energy Balance.

With respect to PM10 and heavy metals emissions from Public Electricity and Heat Production category (1A1a) while PM10 emissions are updated every year on the basis of data submitted by the plants in the framework of the EPRTR registry, Large Combustion Plants Directive and Environmental Reports, heavy metals emission factors time series have been reconstructed from 1990 to 2001 on the basis of a study conducted by ENEL (major company in Italy) which reports heavy metals emissions measurements by fuel and technology (with or without PM10 abatement technologies) of relevant national plants. From 2001 these emission factors have not been updated. Heavy metals emission data in the EPRTR registry refer only to few not representative plants and are not sufficient to calculate average emission factors. Further work is planned to update/change emission factors for those pollutants where figures reported in the EPRTR lead to average values significantly different from those currently used.

3.3 MANUFACTURING INDUSTRIES AND CONSTRUCTION

This chapter deals with emissions from Stationary combustion in manufacturing industries and construction, in particular from categories 1A2a Iron and steel, 1A2b Non-ferrous metals, 1A2c Chemicals, 1A2d Pulp, Paper and print, 1A2e Food processing, beverages and tobacco, 1A2f Non-metallic minerals, 1A2g Other.

3.3.1 Methodological issues

For 1A2 categories, estimates for chemical, food processing, and other sectors (as textile, mechanics, extraction) are based on fuel consumption where EMEP/CORINAIR 2007 emission factors at fuel level have been used except for SO_x, NO_x and PM10 which are estimated on the basis of emission and consumption data provided by the relevant plants in the framework of LCP and ETS European Directives and EPRTR Regulation. PM 2.5 is estimated applying the ratio between PM2.5 and PM10 reported in the Tier 2 tables of the EMEP/EEA 2019 Guidebook at fuel level. Emissions of NH₃ have been also included when available at plant level. With regard to heavy metals, country specific emission factors for each fuel have been used to estimate emissions as provided by the main national operator in relation with the technologies, while for PCB emission factors for coal, oil products and wood biomass from the EMEP/EEA Guidebook 2019 have been used following the recommendation of the review process (EEA,2019). Emission factors for the PAH, Dioxin and HCB for Italy are from a study of TNO at European level (Berdowski et al, 1997). For the iron and steel, non-ferrous metal, pulp and paper and non-metallic minerals sectors emission estimates are based on production data at SNAP category level. SO_x, NO_x and PM10 emission factors time series are estimated based on the communication from operators in the framework of LCP Directive and EPRTR Regulation and industrial association at SNAP activity code level. For NMVOC, default EFs of EMEP/CORINAIR 2007 Guidebook are prevalently used except for glass and lead production where country specific emission factors are used; emission factors provided in the EMEP/EEA 2019 Guidebook are not appropriate because of they are calculated for small combustion boilers while emissions in this category refer prevalently to boilers >20 MWt for auto-production of energy and heat in the industrial sectors.

The Institute, specifically the same unit responsible for the inventory, also collects data in the context of the European Emissions Trading Scheme, the National Pollutant Release and Transfer Register (Italian PRTR) and the Large Combustion Plants (LCP) Directives. All these data are managed and used to compile the inventory. Figures are cross checked to develop country-specific emission factors and input activity data; whenever data cannot be straight used for the inventory compilation, they are considered as

verification. EPER/EPRTTR data are yearly available from 2002 while ETS data from 2005 and LCP data from 1990 all on yearly basis. In the EPRTTR registry total emissions divided by category are reported by plants if they exceed the relevant ceiling for each pollutant. LCP data refer only to SO_x, NO_x and PM emissions that are collected in stacks over 50 MWth and could result in figures lower than those reported in the EPRTTR. In the ETS only CO₂ and fuel consumption data are reported. QA/QC checks at plants level are directed to check the submissions of data in the different context and evaluate the differences if any. For example, if emissions submitted by a plant under LCP are higher than those submitted under the EPRTTR, ISPRA asks the operator of the reporting plant for an explanation and the verification of data submitted. In addition, on the basis of fuel consumption supplied under the ETS and average emission factor by fuel the energy experts estimate emissions at plant level and compare them with those submitted in the EPRTTR and LCP. Also in this case ISPRA ask for clarifications to the reporting plant if necessary.

3.3.1.1 Iron and steel

More in detail 1A2a includes combustion activities from the iron and steel sector as blast furnace cowpers, sinter plants and reheating furnaces. In 1990 there were four integrated iron and steel plants in Italy. In 2022, there is only one of the above-mentioned plants; oxygen steel production represents about 15.7% of the total production and the arc furnace steel the remaining 84.3% (FEDERACCIAI, several years). Also in 2022, Italy confirmed itself as the first European producer of cast steel from electric furnaces in absolute terms, contributing over 30% to the Union's electro-steel production, followed by Germany with 18.5% and Spain with 13.1% (FEDERACCIAI, several years). Currently, long products represent about 49.0% of steel production in Italy, flat products about 38.9%, and pipe the remaining 12.0%. Generally, the only integrated iron and steel plant is oriented towards flat production while, in steel plants equipped with electric ovens almost all located in the northern regions, long products are predominantly produced (e.g carbon steel, stainless steels) and seamless pipes (only one plant) (FEDERACCIAI, several years). Dioxins, PCB, HCB, PAH, Cd and Pb emissions are estimated on the basis of country specific emission factors at activity level, especially referring to sinter plants production, as provided by the main national operators. In particular, HCB emissions come from sinter plant productions and the emission factor is from the 2006 EMEP/CORINAIR Guidebook and it is coherent with data provided by the main national operator, at least for past years. The update of HMs and POPs emission factors with last available data started in the 2022 submission. On the basis of data collected in the framework of the IPPC permits, emission factors from point sources (sinter plants and blast furnaces) have been updated for As, Ni, Pb, Se, Cu, Hg, Zn, in the case of blast furnace it was also possible to estimate the PM10 emission factor. Again, on the basis of the IPPC data, the emission factors of NO_x, SO_x and PM10 from reheating furnaces have been estimated. Emission factors derived from this survey on the IPPC permits have been applied since 2016. As regards Cd emissions, these refer to blast furnaces, sinter and reheating activities; emission factors are those reported by the main Italian plant and emissions have been revised since 2003 based on the last review process and of E-PRTR data. For Hg and the other HMs emission factors are from the IPPC Bref sectoral report (JRC, 2013) and/or EMEP/EEA Guidebook 2006.

In the following Table 3.5 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework of EPRTTR, LCP, ETS or specific studies. Data reported in the dark blue cells represent country specific and fuel based emission factors. PM10 emission factors too, are country specific while PM2.5 factors have been derived from PM10 and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019.

Table 3.5 Main emission factors 1A2 Iron and steel - 2022

	SO _x	NO _x	NMVOC	CO	Cd	Hg	Pb	PM2.5	PAH	Dioxins
	Kg/Mg product				g/Mg product					mg/Mg
Blast furnace cowpers										
areal sources	0.300	0.100	0.0003	0.189	0.001	0.026	0.100	9.938	-	-
point sources	0.164	0.102	0.0003	0.189	0.001	0.001	0.007	0.754	-	-
Sinter and pelletizing plants										

	SO _x	NO _x	NMVOC	CO	Cd	Hg	Pb	PM2.5	PAH	Dioxins
	Kg/Mg product				g/Mg product					mg/Mg
areal sources	0.515	0.859	0.090	16.105	0.023	0.020	3.700	54.593	0.025	0.0009
point sources	0.430	0.603	0.090	14.750	0.002	0.004	0.323	6.397	0.025	0.0002
Reheating furnaces steel and iron										
areal sources	0.036	0.103	0.010	0.050	0.017	0.016	0.117	5.444	-	-
point sources	0.040	0.116	0.005	0.272	0.017	0.016	0.117	0.329	-	-

3.3.1.2 Non-ferrous metals

1A2b, non-ferrous metal sector, includes emissions from grey iron foundries, lead, zinc, copper and secondary aluminium production. In particular, emission from the production of lead and zinc at the moment are entirely reported in the energy sector because up to now there was no information to distinguish between energy and process emissions and, above all, these processes are considered combustion processes with contact, consequently, emissions are dependent on the combustion process. In particular, in Italy no production of primary copper has ever occurred while, as regards lead and zinc, there is a sole integrated plant for the primary productions, and this makes it difficult to ensure a good breakdown. Consequently, the issue related to the allocation of emissions is not only about combustion and process but also about the different productions of different metals in the same factory. To resolve this issue, an in-depth investigation has been started with the aim to better specify the technology used on the basis of E-PRTR and IPPC permits. The first result of this investigation has been the update of certain EFs since 2014 (ISPRA, 2023 [d]).

HCB emission factors available in the Guidebook refer to the consumption of coal and other solid fuels and wood biomass while in Italy only natural gas and small amount of LPG and fuel oil are used so that HCB and PCB emissions from secondary aluminum production have been estimated and reported based on the information available in a national study. Dioxin emissions from this category is driven by emissions from secondary aluminum production where country specific emission factors are used from a research project of 2002 based on measurements at production plant level; such emissions are due prevalently to the role played by recycled material and there is no evidence of changes in the quality of the aluminum scraps as well as in the pretreatment process. Average emission factor is equal to 69 micrograms per Mg, and it is in the range of values of the Guidebook reported in the IPPU relevant sector but representing total emissions from this category. Dioxins emissions reported in 1A2b occur also for secondary production of lead, zinc and copper but in total their emissions are less than 10% of the total of 1A2b, one order of magnitude lower than those from aluminum production. An investigation is ongoing with the aim to report these emissions separately in the energy and IPPU sectors. Because emissions are up to now reported in the 1.A.2.b category, notation key IE has been provided for 2.C.3 category. For Hg emission factors are from EMEP/CORINAIR 2007 Guidebook. Moreover, up to 2013, for primary and secondary lead production, emission factors for SO_x, NO_x, NMVOC, CO, Pb, PM10 are country specific, from a sectoral technical survey (ENEA, 2000) and from the communication of the operators, as well as for PAH e dioxins (ENEA-AIB-MATTM, 2002). For the other pollutants emission factors are from EMEP/CORINAIR 2007 but they have been shared and checked with the main operator. For primary zinc production, up to 2013, SO_x, CO, Pb, PM, Zn and Cd emission factors are country specific as provided by the only operator while for the other pollutants are from the EMEP/CORINAIR Guidebook 2007 taking in account the weight of the different production processes, electrolytic and Imperial Smelting Furnace. Thanks to the investigation above mentioned emission factors for NO_x, SO_x, PM10, Pb, Cd e Zn from zinc and lead production have been updated on the basis of data at plant level (ISPRA, 2023 [d]). For secondary aluminium production PAH and dioxins country specific emission factors have been used (ENEA-AIB-MATTM, 2002).

In the following Table 3.6 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework

of EPRT, LCP, ETS or specific studies. PM10 emission factors too are country specific while PM2.5 emission factors have been derived from PM10 emission factors and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. The remaining emission factors derive from the EMEP/EEA Guidebook.

Table 3.6 Main emission factors 1A2 Non-Ferrous metals - 2022

	SO _x	NO _x	NM VOC	CO	Cd	Hg	Pb	PM2.5	PAH	Dioxins
	Kg/Mg product				g/Mg product				mg/Mgproduct	
Grey iron foundries	0.125	0.160	0.090	9.500	0.140		7.200	423.529		
Primary lead production	0.748	0.104	0.000	0.000	0.312	3.000	14.270	18.860		
Primary zinc production	0.071	0.643		6.953	0.106	6.120	0.819	16.094		
Primary copper production										
Secondary lead production	2.012	0.191	0.000	0.000	0.000		1.563	11.675	0.006	0.005
Secondary zinc production	0.077	0.777	0.000	11.336	0.093	0.020	1.460	13.291		0.065
Secondary copper production		0.200	2.000		5.000		50.000	967.989		0.020
Secondary aluminium production	1.300	0.400	1.250					258.008	0.189	0.068
Other										
copper manufactures		0.200	2.000							
zinc-copper & brass manufactures		0.200	2.000							

3.3.1.3 Chemicals

Category 1A2c refer to the stationary combustion emissions from the chemical and petrochemical sectors. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas and petrochemical gases are prevalently used in gas turbine plants. It is not level key category for any pollutant, while 1A2c is a key category at trend assessment for NO_x and SO_x. Emissions from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRT and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.4 Pulp, paper and print

Category 1A2d refer to the stationary combustion emissions from the pulp and paper sector. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas is prevalently used in gas turbine plants. It is not level and trend key category for any pollutant. Emissions from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRT and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.5 Food processing, beverages and tobacco

Category 1A2e refer to the stationary combustion emissions from the food and drink sector. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas, and biogas are prevalently used in gas turbine plants. It is not level and trend key category for any pollutant. Emissions from the production processes are reported in the relevant IPPU sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.1.6 Non-metallic minerals

Category 1A2f, stationary combustion in non-metallic mineral industry, refers to a multitude of production activities such as cement, lime, glass, brick and tiles, ceramics, and asphalt production which means a multitude of different emission factors. For cement production, PM emissions from kilns are reported in this category where emissions from mills are reported in IPPU (emission factor from USEPA 1991 emission factor handbook) while for lime production PM emission factors referring to the complete process are used (from USEPA 1996 emission factor handbook) and emissions are distributed between energy and IPPU. For Hg, emission factors are country specific (especially cement production which is the emission driver of this category); for Dioxin, HCB, PCB and Cd emission factors are from the relevant Bref reports or EMEP/EEA 2007 Guidebook; for Pb, emission factors are country specific for ceramic production and from the Bref report or EMEP 2006 Guidebook for glass, cement and lime productions.

In the following Table 3.7 the main emission factors for the sector are reported. The data shown in the light blue cells are country specific based on emission data at plant level communicated in the framework of EPRTTR, LCP, ETS or specific studies. PM10 emission factors too are country specific while PM2.5 emission factors have been derived from PM10 emission factors and the ratio PM2.5/PM10 per fuel as reported in the EMEP/EEA Guidebook 2019. The remaining emission factors derive from the EMEP/EEA Guidebook.

Table 3.7 Main emission factors 1A2 Non-Metallic minerals - 2022

	SO _x	NO _x	NMVOC	CO	Cd	Hg	Pb	PM2.5	PAH	Dioxins
	Kg/Mg product				g/Mg product					mg/Mg product
Plaster furnaces	0.115	0.088	0.009	0.013				0.990		
Cement	0.409	1.049	0.023	0.981	0.008	0.030	0.006	2.183		0.0001
Lime										
areal sources	0.027	0.937	0.004	0.013		0.001		1.200		
point sources	0.119	0.037	0.004	0.013		0.001		0.600		
Asphalt concrete plants	0.002	0.011	0.004	0.002				6.500	0.0004	
Flat glass	0.890	2.183	0.044	0.100			0.330	350.000		
Container glass	1.291	1.464	0.047	0.100	0.090		4.550	448.800		
Glass wool (except binding)	2.500	0.968	0.048	0.081				123.390		
Other glass		2.000	0.100	0.260			10.000	17.778		

	SO _x	NO _x	NM VOC	CO	Cd	Hg	Pb	PM2.5	PAH	Dioxins
	Kg/Mg product				g/Mg product					mg/Mg product
Mineral wool (except binding)	1.500			3.200				493.559		
Bricks and tiles	0.785	0.250	0.007	0.040				49.841		
Fine ceramic materials	0.569	0.174	0.014	0.130			6.000	193.898		

3.3.1.7 Other

Category 1A2g refer to the stationary combustion emissions from different industrial sectors as machinery, mining and quarrying, construction, textile and others. Emissions from this category are calculated on the basis of fuel consumption and the relevant emission factors and they are due to the auto production of energy and heat for the production processes. Natural gas is prevalently used in gas turbine plants. Emissions from off-roads vehicles used in industry are also reported under this category. Fugitive emissions from the oil and gas extraction activities are reported in the relevant energy sub sector. SO_x, NO_x and PM10 emission factors are country specific and derive from the communications of the main operators in the framework of the EPRTR and LCP framework while fuel consumption at plant level are provided in the ETS together with CO₂ emissions.

3.3.2 Time series and key categories

In 2022, manufacturing industries and construction (1A2) is a main source of heavy metals and POPs, accounting for about 54% of HCB total emissions, 38% for lead, 32% for SO_x, 29% for cadmium, 23% for Hg and 20% for dioxin. The sector is key category also for PM10 and PM2.5 (4% and 5% respectively) as well as NO_x and CO, about 8% and 5% of total emissions. In 2022 the main sectors are Stationary combustion in iron and steel industries (1A2a), which is key for CO, Cd, and Hg, Stationary combustion in non-ferrous metal industries (1A2b), key for Dioxin and HCB, and stationary combustion in non-metallic mineral industries (1A2f) that is key category for SO_x, NO_x, PM10, PM2.5, Pb, Cd and Hg.

As already reported for 1A1 category, a strong reduction of SO_x, NO_x and PM10 emissions is observed for this category along the time series. The introduction of two regulatory instruments: the DPR 203/88 (Decree of President of the Republic of 24th May 1988), laying down rules concerning the authorization of plants, and the Ministerial Decree of 12th July 1990, which introduced plant level limits to emissions of PM10, NO_x and SO_x for new plants and required old plants to conform to the limit by 1997, explained the emission reduction in the nineties. The shift from fuel oil to natural gas combined with the increase of energy efficiency of the plants and the introduction of PM10 abatement technologies have been implemented to comply with the emission limit values. From 2000 lower limits to emissions at the stacks have been introduced, in the framework of environmental integrated authorizations, for the authorization of new plants and the implementation of the old ones, especially for those facilities located in areas with air quality critical values. For this reason, the plants have increased the use of natural gas heat and power combined technology.

3.3.2.1 Iron and steel

The category 1A2a in 2022 is a key category for CO, Cd and Hg. The trend of emissions is linked above all to the production levels and then to the abatement technologies.

In fact, for the majority of pollutants, in 2009 a strong reduction of emissions is observed due to the effects of the economic recession that in 2010 and 2011 has partially recovered. In 2012 a further drop occurred for the economic crisis and for environmental constrains of the main iron and steel integrated plants that should reduce its productions. In 2015 a drop is still observed consistently with the production

activities reduction of the main iron and steel integrated plants. There were four integrated steel plants in 1990 that from 2005 are reduced to two, with another plant that still has a limited production of pig iron. Nevertheless, the steel production in integrated plants has not changed significantly in the 1990-2008 period due to an expansion in capacity of the two operating plants. From 2015 only one integrated plant remains in operation. The maximum production was around 11 Mt/y in 1995 and in 2005-2008, with lower values in other years and the lowest of 3.4 Mt in 2022.

3.3.2.2 *Non-ferrous metals*

The category 1A2b in 2022 is a key category for dioxins and HCB.

In Italy, the production of primary aluminium stopped in 2013 (and was 232 Gg in 1990) while secondary aluminium accounts for 350 Gg in 1990 and 717 Gg in 2022. These productions, however, use electricity as the primary energy source so the emissions due to the direct use of fossil fuels are limited. The sub sector comprises also the production of other non-ferrous metals, both primary and secondary copper, lead, zinc and others; as above reported, point data by plant authorization for the sole integrated Zn/Pb plant led to more accurate EFs for NO_x, SO_x, CO, PM, Cd, Pb and Zn and, as a consequence, some limited recalculations.

3.3.2.3 *Chemicals*

In Italy there are petrochemical plants integrated with a nearby refinery and stand-alone plants that get the inputs from the market. Main products are Ethylene, Propylene, Styrene. In particular, ethylene and propylene are produced in petrochemical industry by steam cracking. Ethylene is used to manufacture ethylene oxide, styrene monomer and polyethylene. Propylene is used to manufacture polypropylene but also acetone and phenol. Styrene, also known as vinyl benzene, is produced on industrial scale by catalytic dehydrogenation of ethyl benzene. Styrene is used in the rubber and plastic industry to manufacture through polymerisation processes such products as polystyrene, ABS, SBR rubber, SBR latex. Except for ethylene oxide, whose production has stopped in 2002, the other productions of the above-mentioned chemicals still occur in Italy. Activity data are stable from 1990 to 2012, with limited yearly variations along the timeseries and a reduction in the last years. Chemical industry includes non-organic chemicals as chlorine/soda, sulphuric acid, nitric acid, ammonia. A limited production of fertilizers is also present in Italy. From 1990 to 2022 the sum of productions of this source category has greatly reduced: in 2022 it was about 40% of the production in 1990.

3.3.2.4 *Pulp, paper and print*

Emissions from the manufacturing of paper are included in this source category. In Italy the manufacture of virgin paper pulp is rather limited, with a production feeding less than 5% of the produced paper. Most of the pulp was imported in 1990, while in 2022 half of the pulp used is produced locally from recycled paper. The paper production is expanding, and activity data (total paper produced) were 6.2 Mt in 1990 and 8.7 Mt in 2022. The printing industry represents a minor part of the source category emissions.

3.3.2.5 *Food processing, beverages and tobacco*

In Italy the food production industry is expanding. A comprehensive activity data for this sector is not available; more in detail while energy data are those reported in the national energy balance for this sector, information at subsector and technological level is not available and only few plants are part of the ETS; energy fuel consumption was estimated to be 62 PJ in 1990 and 120 PJ in 2022, about half of energy consumptions derives from biomass (99.5% biogas). Value added at constant prices has increased of 0.6% per years from 1990 to 2003 and almost constant from 2004.

3.3.2.6 *Non-metallic minerals*

The category 1A2f in 2022 is a key category for SO_x, NO_x, PM10, PM2.5, Pb, Cd and Hg. This sector, which refers to construction materials, is quite significant in terms of emissions due to the energy intensity of the processes involved. Construction materials subsector includes the production of cement, lime, bricks, tiles and glass. It comprises thousands of small and medium size enterprises, with only few large operators, mainly connected to cement production. Some of the production is also exported.

3.3.2.7 *Other*

This sector comprises emissions from many different industrial subsectors, some of which are quite significant in Italy in terms of both value added and export capacity. In particular, engineering sectors (vehicles and machines manufacturing) is the main industrial sub sector in terms of value added and revenues from export and textiles was the second subsector up to year 2000. The remaining “other industries” include furniture and other various “made in Italy” products that produce not negligible amounts of emissions.

3.3.3 QA/QC and verification

As already reported, QA/QC checks at plants level are directed to check the submissions of data in the different context and evaluate the differences if any. For example, if emissions submitted by a plant under LCP are higher than those submitted under the EPRTR, ISPRA asks the operator of the reporting plant for an explanation and the verification of data submitted. In addition, on the basis of fuel consumption supplied under the ETS and average emission factor by fuel energy, experts estimate emissions at plant level and compare them with those submitted in the EPRTR and LCP. Also, in this case ISPRA asks for clarifications to the reporting plant if necessary. As for 1A1 point sources, the analysis of data collected from point sources allowed to distribute emissions at local level and to check and verify data at local level.

3.3.4 Recalculations

For 1.A.2 category some recalculations occurred in this submission, as a follow up to the review process. Recalculations in 2021 are due to the update of activity data reported in the National Energy Balance.

Ammonia emissions from 1A2e -Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco have been revised on the consequence of NECD review process (EEA, 2023) to introduce NH₃ emissions from solid biomass combustion.

For 1.A.2.b, zinc and lead production emissions only minor recalculations from 2014 for Pb, Cd, PM.

Minor recalculations (<0.5%) occur because the update heavy residual hydrocarbons data for 2010 and 2012-2020.

3.3.5 Planned improvements

Specific improvements are detailed in the 2024 QA/QC plan (ISPRA, 2024[b]).

Following the update of lead and zinc production the reconstruction of the time series ante 2014 is planned to ensure the consistency.

For the energy sector, a major progress regards the management of the information system where data collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, are gathered together thus highlighting the main discrepancies in information and detecting potential errors. Moreover, the complete use of the energy data provided by the Ministry of Environment to the Joint Questionnaire IEA/OECD/EUROSTAT is planned in substitution of the national energy balances used till now; liquid, gaseous and solid fuel are now aligned for the whole time series and renewable fuels and biomass too have been used as possible.

3.4 AVIATION (NFR SUBSECTOR 1.A.3.A)

3.4.1 Overview

Emissions from categories 1.A.3.a.i International Aviation and 1.A.3.a.ii Domestic Aviation are estimated, including figures both for landing and take-off cycles (LTO) and for the cruise phase of the flight (the latter reported as memo items and not included in the national totals).

3.4.2 Methodological issues

According to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2006; IPCC, 2000) and the EMEP/EEA Guidebook 2023 (EMEP/EEA, 2023), a national technique has been developed and applied to estimate emissions. The current method estimates emissions from the following assumptions and information. Activity data comprise both fuel consumptions and aircraft movements, which are available in different level of aggregation and derive from different sources as specified here below:

- Total inland deliveries of aviation gasoline and jet fuel are provided in the national energy balance (MSE, several years (a)). This figure is the best approximation of aviation fuel consumption, for international and domestic use, but it is reported as a total and not split between domestic and international.
- Data on annual arrivals and departures of domestic and international landing and take-off cycles at Italian airports are reported by different sources: National Institute of Statistics in the statistics yearbooks (ISTAT, several years), Ministry of Transport in the national transport statistics yearbooks (MIMS, several years), the Italian civil aviation in the national aviation statistics yearbooks (ENAC/MIMS, several years), EUROCONTROL flights data time series 2002–2022 (EUROCONTROL, several years).

An overall assessment and comparison with EUROCONTROL emission estimates was carried out over the years and that lead to an update of the methodology used by Italy for this category. Data on the number of flights, fuel consumption and emission factors were provided by EUROCONTROL in the framework of a specific project funded by the European Commission, and quality checked by the European Environmental Agency and its relevant Topic Centre (ETC/ACM), aimed at improving the reporting and the quality of emission estimates from the aviation sector of each EU Member State under both the UNFCCC and LRTAP conventions. The Advanced Emissions Model (AEM) was applied by EUROCONTROL to derive these figures, according to a Tier 3 methodology (EMEP/EEA, 2023). EUROCONTROL fuel and emissions time series cover the period 2005–2022, while the number of flights is available since 2002. For the time series from 1990 to 1999, figures for emission and consumption factors are derived by the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007), both for LTO cycles and cruise phases, taking into account national specificities.

These specificities derived from the results of a national study which, taking into account detailed information on the Italian air fleet and the origin–destination flights for the year 1999, calculated national values for both domestic and international flights (Romano et al., 1999; ANPA, 2001; Trozzi et al., 2002 (a)) on the basis of the default emission and consumption factors reported in the EMEP/CORINAIR guidebook. National average emissions and consumption factors were therefore estimated for LTO cycles and cruise both for domestic and international flights from 1990 to 1999. Specifically, for the year referred to in the survey, the method estimates emissions from the number of aircraft movements broken down by aircraft and engine type (derived from ICAO database if not specified) at each of the principal Italian airports; information about whether the flight is international or domestic and the related distance travelled has also been considered. A Tier 3 method has been applied for 1999. In fact, figures on the number of flights, destination, aircraft fleet and engines have been provided by the local airport authorities, national airlines and EUROCONTROL, covering about 80% of the national official statistics on aircraft movements for the relevant years. Data on ‘Times in mode’ have also been supplied by the four principal airports and estimates for the other minor airports have been carried out on the basis of previous sectoral studies at local level. Consumption and emission factors are those derived from the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). Based on sample information, estimates have been

carried out at national level from 1990 to 1999 considering the official statistics of the aviation sector (ENAC/MIMS, several years) and applying the average consumption and emission factors.

From 2005, fuel consumption and emission factors were derived from the database made available to EU Member States by EUROCONTROL, as previously described. These data were used for updating fuel consumption factors, and emission factors of all pollutants. For the period between 1999 and 2005, interpolation has been applied to calculate these parameters. Estimates were carried out applying the consumption and emission factors to the national official aviation statistics (ENAC/MIMS, several years) and EUROCONTROL data on movements from 2002 (EUROCONTROL, several years). In general, to carry out national estimates of greenhouse gases and other pollutants for LTO cycles, both domestic and international, consumptions and emissions are calculated for the complete time series using the average consumption and emission factors multiplied by the total number of flights. The same method is used to estimate emissions for domestic cruise; on the other hand, for international cruise, consumptions are derived by difference from the total fuel consumption reported in the national energy balance and the estimated values as described above and emissions are therefore calculated.

The fuel split between national and international fuel use in aviation is then supplied to the Ministry of the Economic Development to be included in the official international submission of energy statistics to the IEA in the framework of the Joint Questionnaire OECD/EUROSTAT/IEA compilation together with other energy data. Data on domestic and international aircraft movements from 1990 to 2022 are shown in Table 3.8 where domestic flights are those entirely within Italy. Since 2002, EUROCONTROL flights data have been considered, accounting for departures from and arrivals to all airports in Italy, regarding flights flying under instrument flight rules (IFR), including civil helicopters flights and excluding flights flagged as military, when the above flights can be identified. Total fuel consumptions, both domestic and international, are reported by LTO and cruise in Table 3.9.

Table 3.8 Aircraft Movement Data (LTO cycles)

Number of flights	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Domestic flights	172,148	185,220	319,748	350,140	354,520	280,645	288,470	151,156	209,721	286,496
International flights	147,875	198,848	303,608	381,206	406,990	425,410	502,764	172,835	232,750	419,862

Source: ISTAT, several years; ENAC/MIMS, several years; Eurocontrol, several years.

Table 3.9 Aviation jet fuel consumptions for domestic and international flights (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
Domestic LTO	111	120	208	233	227	168	180	89	123	176
International LTO	130	175	258	269	296	328	399	130	162	301
Domestic cruise	357	384	654	666	704	526	580	291	421	620
International cruise	1,246	1,688	2,297	2,456	2,534	2,745	3,584	1,087	1,432	2,625

Source: ISPRA elaborations

Emissions from military aircrafts are also estimated and reported under category 1.A.5 Other. The methodology to estimate military aviation emissions is simpler than the one described for civil aviation since LTO data are not available in this case. As for activity data, total consumption for military aviation is published in the petrochemical bulletin (MSE, several years (b)) by fuel. Emission factors are those provided in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). Therefore, emissions are calculated by multiplying military fuel consumption data for the EMEP/CORINAIR default emission factors.

3.4.3 Time series and key categories

Emission time series of NO_x, NMVOC, SO_x, TSP, CO, Pb are reported in Table 3.10, Table 3.11, Table 3.12, Table 3.13, Table 3.14 and Table 3.15, respectively.

An upward trend in emission levels for civil aviation is observed from 1990 to 2019, which is explained by the increasing number of LTO cycles. Nevertheless, the propagation of more modern aircrafts in the fleet slows down the trend in the most recent years. There has also been a decrease in the number of domestic flights from 2000, although a new increasing trend in the last couple of years has been registered. Year 2020 is to be considered separately, because the aviation sector was severely hit by the pandemic measures: domestic flights in 2020 were about the 52% of the 2019, while international flights during 2020 were the 34% of the flights of the previous year. In year 2022 domestic flights have increased by 37%, while international flights have increased by 80% in comparison with year 2021. Aviation is not a key category.

Table 3.10 Time series of NO_x (Gg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	1.36	1.47	2.50	2.55	2.71	2.12	2.32	1.15	1.59	2.36
1 A 3 a i (i) International aviation LTO (civil)	1.60	2.16	3.20	3.47	3.99	4.55	5.60	1.85	2.33	4.32
1 A 3 a Civil Aviation (LTO)	2.97	3.62	5.70	6.02	6.70	6.68	7.92	3.01	3.92	6.68
1A3 a ii (ii) Domestic aviation cruise (civil)	5.23	5.63	9.43	8.71	10.16	8.09	8.78	4.25	5.68	8.55
1A3a i (ii) International aviation cruise (civil)	18.85	26.83	38.99	36.55	41.02	47.05	59.65	18.56	19.05	34.51
1 A 5 b Other, Mobile (including military, land based and recreational boats)	11.16	11.99	7.24	13.50	6.11	3.29	2.73	3.48	1.94	2.79

Table 3.11 Time series of NMVOC (Gg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	0.12	0.13	0.23	0.25	0.33	0.27	0.30	0.14	0.18	0.25
1 A 3 a i (i) International aviation LTO (civil)	0.15	0.20	0.31	0.39	0.45	0.48	0.52	0.16	0.20	0.36
1 A 3 a Civil Aviation (LTO)	0.27	0.33	0.54	0.64	0.78	0.75	0.82	0.31	0.39	0.61
1A3 a ii (ii) Domestic aviation cruise (civil)	0.10	0.10	0.18	0.20	0.37	0.34	0.38	0.19	0.26	0.34
1A3a i (ii) International aviation cruise (civil)	0.25	0.36	0.55	0.69	0.81	0.89	0.94	0.29	0.37	0.69

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 5 b Other, Mobile (including military, land based and recreational boats)	3.00	3.13	1.90	3.00	1.05	0.66	0.58	0.77	0.41	0.62

Table 3.12 Time series of SO_x (Gg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	0.11	0.12	0.21	0.23	0.23	0.17	0.18	0.09	0.12	0.18
1 A 3 a i (i) International aviation LTO (civil)	0.13	0.17	0.26	0.27	0.30	0.33	0.40	0.13	0.16	0.30
1 A 3 a Civil Aviation (LTO)	0.24	0.29	0.47	0.50	0.52	0.50	0.58	0.22	0.28	0.48
1A3 a ii (ii) Domestic aviation cruise (civil)	0.36	0.38	0.65	0.67	0.70	0.53	0.58	0.29	0.42	0.62
1A3a i (ii) International aviation cruise (civil)	1.25	1.78	2.60	2.45	2.65	2.95	3.69	1.16	1.36	2.45
1 A 5 b Other, Mobile (including military, land based and recreational boats)	1.19	0.81	0.21	0.17	0.13	0.12	0.13	0.19	0.09	0.16

Table 3.13 Time series of TSP (Gg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	0.01	0.01	0.02	0.03	0.02	0.02	0.02	0.01	0.01	0.02
1 A 3 a i (i) International aviation LTO (civil)	0.02	0.03	0.04	0.05	0.03	0.04	0.04	0.01	0.02	0.03
1 A 3 a Civil Aviation (LTO)	0.03	0.04	0.06	0.07	0.06	0.05	0.06	0.02	0.03	0.05
1A3 a ii (ii) Domestic aviation cruise (civil)	0.07	0.08	0.13	0.10	0.10	0.07	0.08	0.04	0.06	0.09
1A3a i (ii) International aviation cruise (civil)	0.36	0.52	0.75	0.71	0.83	0.91	0.96	0.30	0.37	0.71
1 A 5 b Other, Mobile (including military, land based and recreational boats)	1.30	1.57	0.91	1.63	0.83	0.48	0.43	0.56	0.30	0.46

Table 3.14 Time series of CO (Gg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	1.23	1.33	2.26	2.33	2.28	1.69	1.88	0.94	1.24	1.73
1 A 3 a i (i) International aviation LTO (civil)	1.73	2.32	3.33	2.86	3.00	3.25	3.89	1.25	1.48	2.73
1 A 3 a Civil Aviation (LTO)	2.96	3.64	5.59	5.19	5.28	4.94	5.77	2.19	2.72	4.46
1A3 a ii (ii) Domestic aviation cruise (civil)	1.31	1.41	2.43	2.66	3.07	2.35	2.60	1.47	2.18	2.83
1A3a i (ii) International aviation cruise (civil)	2.03	2.89	4.42	5.55	5.74	6.07	7.23	2.49	3.65	6.47
1 A 5 b Other, Mobile (including military, land based and recreational boats)	65.12	79.02	45.49	54.48	17.33	16.49	17.42	24.93	11.98	20.34

Table 3.15 Time series of Pb (Mg)

Source categories for NFR Subsector 1.A.3.a, 1.A.5.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Mg									
1 A 3 a ii (i) Domestic aviation LTO (civil)	0.19	0.20	0.35	0.38	0.38	0.30	0.31	0.16	0.23	0.31
1 A 3 a i (i) International aviation LTO (civil)	0.21	0.28	0.43	0.54	0.57	0.60	0.71	0.24	0.33	0.59
1 A 3 a Civil Aviation (LTO)	0.39	0.48	0.77	0.91	0.96	0.90	1.02	0.41	0.55	0.90
1A3 a ii (ii) Domestic aviation cruise (civil)	0.57	0.62	1.06	1.16	1.18	0.93	0.96	0.50	0.70	0.95
1A3a i (ii) International aviation cruise (civil)	2.01	2.86	4.36	5.48	5.85	6.11	7.22	2.48	3.34	6.03
1 A 5 b Other, Mobile (including military, land based and recreational boats)	16.34	4.22	1.16	0.001	NA	0.12	0.18	0.18	0.10	0.09

3.4.4 QA/QC and Uncertainty

Data used for estimating emissions from the aviation sector derive from different sources: local airport authorities, national airlines operators, EUROCONTROL and official statistics by different Ministries and national authorities.

Different QA/QC and verification activities are carried out for this category. As regards past years, the results of the national studies and methodologies, applied at national and airport level, were shared with national experts in the framework of an ad hoc working group on air emissions instituted by the National

Aviation Authority (ENAC). The group, chaired by ISPRA, included participants from ENAC, Ministry of Environment, Land and Sea, Ministry of Transport, national airlines and local airport authorities. The results reflected differences between airports, aircrafts used and times in mode spent for each operation.

Currently, verification and comparison activities regard activity data and emission factors. In particular, number of flights have been compared considering different sources: ENAC, ASSAEROPORTI, ISTAT, EUROCONTROL and verification activities have been performed on the basis of the updated EUROCONTROL data on fuel consumption and emission factors resulting in an update and improving of the national inventory. Furthermore, there is an ongoing collaboration and data exchange with regional environmental agencies on this issue.

3.4.5 Recalculations

No recalculations have been done in this submission.

3.4.6 Planned improvements

Improvements for next submissions are planned on the basis of the outcome of the ongoing quality assurance and quality control activities, in particular with regard to the results of investigation about data and information deriving from different sources, in particular further assessment of EUROCONTROL data, and comparison with information provided by the national institute of statistics, ISTAT, on the number of flights.

3.5 ROAD TRANSPORT (NFR SUBSECTOR 1.A.3.B)

3.5.1 Overview

The road transport sector contributes to the total national emissions in 2022 as follows: nitrogen oxides emissions for 40.6% of the total, emissions of carbon monoxide for 14.8%, non-methane volatile organic compounds for 11.0%, PM10 and PM2.5, for 10.0% and 9.3%, respectively, of the total.

The estimation refers to the following vehicle categories:

- 1.A.3.b.i Passenger cars
- 1.A.3.b.ii Light-duty trucks
- 1.A.3.b.iii Heavy-duty vehicles including buses
- 1.A.3.b.iv Mopeds and motorcycles
- 1.A.3.b.v Gasoline evaporation
- 1.A.3.b.vi Road transport: Automobile tyre and brake wear
- 1.A.3.b.vii Road transport: Automobile road abrasion

3.5.2 Methodological issues

A national methodology has been developed and applied to estimate emissions according to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2000; IPCC, 2006) and the EMEP/EEA Guidebook (EMEP/EEA, 2023).

In general, the annual update of the model is based on the availability of new measurements and studies regarding road transport emissions (for further information see: <https://www.emisia.com/utilities/copert/>). The model COPERT 5 (updated version 5.7.3, January 2024) has been used and applied for the whole time series in 2024 submission. COPERT 5 introduced over the years upgrades both from software and methodological point of view (<https://www.emisia.com/utilities/copert/versions/>). Several new methodological features had been introduced respect to the previous model COPERT 4. As regards fuel, updates concerned: fuel energy instead of fuel mass calculations; distinction between primary and end (blends) fuels, automated energy balance. Regarding vehicle types, updated vehicle category naming, new vehicle types and emission control technology level, have been introduced. As regards emission factors, one function type and the possibility to distinguish between peak/off-peak urban, have been implemented.

Main methodological innovations introduced in version 5.7.3 respect to version 5.6.1, used in last submission, relate: updating emission factors of Euro 6 CNG passenger cars, Euro VI diesel buses, Euro VI diesel hybrid buses, non-exhaust emission factors.

As regards the software, revisions relate: the removal of CO₂ correction, capability of alternative HDVs classification based on REG EU 2017/2400, improved labels of forms and headers.

Furthermore various bugs have been corrected regarding: the correction of the calculation for cold emissions of CO, NO_x, VOC for petrol and diesel PCs and LCVs, the correction of cold start ratio of diesel Euro 6 cars for NO_x and CO, the correction of cold start ratio of petrol-fueled cars and vans for VOC and CO, the correction of cold emissions of Euro 6 CNG passenger cars for SPN23, the correction of bug when importing stock of HDVs in VECTO groups from excel to COPERT.

In addition, software update issues and minor corrections have been implemented.

The model, on the basis of the inputs inserted, gives output results separately for vehicles category and urban (peak/off-peak urban), rural, highway areas, concerning emission estimates of CO, VOC, NMVOC, CH₄, NO_x, N₂O, NH₃, PM2.5, PM10, PM exhaust (the emission factors of particulate matter from combustion refer to particles smaller than 2.5 µm, that implicitly assumes that the fraction of particulate matter with diameter between 2.5 µm and 10 µm is negligible), CO₂, SO₂, heavy metals, NO_x speciation in NO and NO₂, the speciation in elemental and organic carbon of PM, the speciation of NMVOC. Resulting national emission factors at detailed level are available on the following public web address: <https://fetransp.isprambiente.it/#/>. Data on fuel consumption of gasoline, diesel, liquefied petroleum gas (LPG), natural gas (CNG) and biofuels are those reported in the EUROSTAT energy balance; Italian road

vehicles electricity consumption data, introduced recently in COPERT in relation to the evolution of the fleet, derive from Eurostat database (<https://ec.europa.eu/eurostat/data/database>). Time series of consumptions, by fuel and vehicle categories, are detailed in the NFR. Lubricants consumption is estimated and reported in 1A3b.

3.5.2.1 Exhaust emissions

Exhaust emissions from vehicles subsectors are split between cold and hot emissions; estimates are calculated either on the basis of a combination of total fuel consumption and fuel properties data or on the basis of a combination of drive related emission factors and road traffic data.

The calculation of emissions is based on emission factors calculated for the vehicle models most widely and systematically used, distinguishing between the type of vehicle, fuel, engine size or weight class, standard legislation. The legislative standards introduced become more stringent over the years, ensuring that new vehicles emit much less than the older ones as regards the regulated pollutants. With reference to four groups of pollutants, the method of calculation of exhaust emissions is different. The methodology implemented is derived from the EMEP/EEA Emission Inventory Guidebook 2023 (EMEP/EEA, 2023). As regards the first two groups, methods are used leading to high standard detailed emissions data.

The first group includes: CO, NO_x, VOC, CH₄, NMVOC, N₂O, NH₃ and PM. For these pollutants, specific emission factors are applied relating to different engine conditions and urban, rural and highway driving shares.

The second group includes: CO₂, SO₂, Pb, Cd, Cr, Cu, Ni, Se, Zn. The emissions of these pollutants are estimated on the basis of fuel consumption.

For the third group of pollutants, including PAHs and PCDDs and PCDFs, detailed data are not available and then a simplified methodology is applied.

Finally, the fourth group includes pollutants (alkanes, alkenes, alkynes, aldehydes, ketones, cycloalkanes and aromatic compounds) obtained as a fraction of the total emissions of NMVOC, assuming that the fraction of residual NMVOC are PAHs.

Because of the availability in Italy of an extensive and accurate database, a detailed methodology is implemented in the model COPERT 5. Total emissions are calculated as the sum of hot emissions, deriving from the engine when it reaches a hot temperature, and cold emissions produced during the heating process. The different methodological approach is justified by the performance of vehicles in the two different phases. The production of emissions is also closely linked to the driving mode, differentiating for activity data and emission factors, with reference to urban (where it is assumed that almost all cold emissions are produced), rural and highway shares. Several factors contribute to the production of hot emissions such as mileage, speed, type of road, vehicle age, engine capacity and weight. Cold emissions are mainly attributed to urban share, and are attributed only to passenger cars and light duty vehicles. Varying according to the weather conditions and driving behaviour, are related to the specific country.

Emissions of NMVOC, NO_x, CO and PM are calculated on the basis of emission factors expressed in grams per kilometre and road traffic statistics estimated by ISPRA on account of data released from Ministry of Transport, ACI and ANCMA (several years). The emission factors are based on experimental measurements of emissions from in-service vehicles of different types driven under test cycles with different average speeds calculated from the emission functions and speed-coefficients provided by COPERT 5 (EMISIA SA, 2024). This source provides emission functions and coefficients relating emission factors (in g/km) to average speed for each vehicle type and Euro emission standard derived by fitting experimental measurements to polynomial functions. These functions were then used to calculate emission factor values for each vehicle type and Euro emission standard at each of the average speeds of road and area types. As regards the speciation of PM into elemental (EC, assumed to be equal to black carbon for road transport) and organic carbon (OC), considering the organic material (OM) as the mass of organic carbon corrected for the hydrogen content of the compounds collected, since the estimates are based on the assumption that low-sulphur fuels are used, when advanced after treatments are used, EC and OM do not add up to 100%, assuming that the remaining fraction consists of ash, nitrates, sulphates, water and ammonium salts (EMEP/EEA 2023).

Emissions of fuel dependent pollutants have been estimated applying a different approach. Data on consumption of various fuels are derived from official statistics aggregated at national level and then estimated in the detail of vehicle categories, emission regulation and road type in Italy. The resulting error of approximation deriving from the comparison between the calculated value and the statistical value of the total fuel consumption, is corrected by applying a normalisation procedure to the breakdown of fuel consumption by each vehicle type calculated on the basis of the fuel consumption factors added up, with reference to the BEN figures for total fuel consumption in Italy (adjusted for off-road consumption). The 1990-2022 inventory uses fuel consumption factors expressed as MJ per kilometre for each vehicle type and average speed calculated from the emission functions and speed-coefficients provided by the model COPERT 5, version 5.7.3. Emissions of sulphur dioxide and heavy metals are calculated applying specific factors to consumption of gasoline, diesel, liquefied petroleum gas (LPG) and natural gas (CNG), taken from the BEN (MASE, several years (a)), updated since 2017 according to EUROSTAT methodology.

Emissions of SO₂ are based on the sulphur content of the fuel. Values for SO₂ vary annually as the sulphur-content of fuels change and are calculated every year for gasoline and gas oil and officially communicated to the European Commission in the framework of the European Directives on fuel quality; these figures are also published by the refineries industrial association (UNEM, several years). Fuel specifications for gasoline, diesel fuel and LPG, derive from ad hoc studies about the properties of transportation fuels sold in Italy and whose results are representative and applicable with reference to three different time phases: 1990 – 1999; 2000 – 2012; 2013 – 2019; since 2020 (Innovhub – Fuel Experimental Station surveys, several years). In particular the results obtained for biodiesel in 2020 have been used to update the entire fuel specifications historical series (previously, in the absence of country specific parameters, Eurostat Energy Balance parameters were used).

As regards natural gas, the national market is characterized by the commercialisation of gases with different chemical composition in variable quantities from one year to the other. Each year the quantities of natural gas imported or produced in Italy are published on the web: <https://dgsaie.mise.gov.it/bilancio-gas-naturale> (Ministry of Environment, Ministry of Industry, several years).

In Italy, biodiesel, biogasoline and biogas are used in road transportation and the respective emissions have been estimated in the Inventory. As regards biodiesel and biogasoline, almost all the commercial gasoline is practically still substantially an E0 (in 2022 the share of biogasoline is 0.4%, respect to the total road gasoline consumption), while the distributed diesel reaches up to 5-7% by volume of biodiesel in diesel fuel (in 2022 the share of biodiesel is 5.8%, respect to the total road diesel consumption). That is because Italian producers/refineries have decided since the beginning of the introduction of the obligations on biofuels to focus on biodiesel rather than on ethanol to comply with the European/Italian obligations to introduce bio-fuels on the market. Biogasoline represents to date a minimum percentage out of the total gasoline including biogasoline consumption. According to the Renewable energy Directive (2009/28/EC) the amount of biogasoline reported in the Energy balance is equal to the renewable part of the fuel, calculated as the 37% of the total volume placed on the market. Biodiesel has been tested since 1994 to 1996 before entering in production since 1998. Moreover biogas road consumption has been taken into account in the Inventory, representing about 11%, 15% and 26% of total road natural gas consumption in 2020, 2021 and 2022 respectively. It is reported since year 2020 in the IEA - Eurostat – UNECE Energy Questionnaire (as element not covered by Regulation EC No 1099/2008 on energy statistics, therefore it is not mandatory for Countries to transmit it to Eurostat).

Emissions of heavy metals are estimated on the basis of data regarding the fuel and lubricant content and the engine wear; as reported in the EMEP/EEA Emission Inventory Guidebook 2023, these apparent fuel metal content factors originate from the work of Winther and Slentø, 2010, and have been reviewed by the TFEIP expert panel in transport and because of the scarce available information, the uncertainty in the estimate of these values is still considered quite high. In COPERT model heavy metals emission factors have been then updated focusing on the distinction between exhaust and non exhaust share. Non exhaust emissions of PAHs have also been estimated on the basis of brake and tyres debris-bound values resulting from the EMEP/EEA guidebook 2023.

3.5.2.2 *Evaporative emissions*

As regards NMVOC, the share of evaporative emissions is provided. These emissions are calculated only for gasoline vehicles: passenger cars, light duty vehicles, mopeds and motorcycles. Depending on temperature and vapour pressure of fuel, evaporative emissions have shown a growth over the years, nevertheless recently the contribution has been reduced by the introduction of control systems such as the canister. The estimation procedure is differentiated according to the processes of diurnal emission, running losses and hot soak emissions (EMEP/EEA, 2023).

3.5.2.3 *Emissions from automobile tyre and brake wear*

Not exhaust PM emissions from road vehicle tyre and brake wear are estimated. The focus is on the primary particles, deriving directly from tyre and brake wear. The material produced by the effects of wear and attrition between surfaces is subject to evaporation at high temperatures developed by the contact. Emissions are influenced by, as regards tyres, composition and pressure of tyres, structure and characteristics of vehicles, the peculiarities of the road and, as regards brakes, by the composition of the materials of the components, the position, the configuration systems, and the mechanisms of actuation (EMEP/EEA, 2023).

3.5.2.4 *Emissions from automobile road abrasion*

Particulate non-exhaust emissions deriving from road surface wear have been introduced in COPERT model, according to the Guidebook methodology (EMEP/EEA, 2023). Emissions depend on the type of asphalt-based and concrete-based road surfaces, taking into account that composition can vary widely, both from country to country and within countries. The type of tyres used also affect emissions, for instance the wear of the road surface, and the resulting PM concentrations due to resuspension, are considerably high when studded tyres are extensively used during the winter. The wear of the road surface increases with moisture level, also increasing after salting of the road, since the surface remains wet for longer periods. Other influencing factors are vehicle speed, tyre pressure and air temperature. As a consequence of the decrease of temperature, tyres become less elastic, causing the increase of the road surface wear rates (EMEP/EEA, 2023).

3.5.3 Activity data

The road traffic data used are vehicle-kilometre estimates for the different vehicle types and different road classifications in the national road network. These data have to be further broken down by composition of each vehicle fleet in terms of the fraction of different fuels types powered vehicles on the road and in terms of the fraction of vehicles on the road set by the different emission regulations which applied when the vehicle was first registered. These are related to the age profile of the vehicle fleet.

Basic data derive from different sources. Detailed data on the national fleet composition are found in the yearly report from ACI (ACI, several years), used from 1990 to 2006, except for mopeds for which estimates have been elaborated, for the whole time series, on the basis of National Association of Cycle-Motorcycle Accessories data on mopeds fleet composition and mileages (ANCMA, several years).

The Ministry of Transport provides specific fleet composition data for all vehicle categories from 2007 onwards, starting from 2013 submission. The Ministry of Transport in the national transport yearbook (MIT, several years) reports mileages time series. Furthermore since 2015 Ministry of Transport supplies information relating the distribution of old gasoline cars over the detailed vehicles categories (PRE ECE; ECE 15/00-01; ECE 15/02; ECE 15/03; ECE 15/04; information obtained from the registration year; data used for the updating of the time series since 2007). Ministry of Transport data are used relating to: the passenger cars (battery electric, petrol hybrid and diesel hybrid passenger cars are introduced from 2007 onwards, the detailed "Gasoline < 0.8 l" passenger cars subsector is introduced since 2012 and "Diesel < 1.4 l" subsector since 2007 onwards, Petrol PHEV passenger cars are introduced since 2013, in addition to the gasoline, diesel, LPG, CNG traditional ones); the diesel and gasoline light commercial vehicles; the breakdown of the heavy duty trucks, buses and coaches fleet according to the different weight classes and fuels (diesel almost exclusively for HDT, a negligible share consists of gasoline vehicles; diesel for coaches; diesel, diesel hybrid and CNG for urban buses); the motorcycles fleet in the detail of subsector

and legislation standard of both 2-stroke and 4-stroke categories. Fleet values for mopeds are updated according to the revisions of data published by ANCMMA and Ministry of Transport data; fleet values for diesel buses are updated according to the updating of the data on urban public buses, published on CNIT. The National Institute of Statistics carries out annually a survey on heavy goods vehicles, including annual mileages (ISTAT, several years). The National Association of concessionaries of motorways and tunnels produces monthly statistics on highway mileages by light and heavy vehicles (AISCAT, several years). The National General Confederation of Transport and Logistics (CONFETRA, several years) and the national Central Committee of road transporters (Giordano, 2007) supplied useful information and statistics about heavy goods vehicles fleet composition and mileages.

Fuel consumption data derive basically from the National Energy Balance (MASE, several years (a)); supplementary information is taken from the Oil Bulletin (MASE, several years (b)). As regards biofuels, the consumption has increased in view of the targets to be respected by Italy and set in the framework of the European directive 20-20-20. The trend of biodiesel is explained by the fact that this biofuel has been tested since 1994 to 1996 before entering in production since 1998. The consumption of bioethanol is introduced since 2008, according to data resulting on the BEN. Biogas is introduced since 2020 in the IEA - Eurostat – UNECE Energy Questionnaire.

Emissions are calculated from vehicles of the following types:

- Gasoline passenger cars
- Diesel passenger cars
- LPG passenger cars
- CNG passenger cars
- Petrol Hybrid passenger cars
- Petrol PHEV passenger cars
- Diesel Hybrid passenger cars
- Battery electric passenger cars
- Gasoline Light Commercial Vehicles (Gross Vehicle Weight (GVW) <= 3.5 tonnes)
- Diesel Light Commercial Vehicles (Gross Vehicle Weight (GVW) <= 3.5 tonnes)
- Rigid-axle Heavy Duty Trucks (GVW > 3.5 tonnes)
- Articulated Heavy Duty Trucks (GVW > 3.5 tonnes)
- Diesel Buses and coaches
- Diesel Hybrid Buses
- CNG Buses
- Mopeds and motorcycles.

In Table 3.16 the historical series of annual consumption data (Mg) for the different fuel types is reported.

Table 3.16 Annual fuel consumption data (Mg)

Fuel	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gasoline Leaded	12,280,212	10,112,250	4,542,113	-	-	-	-	-	-	-
Gasoline Unleaded	639,115	7,060,391	12,175,814	13,482,132	9,806,890	7,809,940	7,353,002	5,797,395	7,024,957	7,860,951
Diesel	15,278,022	14,445,441	17,059,010	22,327,864	21,557,266	21,128,587	20,985,104	17,103,910	21,033,279	21,799,554
LPG	1,342,000	1,478,000	1,422,000	1,029,000	1,214,000	1,654,000	1,653,017	1,309,461	1,406,968	1,535,280
CNG	183,770	216,804	292,214	342,756	610,502	787,148	832,667	671,443	795,900	613,184
of which biogas								71,340	119,044	160,953
Biodiesel	-	44,491	64,723	200,000	1,468,086	1,292,079	1,409,548	1,408,889	1,571,059	1,532,236
Biogasoline	-	-	-	-	142,106	30,420	35,401	22,841	31,524	40,718

Source: ISPRA elaborations on BEN, BP, UNEM data

The final reports on the physic-chemical characterization of fossil fuels used in Italy, carried out by the Fuel Experimental Station, that is an Italian Institute operating in the framework of the Department of Industry, are used with the aim to improve fuel quality specifications (surveys conducted in 2000, in 2012

– 2013 and in 2020). Fuel information has also been updated for the entire time series on the basis of the annual reports published by ISPRA about the fuel quality in Italy. Monitoring of the carbon content of the fuels used in Italy is an ongoing activity at ISPRA (Italian Institute for Environmental Protection and Research). The purpose is to analyse regularly the chemical composition of the used fuels or relevant commercial statistics to estimate the carbon content/emission factor (EF) of the fuels. With reference to the whole inventory, for each primary fuel, a specific procedure has been established.

As regards road transport, Italy fuel specifications values for gasoline, diesel fuel and LPG, derive from Fuel Experimental Station analysis about the properties of transportation fuels sold in Italy and whose results are representative and applicable with reference to four different time phases: 1990 – 1999; 2000 – 2012; 2013 – 2019; since 2020 (Innovhub – Fuel Experimental Station surveys, several years). As regards natural gas, the national market is characterized by the commercialisation of gases with different chemical composition in variable quantities from one year to the other. The methodology used to estimate the average EF for natural gas per year is based on the available consumption data, referring to the lower heat value (each year the quantities of natural gas imported or produced in Italy are published on the web by the Ministry of Environment (<https://dgsaie.mise.gov.it/bilancio-gas-naturale>). A normalisation procedure is applied to ensure that the breakdown of fuel consumption by each vehicle type calculated on the basis of the fuel consumption factors then added up matches the BEN figures for total fuel consumption in Italy (adjusted for off-road consumption). The automatic energy balance process, introduced by COPERT 5, has been applied. The simulation is started up having the target to equalize calculated and statistical consumptions, separately for fuel, at national level, with the aim to obtain final estimates the most accurate as possible. Once all data and input parameters have been inserted and all options have been set reflecting the peculiar situation of the Country, emissions and consumptions are calculated by the model in the detail of the vehicle category legislation standard; then the aggregated consumption values so calculated are compared with the input statistical national aggregated values (deriving basically from the National Energy Balance, as described above), with the aim to minimize the deviation.

In the following Table 3.17, Table 3.18, Table 3.19 and Table 3.20 detailed data on the relevant vehicle mileages in the circulating fleet are reported, subdivided according to the main emission regulations (ISPRA elaborations on ACI, ANCMA and Ministry of Transport data).

Table 3.17 Passenger Cars technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
a. Gasoline cars technological evolution										
PRE ECE, pre-1973	0.04	0.03	0.01	0.01	0.00	0.00	0.00	0.00	0.00	0.00
ECE 15/00-01, 1973-1978	0.10	0.04	0.01	0.00	0.00	0.00	0.00	0.00	0.00	0.00
ECE 15/02-03, 1978-1984	0.30	0.15	0.03	0.01	0.01	0.01	0.01	0.01	0.00	0.00
ECE 15/04, 1985-1992	0.55	0.55	0.28	0.10	0.04	0.03	0.02	0.02	0.02	0.02
PC Euro 1 - 91/441/EEC, from 1/1/93	0.00	0.24	0.27	0.17	0.05	0.02	0.01	0.01	0.01	0.01
PC Euro 2 - 94/12/EEC, from 1/1/97	-	-	0.39	0.32	0.21	0.11	0.07	0.06	0.05	0.04
PC Euro 3 - 98/69/EC Stage2000, from 1/1/2001	-	-	-	0.31	0.20	0.15	0.09	0.08	0.07	0.07
PC Euro 4 - 98/69/EC Stage2005, from 1/1/2006	-	-	-	0.09	0.44	0.41	0.28	0.27	0.25	0.24
PC Euro 5 - EC 715/2007, from 1/1/2011	-	-	-	-	0.04	0.21	0.18	0.17	0.17	0.17
PC Euro 6 (Since EC 715/2007, from 9/1/2015)										
- Euro 6 a/b/c	-	-	-	-	-	0.06	0.25	0.24	0.23	0.23
- Euro 6 d-temp (2019 - 2020)	-	-	-	-	-	-	0.09	0.14	0.14	0.13
- Euro 6 d (since 2021)	-	-	-	-	-	-	-	-	0.05	0.09
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
b. Diesel cars technological evolution										

f. Petrol PHEV cars technological evolution (from 2013 onwards)											
- Euro 6 a/b/c			1.00	1.00	1.00	0.14	0.08	0.05	0.03		
- Euro 6 d-temp (2017-2019)			-	-	-	0.86	0.41	0.24	0.16		
- Euro 6 d (2020 and later)			-	-	-	-	0.50	0.71	0.81		
Total			1.00	1.00	1.00	1.00	1.00	1.00	1.00		
			2007	2008	2009	2010	2015	2019	2020	2021	2022
g. Hybrid Diesel cars technological evolution (from 2007 onwards)											
PC Euro 6 (Since EC 715/2007, from 9/1/2015)											
- Euro 6 a/b/c			1.00	1.00	1.00	1.00	1.00	0.11	0.04	0.02	0.01
- Euro 6 d-temp (2017-2019)			-	-	-	-	-	0.89	0.37	0.17	0.11
- Euro 6 d (2020 and later)			-	-	-	-	-	-	0.59	0.81	0.88
Total			1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

Source: ISPRA elaborations on ACI and Ministry of Transport data

Table 3.18 Light Commercial Vehicles technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
a. Gasoline Light Commercial Vehicles technological evolution										
Conventional, pre 10/1/94	1.00	0.93	0.63	0.35	0.08	0.05	0.03	0.03	0.03	0.03
LCV Euro 1 - 93/59/EEC, from 10/1/94	-	0.07	0.21	0.17	0.10	0.05	0.02	0.02	0.02	0.02
LCV Euro 2 - 96/69/EEC, from 10/1/98	-	-	0.16	0.15	0.30	0.17	0.06	0.05	0.04	0.04
LCV Euro 3 - 98/69/EC Stage2000, from 1/1/2002	-	-	-	0.31	0.26	0.20	0.11	0.10	0.07	0.08
LCV Euro 4 - 98/69/EC Stage2005, from 1/1/2007	-	-	-	0.01	0.25	0.31	0.24	0.23	0.20	0.17
LCV Euro 5 - 2008 Standards 715/2007/EC, from 1/1/2012	-	-	-	-	0.00	0.21	0.17	0.16	0.15	0.13
LCV Euro 6 (Since 2007/715/EC, from 9/1/2016)										
- Euro 6 a/b/c	-	-	-	-	-	0.02	0.11	0.10	0.09	0.08
- Euro 6 d-temp (2018 - 2020)	-	-	-	-	-	-	0.25	0.30	0.31	0.26
- Euro 6 d (since 2021)	-	-	-	-	-	-	-	-	0.09	0.20
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
b. Diesel Light Commercial Vehicles technological evolution										
Conventional, pre 10/1/94	1.00	0.92	0.54	0.23	0.07	0.03	0.01	0.01	0.01	0.01
LCV Euro 1 - 93/59/EEC, from 10/1/94	-	0.08	0.22	0.11	0.05	0.03	0.01	0.01	0.00	0.00
LCV Euro 2 - 96/69/EEC, from 10/1/98	-	-	0.24	0.20	0.18	0.09	0.02	0.01	0.01	0.01
LCV Euro 3 - 98/69/EC Stage2000, from 1/1/2002	-	-	-	0.45	0.34	0.22	0.06	0.05	0.05	0.04
LCV Euro 4 - 98/69/EC Stage2005, from 1/1/2007	-	-	-	0.01	0.34	0.33	0.19	0.18	0.18	0.16
LCV Euro 5 - 2008 Standards 715/2007/EC, from 1/1/2012	-	-	-	-	0.01	0.30	0.32	0.29	0.26	0.24
LCV Euro 6 (Since 2007/715/EC, from 9/1/2016)										
- Euro 6 a/b/c	-	-	-	-	0.00	0.01	0.17	0.16	0.14	0.13
- Euro 6 d-temp (2018 - 2020)	-	-	-	-	-	-	0.22	0.29	0.26	0.25
- Euro 6 d (since 2021)	-	-	-	-	-	-	-	-	0.09	0.16
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

Table 3.19 Heavy Duty Trucks and Buses technological evolution: circulating fleet calculated as stock data multiplied by actual mileage (%)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
a. Heavy Duty Trucks technological evolution										
Conventional, pre 10/1/93	1.00	0.90	0.68	0.39	0.20	0.02	0.02	0.01	0.01	0.01
HDT Euro I - 91/542/EEC Stage I, from 10/1/93	-	0.10	0.10	0.06	0.04	0.01	0.01	0.00	0.00	0.00
HDT Euro II - 91/542/EEC Stage II, from 10/1/96	-	-	0.22	0.27	0.15	0.08	0.05	0.04	0.04	0.04
HDT Euro III - 2000 Standards, 99/96/EC, from 10/1/2001	-	-	-	0.27	0.36	0.34	0.23	0.20	0.18	0.17
HDT Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006	-	-	-	-	0.07	0.09	0.07	0.07	0.06	0.06
HDT Euro V - 2008 Standards, 99/96/EC, from 10/1/2009	-	-	-	-	0.18	0.39	0.33	0.32	0.30	0.28
HDT Euro VI (Since 2009/595/EC, from 12/31/2013)										
- Euro VI A/B/C	-	-	-	-	-	0.07	0.24	0.24	0.22	0.21
- Euro VI D/E (2019 and later)	-	-	-	-	-	-	0.06	0.12	0.18	0.24
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
b. Diesel Buses technological evolution										
	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Conventional, pre 10/1/93	1.00	0.93	0.65	0.34	0.13	0.01	0.00	0.00	0.00	0.00
Buses Euro I - 91/542/EEC Stage I, from 10/1/93	-	0.07	0.07	0.08	0.04	0.01	0.00	0.01	0.00	0.00
Buses Euro II - 91/542/EEC Stage II, from 10/1/96	-	-	0.28	0.32	0.27	0.14	0.09	0.07	0.06	0.05
Buses Euro III - 2000 Standards, 99/96/EC, from 10/1/2001	-	-	-	0.26	0.34	0.38	0.25	0.26	0.21	0.20
Buses Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006	-	-	-	-	0.12	0.13	0.11	0.11	0.10	0.09
Buses Euro V - 2008 Standards, 99/96/EC, from 10/1/2009	-	-	-	-	0.11	0.28	0.26	0.25	0.26	0.26
Buses Euro VI (Since 2009/595/EC, from 12/31/2013)	-	-	-	-	-	0.05	0.21	0.20	0.21	0.20
- Euro VI A/B/C	-	-	-	-	-	-	0.06	0.10	0.15	0.19
- Euro VI D/E (2019 and later)	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
c. CNG Buses technological evolution										
	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Urban CNG Buses Conventional, pre 10/1/93; Urban CNG Buses Euro I - 91/542/EEC Stage I, from 10/1/93	1.00	1.00	0.11	0.01	0.00	0.00	0.00	0.00	0.00	0.00
Urban CNG Buses Euro II - 91/542/EEC Stage II, from 10/1/96	-	-	0.89	0.20	0.10	0.05	0.03	0.02	0.01	0.01
Urban CNG Buses Euro III - 2000 Standards, 99/96/EC, from 10/1/2001; Urban CNG Buses Euro IV - 2005 Standards, 99/96/EC, from 10/1/2006	-	-	-	0.79	0.09	0.07	0.06	0.05	0.04	0.04
Euro V - 2008 Standards, 99/96/EC, from 10/1/2009; EEV (Enhanced environmentally friendly vehicle; ref. 2001/27/EC and 1999/96/EC line C, optional limit emission values); Urban CNG Buses Euro VI - EC 595/2009, from 12/31/2013	-	-	-	-	0.81	0.88	0.91	0.93	0.94	0.95
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00
d. Diesel Hybrid Buses technological evolution (from 2007 onwards)										
	2007	2008	2009	2010	2015	2019	2020	2021	2022	
Buses Euro VI (Since 2009/595/EC, from 12/31/2013)										
- Euro VI A/B/C	1.00	1.00	1.00	1.00	1.00	0.21	0.16	0.10	0.08	
- Euro VI D/E	-	-	-	-	-	0.79	0.84	0.90	0.92	
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	

Table 3.20 Mopeds and Motorcycles technological evolution: stock data multiplied by actual mileage (%)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Mopeds and motorcycles										
- Conventional	1.00	1.00	0.88	0.43	0.18	0.12	0.11	0.09	0.09	0.09
Mopeds and motorcycles										
- Euro 1	-	-	0.12	0.30	0.20	0.14	0.11	0.10	0.09	0.09
Mopeds and motorcycles										
- Euro 2	-	-	-	0.22	0.35	0.32	0.23	0.22	0.20	0.19
Mopeds and motorcycles										
- Euro 3	-	-	-	0.04	0.27	0.41	0.41	0.40	0.38	0.36
Mopeds and motorcycles										
- Euro 4	-	-	-	-	-	-	0.14	0.18	0.18	0.17
Mopeds and motorcycles										
- Euro 5	-	-	-	-	-	-	0.00	0.01	0.05	0.10
Total	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00

Average emission factors are calculated for average speeds by three driving modes (urban, rural and motorway) combined with the vehicle kilometres travelled and vehicle categories. ISPRA estimates total annual vehicle kilometres for the road network in Italy by vehicle type, based on data from various sources:

- Ministry of Transport (MIT, several years) for rural roads and on other motorways; the latter estimates are based on traffic counts from the rotating census and core census surveys of ANAS (management authority for national road and motorway network);
- highway industrial association for fee-motorway (AISCAT, several years);
- local authorities for built-up areas (urban).

Table 3.21 Evolution of fleet consistency and mileage

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
All passenger vehicles (including moto), total mileage (10 ⁹ veh-km/y)	350	412	454	453	447	443	442	341	422	439
Car fleet (10 ⁶)	27	30	32	34	36	37	40	40	40	40
Moto, total mileage (10 ⁹ veh-km/y)	30	41	42	42	34	32	26	24	25	26
Moto fleet (10 ⁶)	7	7	9	9	9	10	10	10	10	10
Goods transport, total mileage (10 ⁹ veh-km/y)	69	76	81	104	85	70	65	63	72	79
Truck fleet (10 ⁶), including LDV	2	3	3	4	5	5	5	5	5	5

Source: ISPRA elaborations

Notes: The passenger vehicles include passenger cars, buses and moto; the moto fleet includes mopeds and motorcycles; in the goods transport light commercial vehicles and heavy duty trucks are included.

3.5.4 Time series and key categories

The analysis of time series on transport data shows a trend that is the result of the general growth in mobility demand and consumptions until 2007, followed by a decrease basically due to the economic crisis on one side, and of the introduction of advanced technologies limiting emissions in modern vehicles in recent years, on the other side; then the growth in the years 2018-2019 is followed by a sharp decrease in 2020 due to the pandemic crisis and a subsequent partial recovery from 2020 to 2021.

More in details, passenger cars and light duty vehicles emissions trends are driven by a gradual decrease over the years of gasoline fuel consumption balanced by an increase of diesel fuel which is the main driver for NO_x and PM emissions. At pollutant level emission trends are driven not only by fuel but also by changes in technologies which are reflected in the COPERT model by the evolution of the annual vehicle fleet. Due to the penetration of new vehicles with more stringent pollutant limits, some pollutant emissions decreased faster than other. An important role has been played also by the distribution between diesel and gasoline fuel consumptions. In the last years an increase of diesel fuelled vehicles and a decrease of gasoline ones have been registered and diesel fuel new technologies resulted in a slower

decrease of NO_x emission than expected. Regarding heavy duty vehicles emissions trends are explained by the variations estimated in mileages time series data correlated to the variations registered in fuel consumptions; annual variation is explained by the general trend of national economic growth and in particular commercial and industrial activities. Emissions trends regarding mopeds and motorcycles are explained by the variations estimated in mileages time series data correlated to the variations registered in gasoline consumptions. The annual penetration of new technologies explains annual emission trends. The general decrease between 2019 and 2020 is explained by the pandemic crisis.

In Table 3.22 the list of key categories by pollutant identified for road transport in 2022, 1990 and at trend assessment is reported.

Table 3.22 List of key categories for pollutant in the road transport in 2022, 1990 and in trend

	Key categories in 2022			Key categories in 1990			Key categories in trend		
NO _x	1A3bi	1A3bii	1A3biii	1A3bi	1A3bii	1A3biii	1A3bi	1A3bii	1A3biii
NMVOOC	1A3biv	1A3bv		1A3bi	1A3biv	1A3bv	1A3bi	1A3biv	
NH ₃							1A3bi		
CO	1A3bi	1A3biv		1A3bi	1A3biv		1A3bi		
PM10	1A3bvi	1A3bvii		1A3bi	1A3bii	1A3biii	1A3bvi	1A3bi	1A3bii
PM2.5	1A3bi	1A3bvi		1A3bi	1A3bii	1A3biii	1A3bvi	1A3bi	1A3bii
BC	1A3bi	1A3biii	1A3bvi	1A3bi	1A3bii	1A3biii		1A3bi	1A3bii
Pb	1A3bvi			1A3bi	1A3biv			1A3bi	1A3bvi
Cd	1A3bi	1A3bvi						1A3bi	

Source: ISPRA elaborations

In 2022 key categories are identified for the following pollutants: nitrogen oxides, non methane volatile organic compounds, carbon monoxide, particulate matter with diameter less than 10 µm, particulate matter with diameter less than 2.5 µm, black carbon, lead and cadmium.

Nitrogen oxides emissions show a decrease since 1990 of -74.8%. Emissions are mainly due to diesel vehicles. The decrease observed since 1990 in emissions relates to all categories except for diesel passenger cars, hybrid categories and CNG buses. In 2022, emissions of nitrogen oxides (Table 3.23) from passenger cars, light-duty vehicles and heavy-duty trucks including buses are key categories. The same categories are identified as key categories in 1990 and in trend.

Table 3.23 Time series of nitrogen oxides emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
1.A.3.b.i Passenger cars	590.9	628.3	388.2	239.4	178.8	169.6	146.6	105.2	125.1	124.5
1.A.3.b.ii Light-duty vehicles	60.6	69.5	64.0	75.3	54.8	48.8	39.8	36.5	38.2	40.2
1.A.3.b.iii Heavy-duty vehicles including buses	340.3	336.1	328.6	307.6	193.8	123.5	93.3	77.6	85.3	84.4
1.A.3.b.iv Mopeds and motorcycles	4.3	5.6	5.9	6.8	4.8	4.1	2.8	2.4	2.4	2.4
Total emissions	996.1	1,039.5	786.8	629.1	432.2	346.0	282.5	221.8	251.1	251.5

As regards non methane volatile organic compounds, emissions from mopeds and motorcycles and gasoline evaporation are key categories in 2022; emissions from passenger cars, mopeds and motorcycles and gasoline evaporation are key categories in 1990; emissions from passenger cars and mopeds and motorcycles are key categories in trend. Despite the decline of about -88.2% since 1990 of emissions of non-methane volatile organic compounds from this category, road transport (Table 3.24) is the fourth source at national level after the use of solvents, the not industrial combustion and agriculture; this trend is due to the combined effects of technological improvements that limit VOCs from tail pipe and evaporative emissions (for cars) and the expansion of two-wheelers fleet. In Italy there is in fact a remarkable fleet of motorbikes and mopeds (about 10 million vehicles in 2022) that uses gasoline and it is increased of about 56.5% since 1990 (this fleet not completely complies with strict VOC emissions controls).

Table 3.24 Time series of non methane volatile organic compounds emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
1.A.3.b.i Passenger cars	430.4	468.2	265.0	122.2	47.3	26.5	19.2	13.0	13.9	13.6
1.A.3.b.ii Light-duty vehicles	15.6	18.5	12.4	11.6	5.9	2.7	0.8	0.7	0.7	0.7
1.A.3.b.iii Heavy-duty vehicles including buses	26.3	25.2	21.7	17.4	8.6	3.7	2.8	2.3	2.6	2.6
1.A.3.b.iv Mopeds and motorcycles	175.5	248.8	225.2	152.2	73.0	50.0	29.8	25.1	25.1	25.2
1.A.3.b.v Gasoline evaporation	119.2	119.1	87.9	55.5	44.3	45.7	43.9	45.7	44.4	48.6
Total emissions	767.0	879.7	612.2	358.9	179.1	128.6	96.5	86.7	86.7	90.6

Source: ISPRA elaborations

Carbon monoxide emissions from passenger cars and mopeds and motorcycles are key categories in 2022 and 1990; passenger cars are also key category in trend. The time series of CO emissions is reported in Table 3.25.

Table 3.25 Time series of carbon monoxide emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
1.A.3.b.i Passenger cars	4,124.6	4,171.4	2,191.3	1,087.4	467.6	271.4	214.1	149.1	164.3	163.9
1.A.3.b.ii Light-duty vehicles	175.1	207.7	124.7	98.3	45.7	21.6	7.6	6.3	6.9	7.4
1.A.3.b.iii Heavy-duty vehicles including buses	81.4	79.8	74.2	70.4	48.7	36.3	28.6	24.5	27.1	26.9
1.A.3.b.iv Mopeds and motorcycles	493.4	667.3	619.3	433.8	195.7	141.0	98.1	81.6	81.9	82.1

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Total emissions	4,874.5	5,126.1	3,009.5	1,689.8	757.7	470.3	348.3	261.4	280.3	280.3

Source: ISPRA elaborations

A strong contribution to total emissions is given by gasoline vehicles (about 76.7% in 2022, although since 1990 a decrease of about -95.2% is observed); since 1990 to 2022 a general decrease, of about -94.2%, is observed.

Emissions of PM10 (Table 3.26) deriving from passenger cars, light-duty vehicles, heavy-duty vehicles including buses, road vehicle tyre and brake wear are key categories in 1990; emissions from road vehicle tyre and brake wear and emissions from road surface wear are key categories in 2022; emissions from passenger cars, light-duty vehicles, heavy-duty vehicles including buses and from road vehicle tyre and brake wear are key category in trend.

As regards PM2.5 (Table 3.27), in 2022 emissions from passenger cars and from road vehicle tyre and brake wear are key categories; emissions from passenger cars, light-duty vehicles, heavy-duty vehicles including buses and road vehicle tyre and brake wear are key categories in 1990 and in trend.

With regard to particulate matter, the relative weight of the non-exhaust component in total emissions becomes progressively more significant over the years.

Table 3.26 Time series of particulate matter with diameter less than 10 µm emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
1.A.3.b.i Passenger cars	19.3	14.1	11.9	9.8	8.5	5.8	3.7	2.5	2.6	2.8
1.A.3.b.ii Light-duty vehicles	10.4	11.5	9.2	8.9	4.8	2.3	0.8	0.6	0.7	0.6
1.A.3.b.iii Heavy-duty vehicles including buses	13.6	13.3	11.4	9.1	4.7	2.3	1.6	1.3	1.5	1.4
1.A.3.b.iv Mopeds and motorcycles	3.7	5.3	4.8	3.2	1.4	1.0	0.6	0.5	0.5	0.5
1.A.3.b.vi Road Transport, Automobile tyre and brake wear	8.9	10.1	11.0	12.0	11.4	11.4	10.0	8.2	9.9	10.4
1.A.3.b.vii Road transport: Automobile road abrasion	4.1	4.6	5.0	5.4	4.8	4.5	4.6	3.8	4.6	4.8

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Total emissions	59.9	58.9	53.4	48.4	35.6	27.3	21.4	16.8	19.7	20.6

Source: ISPRA elaborations

Table 3.27 Time series of particulate matter with diameter less than 2.5 µm emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
1.A.3.b.i Passenger cars	19.3	14.1	11.9	9.8	8.5	5.8	3.7	2.5	2.6	2.8
1.A.3.b.ii Light-duty vehicles	10.4	11.5	9.2	8.9	4.8	2.3	0.8	0.6	0.7	0.6
1.A.3.b.iii Heavy-duty vehicles including buses	13.6	13.3	11.4	9.1	4.7	2.3	1.6	1.3	1.5	1.4
1.A.3.b.iv Mopeds and motorcycles	3.7	5.3	4.8	3.2	1.4	1.0	0.6	0.5	0.5	0.5
1.A.3.b.vi Road Transport Automobile tyre and brake wear	4.6	5.3	5.7	6.2	5.9	5.9	5.3	4.3	5.2	5.5
1.A.3.b.vii Road transport Automobile road abrasion	2.2	2.5	2.7	2.9	2.6	2.4	2.5	2.0	2.5	2.6
Total emissions	53.7	51.9	45.8	40.1	27.9	19.6	14.5	11.2	12.9	13.4

Source: ISPRA elaborations

Emissions of particulate matter with diameter less than 10µm and less than 2.5µm show a decreasing trend since 1990 respectively of about -65.6% and -75.0%. Despite the decrease, diesel vehicles (passenger cars, light duty vehicles and heavy duty trucks including buses) are mainly responsible for road transport emissions giving a strong contribution to total emissions, in 2022 about 76.0% and 78.4% out of the total for PM10 and PM2.5 respectively.

Emissions of black carbon are reported in Table 3.28. Emissions from passenger cars, heavy-duty trucks including buses and road vehicle tyre and brake wear are key categories in 2022; emissions from passenger cars, light-duty vehicles and heavy-duty trucks including buses are key categories in 1990; emissions from passenger cars, light-duty vehicles and heavy-duty trucks including buses and road vehicle tyre and brake wear are key categories in trend. The emissions trend is generally decreasing (-78.9% since 1990). The main contribution to total emissions is given by diesel vehicles, in 2022 equal to 90.3% out of the total. Despite of the decrease, road transport is the second source of emissions (the main source is non industrial combustion) at national level in 2022 (28.6%).

Table 3.28 Time series of black carbon emissions in road transport (Gg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Gg									
1.A.3.b.i Passenger cars	10.4	7.6	7.3	7.3	7.0	4.8	3.1	2.0	2.1	2.2
1.A.3.b.ii Light-duty vehicles	5.7	6.4	5.5	5.9	3.6	1.8	0.6	0.5	0.5	0.5
1.A.3.b.iii Heavy-duty vehicles including buses	6.8	6.8	6.0	5.2	2.9	1.5	1.1	0.9	1.0	1.0
1.A.3.b.iv Mopeds and motorcycles	0.7	0.9	0.9	0.6	0.2	0.2	0.1	0.1	0.1	0.1
1.A.3.b.vi Road Transport Automobile tyre and brake wear	1.1	1.2	1.3	1.4	1.3	1.3	1.2	1.0	1.2	1.3
1.A.3.b.vii Road transport Automobile road abrasion	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Total emissions	24.7	23.0	21.0	20.5	15.1	9.7	6.2	4.6	5.0	5.2

Source: ISPRA elaborations

Emissions of cadmium are reported in Table 3.29. Cadmium emissions from passenger cars are key categories in 2022 and in trend. Emissions from automobile tyre and brake wear are also key categories in 2022. Emissions show an increase since 1990 of about +4.1%, representing in 2022 the 13.0% of the national total. In 2022 most of the emissions derive from passenger cars (72.6%); non exhaust emissions from automobile tyre and brake wear are equal to 32.4% of the total.

Table 3.29 Time series of Cadmium emissions in road transport (Mg)

Source categories for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	Mg									
1.A.3.b.i Passenger cars	0.2	0.2	0.3	0.3	0.3	0.3	0.3	0.2	0.3	0.3
1.A.3.b.ii Light-duty vehicles	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.3.b.iii Heavy-duty vehicles including buses	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
1.A.3.b.iv Mopeds and motorcycles	0.1	0.1	0.1	0.1	0.1	0.1	0.0	0.0	0.0	0.0
1.A.3.b.vi Road Transport Automobile tyre and brake wear	0.1	0.2	0.2	0.2	0.2	0.2	0.2	0.1	0.2	0.2
1.A.3.b.vii Road transport Automobile road abrasion	NE	NE	NE	NE	NE	NE	NE	NE	NE	NE
Total emissions	0.5	0.6	0.6	0.7	0.6	0.6	0.5	0.4	0.5	0.5

Source: ISPRA elaborations

Emissions of SO_x, NH₃ and Pb are shown in Table 3.30. SO_x are not key categories. Pb emissions from road vehicle tyre and brake wear are key category in 2022, Pb emissions from passenger cars and from mopeds and motorcycles are key categories in 1990, Pb emissions from passenger cars and road vehicle tyre and brake wear are key categories in trend. Emissions of NH₃ from passenger cars are key categories in trend. In 2022 emissions of these pollutants deriving from road transport are less important compared to other sectors. Emissions of SO_x and Pb show strong decreases. Since 2002, due to limits on fuels properties imposed by legislation, Pb resulting emissions are almost completely non exhaust (road vehicle tyre and brake wear Pb emissions increase of about 13.1% since 1990); total Pb emissions decrease of -99.0% since 1990, representing in 2022 about 18.8% of the national total. SO_x emissions decrease by -99.7%, representing 0.5% of the total in 2022. Emissions of NH₃, despite the strong increase since 1990, in 2022 account for just 1.6% out of the total.

Table 3.30 Time series of sulphur oxides, ammonia and lead emissions in road transport

SO_x, NH₃, Pb Total Emissions for NFR Subsector 1.A.3.b	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
SO _x (Gg)	129.3	71.6	11.9	2.2	0.4	0.4	0.4	0.3	0.4	0.4
NH ₃ (Gg)	0.8	5.3	20.6	15.7	9.5	6.1	5.5	4.3	5.1	5.5
Pb (Mg)	3,806.2	1,644.8	719.8	46	43.6	44.6	37.3	30.2	36.6	38.4

3.5.5 QA/QC and Uncertainty

Data used for estimating emissions from the road transport sector, derive from different sources, including official statistics providers and industrial associations. A specific procedure undertaken for improving the inventory in the sector regards the establishment of a national expert panel in road transport which involves, on a voluntary basis, different institutions, local agencies and industrial associations cooperating for improving activity data and emission factors accuracy. In this group emission estimates are presented annually and new methodologies are shared and discussed. Reports and data of the meetings can be found at the following address: http://groupware.sinanet.isprambiente.it/expert_panel/library; 2023 meeting has been held in Livorno (<https://www.snambiente.it/snpa/arpa-lombardia/expert-panel-for-polluting-emissions-reduction/>). In addition, road transport emission factors are shared and publicly available on the website <https://fetransp.isprambiente.it/#/>.

Besides, over time recalculations of time series estimates have been discussed with national experts in the framework of an ad hoc working group on air emissions inventories. The group is chaired by ISPRA and includes participants from the local authorities responsible for the preparation of local inventories, sectoral experts, the Ministry of Environment and air quality model experts. Recalculations are comparable with those resulting from application of the model at local level. Top-down and bottom-up approaches have been compared with the aim at identifying the major problems and future possible improvements in the methodology to be addressed (emission estimates at the link: <https://emissioni.sina.isprambiente.it/inventario-nazionale/>). A Montecarlo analysis has been carried out by EMISIA on behalf of the Joint Research Centre (Kouridis et al., 2010) in the framework of the study "Uncertainty estimates and guidance for road transport emission calculations" for 2005 emissions. The study shows an uncertainty assessment, at Italian level, for road transport emissions on the basis of 2005 input parameters of the COPERT 4 model (v. 7.0).

3.5.6 Recalculation

The annual update of the emissions time series from road transport implies a periodic review process.

In 2024 submission the historical series has been revised mainly as a result of the upgrade of COPERT model version used (from version 5.6.1 in last submission to 5.7.3 in submission 2024), which resulted in various methodological updates.

Emission factors of Euro 6 CNG passenger cars, Euro VI diesel buses, Euro VI diesel hybrid buses, non-exhaust emission factors have been updated.

As regards the software, revisions relate: the removal of CO₂ correction tool, capability of alternative HDVs classification based on REG EU 2017/2400, improved labels of forms and headers.

Furthermore various bugs have been corrected regarding: the correction of the calculation for cold emissions of CO, NO_x, VOC for petrol and diesel PCs and LCVs, the correction of cold start ratio of diesel Euro 6 cars for NO_x and CO, the correction of cold start ratio of petrol-fueled cars and vans for VOC and CO, the correction of cold emissions of Euro 6 CNG passenger cars for SPN23, the correction of bug when importing stock of HDVs in VECTO groups from excel to COPERT. Software update issues and minor corrections have been also implemented.

Country specific emission factors for Euro 6 LPG passenger cars, deriving from Innovhub survey, until 2022 submission inserted in the model as user values, are now implemented in COPERT model.

N₂O emission factors for urban CNG buses have been introduced in COPERT model version 5.7.3, respect to the version 5.6.1 applied in last submission.

In consequence of the removal of CO₂ correction tool in COPERT model, previously referred to gasoline and diesel passenger cars from Euro 4 onwards, Country specific hot energy consumption factors have been applied (CNR STEMS, Innovhub SSI, 2020).

Italian road vehicles electricity consumption data derive from Eurostat database (<https://ec.europa.eu/eurostat/data/database>). COPERT input road electricity consumption data have been calibrated on the basis of the vehicles categories using electricity actually included in COPERT classification. In 2024 submission, updated electricity consumption factors have been applied, deriving from Appendix 4 of the 2023 EMEP/EEA air pollutant emission inventory guidebook, resulting in a different balance of mileage and in a revision of estimated electricity consumptions since 2007, respect to previous submission.

Revisions applied affected also the annual balances and, as a consequence, adjustments of mileages were applied in the historical series.

Natural gas estimates have been revised since 2021 according to the revision of road consumption data reported in the National Energy Balance and of parameters applied for the estimation of emissions from this fuel for the whole Inventory. In particular, in the National Energy Balance, road natural gas final consumption has been revised for 2021 respect to last submission; moreover, because of additional data availability for natural gas, density data for 2021 and H:C ratio since 2019 have been also updated.

Biogas road consumption, included in COPERT model in the total road natural gas consumption, has been considered in the Inventory, reported only since 2020 in the IEA - Eurostat – UNECE Energy Questionnaire (about 11%, 15% and 26% of total road natural gas consumption in 2020, 2021 and 2022 respectively).

Estimate of the mopeds circulating fleet in 2021 has been revised consistently with 2022 data supplied by the Ministry of Transport.

Mileage balance has been also revised in consequence of a revision of transport statistics data published by the Ministry of Transport regarding: the circulating urban buses fleet and the passengers-km since 2020; the tons-km as regards freight transport since 2021. In particular the most recent estimate of total on road freight transport data published for 2021, has been revised upwards from the Ministry of Transport respect to last submission and the estimate for 2022 results further higher, being the highest value than in last decade.

3.5.7 Planned improvements

Improvements for the next submission will be connected to the possible new availability of data and information regarding activity data, calculation factors and parameters, new developments of the methodology and the update of the software.

3.6 RAILWAYS (NFR SUBSECTOR 1.A.3.C)

The electricity used by the railways for electric traction is supplied from the public distribution system, so the emissions arising from its generation are reported under category 1.A.1.a Public Electricity. Emissions from diesel trains are reported under the IPCC category 1.A.3.c Railways. Estimates are based on the gasoil consumption for railways reported in BEN (MSE, several years [a]), updated since 2018 according to EUROSTAT methodology (<https://dgsaie.mise.gov.it/bilancio-energetico-nazionale>), and on the methodology Tier1, and emission factors from the EMEP/EEA Emission Inventory Guidebook 2023 (EMEP/EEA, 2023).

Fuel consumption data are collected by the Ministry of Economic Development, responsible of the energy balance, from the companies with diesel railways. The activity is present only in those areas without electrified railways, which are limited in the national territory. The trend reflects the decrease of the use of these railways. Because of low values, emissions from railways do not represent a key category. In Table 3.31 diesel consumptions (TJ) and nitrogen oxides, non-methane volatile organic compounds, Sulphur oxides, ammonia, particulate and carbon monoxide emissions (Gg) are reported. Emissions of Pb from 2002 are reported as 'NA', because of the introduction of unleaded liquid fuels in the market in 2002. In particular, heavy metals contents values derive from the analysis about the physical - chemical characterization of fossil fuels used in Italy (Innovhub, Fuel Experimental Station, several years).

Table 3.31 Consumptions and emissions time series in railways

Consumptions and Emissions for NFR Subsector 1.A.3.c	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Diesel Consumption (TJ)	8,370.3	8,199.4	5,850.6	4,142.4	2,690.4	939.6	1,836.3	1,851.7	1,601.0	475.2
Emissions from diesel trains (Gg)										
NO _x	10.27	10.06	7.18	5.08	3.24	1.00	1.75	1.72	1.49	0.44
NMVOC	0.91	0.89	0.64	0.45	0.29	0.09	0.17	0.17	0.14	0.04
SO _x	1.20	0.77	0.08	0.01	0.001	0.0003	0.001	0.001	0.0005	0.0002
NH ₃	0.001	0.001	0.001	0.001	0.0004	0.0002	0.0003	0.0003	0.0003	0.0001
PM _{2.5}	0.27	0.26	0.19	0.13	0.08	0.03	0.05	0.04	0.04	0.01
PM ₁₀	0.28	0.28	0.20	0.14	0.09	0.03	0.05	0.04	0.04	0.01
TSP	0.29	0.28	0.20	0.14	0.09	0.03	0.05	0.04	0.04	0.01
BC	0.18	0.18	0.13	0.09	0.06	0.02	0.03	0.03	0.02	0.01
CO	2.10	2.05	1.47	1.04	0.67	0.24	0.46	0.46	0.40	0.12

In the review process has been observed the existence of at least one steam engine still operating in Italy. It is an historic train used only for few days per year and probably fueled with biomass nowadays instead of coal. Nor biomass or coal are reported in the energy balance for railways activities. Anyway, this possible source of emission could be considered insignificant. No recalculation occurred in this submission. No specific improvements are planned for the next submission.

3.7 NAVIGATION (NFR SUBSECTOR 1.A.3.D)

3.7.1 Overview

This source category includes all emissions from fuels delivered to water-borne navigation. National navigation is a key category in 2022 with respect to emissions of SO_x, NO_x, PM₁₀, PM_{2.5} and BC.

3.7.2 Methodological issues

Emissions of the Italian inventory from the navigation sector are carried out according to the IPCC Guidelines and Good Practice Guidance (IPCC, 1997; IPCC, 2000) and the EMEP/EEA Guidebook (EMEP/EEA, 2023). In particular, a national methodology has been developed following the EMEP/EEA Guidebook which provides details to estimate emissions from domestic navigation, specifying recreational craft, ocean-going ships by cruise and harbour activities; emissions from international navigation are also estimated and included as memo item but not included in national totals (EMEP/EEA, 2023). Inland, coastal and deep-sea fishing are estimated and reported under 1.A.4.c. International inland waterways do not occur in Italy. The methodology developed to estimate emissions is based on the following assumptions and information.

Activity data comprise both fuel consumptions and ship movements, which are available in different level of aggregation and derive from different sources as specified here below:

- Total deliveries of fuel oil, gas oil and marine diesel oil to marine transport are given in national energy balance (MSE, several years (a)) but the split between domestic and international is not provided;
- Naval fuel consumption for inland waterways, ferries connecting mainland to islands and leisure boats, is also reported in the national energy balance as it is the fuel for shipping (MSE, several years (a));
- Data on annual arrivals and departures of domestic and international shipping calling at Italian harbors are reported by the National Institute of Statistics in the statistics yearbooks (ISTAT, several years (a)) and Ministry of Transport in the national transport statistics yearbooks (MIMS, several years).

As for emission and consumption factors, figures are derived by the EMEP/EEA guidebook (EMEP/EEA, 2019), both for recreational and harbor activities and national cruise, taking into account national specificities. These specificities derive from the results of a national study which, taking into account detailed information on the Italian marine fleet and the origin-destination movement matrix for the year 1997, calculated national values (ANPA, 2001; Trozzi et al., 2002 (b)) on the basis of the default emission and consumption factors reported in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). National average emissions and consumption factors were therefore estimated for harbor and cruise activities both for domestic and international shipping from 1990 to 1999. In 2009 submission the study was updated for the years 2004, 2005 and 2006 in order to consider most recent trends in the maritime sector both in terms of modelling between domestic and international consumptions and improvements of operational activities in harbor (TECHNE, 2009). On the basis of the results, national average emissions and consumption factors were updated from 2000. Specifically, for the years referred to in the surveys, the current method estimates emissions from the number of ships movements broken down by ship type at each of the principal Italian ports considering the information of whether the ship movement is international or domestic, the average tonnage and the relevant distance travelled. For those years, in fact, figures on the number of arrivals, destination, and fleet composition have been provided by the local port authorities and by the National Institute of Statistics (ISTAT, 2009), covering about 90% of the official national statistics on ship movements for the relevant years. Consumption and emission factors are those derived from the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007) and refer to the Tier 3 ship movement methodology that takes into account origin-destination ship movements matrices as well as technical information on the ships, as engine size, gross tonnage of ships and operational times in harbors. Based on sample information, estimates have been carried out at national level for the relevant years considering the official statistics of the maritime sector. Moreover, an update of the emission factor of hoteling phase has been carried out more recently and the emission factors have been updated accordingly (ISPRA, 2023 [c]).

In general, to carry out national estimates of greenhouse gases and other pollutants in the Italian inventory for harbor and domestic cruise activities, consumptions and emissions are calculated for the complete time series using the average consumption and emission factors multiplied by the total number of movements. On the other hand, for international cruise, consumptions are derived by difference from the total fuel consumption reported in the national energy balance and the estimated values as described above and emissions are therefore calculated. For maritime transportation only by Directive 1999/32/EC European Union started to examine environmental impact of navigation and in particular the Sulphur content of fuels. This directive was amended by Directive 2005/33/EC that designated the Baltic Sea, the English Channel and the North Sea as Sulphur emission control areas (SECA) limiting the content of Sulphur in the fuel for these areas and introducing a limit of 0.1% of the Sulphur content in the fuel used in EU harbors from 2010. EU legislation combined with national normative resulted in the introduction of a limit of Sulphur content in maritime gasoil equal to 0.2% (2% before) from 2002 and 0.1% from 2010 while for fuel oil some limits occur only from 2008 (maximum Sulphur content of 1.5 % in harbor) and from 2010, 2% in domestic waters and 1% in harbor. For inland waterways, which include the navigation on the Po river and ferry-boats in the Venice lagoon, the same legislation is applied. Moreover, since January 2020, IMO introduced a limit of Sulphur content in marine fuel oil equal to 0.5% (previously equal to 3.5%) in IMO jurisdiction areas (except in ECAS areas, which have a stricter Sulphur content limit, equal to 0.1%). For international navigation 0.3% of sulphur content in fuel oil has been assumed for the whole time series. Fuel oil accounts for about 98% of total international fuel consumption along the time series.

The composition of the fleet of gasoline fueled recreational craft distinguished in two strokes and four strokes engine distribution is provided by the industrial category association (UCINA, several years); the trend of the average emission factors takes into account the switch from two strokes to four strokes engines of the national fleet due to the introduction in the market of new models. In 2000, the composition of the fleet was 90% two stroke engines equipped and 10% four stroke while in the last year four strokes engines are about 56 % of the fleet. The fuel split between national and international fuel use in maritime transportation is then supplied to the Ministry of Economic Development to be included in the official international submission of energy statistics to the IEA in the framework of the Joint Questionnaire OECD/EUROSTAT/IEA compilation together with other energy data. A discrepancy with the international bunkers reported to the IEA still remains, especially for the nineties, because the time series of the energy statistics to the IEA are not updated. PCB, HCB and Dioxins emissions are estimated with Tier1 emission factors available in the 2023 EMEP/EEA Guidebook (EMEP/EEA, 2023).

3.7.3 Time series and key categories

In Table 3.32 the list of key categories by pollutant identified for navigation in 2022, 1990 and at trend assessment is reported. Navigation is, in 2022, key category for many pollutants: SO_x, NO_x, PM10, PM2.5, BC; furthermore, it is a key driver of the SO_x, NO_x and BC trend.

Table 3.32 List of key categories for pollutant in navigation in 2022, 1990 and in the trend

	Key categories in 2022	Key categories in 1990	Key categories in trend
SO _x	1A3dii	1A3dii	1A3dii
NO _x	1A3dii	1A3dii	1A3dii
PM10	1A3dii	1A3dii	
PM2.5	1A3dii	1A3dii	
BC	1A3dii		1A3dii

Source: ISPRA elaborations

Estimates of fuel consumption for domestic use, in the national harbors or for travel within two Italian destinations, and bunker fuels used for international travels are reported in Table 3.33. An upward trend

in emission levels is observed from 1990 to 2000, explained by the increasing number of ship movements. Nevertheless, the operational improvements in harbor activities and a reduction in ship domestic movements inverted the tendency in the last years.

Table 3.33 Marine fuel consumptions in domestic navigation and international bunkers (Gg) and pollutants emissions from domestic navigation (Gg)

Consumptions and Emissions for NFR Subsector 1.A.3.d	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gasoline for recreational craft (Gg)	182	210	213	199	169	90	87	86	84	66
Diesel oil for inland waterways (Gg)	20	23	83	77	62	54	24	19	22	21
Fuels used in domestic cruise navigation (Gg)	778	706	811	740	725	525	621	667	511	664
Fuel in harbors (dom+int ships) (Gg)	748	693	818	759	844	733	981	1,054	807	1,049
Fuel in international Bunkers (Gg)	1,348	1,241	1,214	2,095	2,562	1,563	1,702	1,321	1,774	1,331
Emissions of NO _x	95.55	87.97	105.16	97.15	100.30	78.33	95.40	102.11	78.89	101.70
Emissions of NMVOC	46.11	52.42	50.47	43.53	32.50	16.45	15.34	15.09	13.25	10.23
Emissions of SO _x	77.94	70.31	81.53	49.73	30.99	22.46	31.34	11.34	7.16	10.91
Emissions of PM2.5	9.30	8.83	9.89	9.14	8.77	6.01	7.85	8.34	6.36	7.87
Emissions of PM10	9.33	8.86	9.93	9.17	8.80	6.04	7.85	8.34	6.36	7.87
Emissions of BC	1.33	1.25	1.56	1.44	1.42	1.10	1.76	1.82	1.60	1.82
Emissions of CO	102.27	115.57	125.46	123.43	110.69	60.20	56.64	55.71	52.38	43.60

3.7.4 QA/QC and Uncertainty

Basic data to estimate emissions are reconstructed starting from information on ship movements and fleet composition coming from different sources. Data collected in the framework of the national study from the local port authorities, carried out in 2009 (TECHNE, 2009), were compared with the official statistics supplied by ISTAT, which are collected from maritime operators with a yearly survey and communicated at international level to EUROSTAT. Differences and problems were analyzed in details and solved together with ISTAT experts. Different sources of data are usually used and compared during the compilation of the annual inventory. Besides, time series resulting from the recalculation have been presented to the national experts in the framework of an ad hoc working group on air emissions inventories. The group is chaired by ISPRA and includes participants from the local authorities responsible for the preparation of local inventories, sectoral experts, the Ministry of Environment, Land and Sea, and air quality model experts. Top-down and bottom-up approaches have been compared with the aim to

identify the potential problems and future improvements to be addressed. There is also an ongoing collaboration and data exchange with regional environmental agencies on this issue.

3.7.5 Recalculations

Recalculations, respect to the previous submission occurred. Natural gas emission factor has been corrected for year 2021. Minor corrections on the figures of the gasoline engines split between 2 stroke and 4 stroke from 2014 onwards. The corrections lead to any relevant impacts on the distribution of the ratio of 2 strokes 4 strokes in the time series. Other minor recalculations affected activity data for 2020 and 2021.

3.7.6 Planned improvements

Further improvements will include a verification of activity data on ship movements with ISTAT, the National Institute of Statistics.

3.8 PIPELINE COMPRESSORS (NFR SUBSECTOR 1.A.3.E)

Pipeline compressors category (1.A.3e) includes all emissions from fuels delivered to the transportation by pipelines and storage of natural gas. Relevant pollutant emissions typical of a combustion process, such as SO_x, NO_x, CO and PM emissions, derive from this category. This category is not a key category.

Emissions from pipeline compressors are estimated on the basis of natural gas fuel consumption used for the compressors and the relevant emission factors. The amount of fuel consumption is estimated on the basis of data supplied for the whole time series by the national operators of natural gas distribution (SNAM and STOGIT) and refers to the fuel consumption for the gas storage and transportation; this consumption is part of the fuel consumption reported in the national energy balance in the consumption and losses sheet. Emission factors are those reported in the EMEP/EEA Guidebook for gas turbines (EMEP/CORINAIR, 2007). Emissions communicated by the national operators in their environmental reports are also taken into account to estimate air pollutants, especially SO_x, NO_x, CO and PM10.

Regarding QA/QC, fuel consumptions reported by the national operators for this activity are compared with the amount of natural gas internal consumption and losses reported in the energy balance as well as with energy consumption data provided by the operators to the emission trading scheme.

Starting from the length of pipelines, the average energy consumptions by kilometre are calculated and used for verification of data collected by the operators. Energy consumptions and emissions by kilometre calculated on the basis of data supplied by SNAM, which is the main national operator, are used to estimate the figures for the other operators when their annual data are not available.

In Table 3.34, nitrogen oxides, non-methane volatile organic compounds, sulphur oxides, particulate and carbon monoxide emissions (Gg) are reported.

Table 3.34 Emissions from pipeline compressors (Gg)

Emissions for NFR Subsector 1.A.3.e	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
NO _x	2.89	4.18	2.96	2.37	1.71	0.36	0.43	0.38	0.57	0.56
NMVOG	0.02	0.03	0.04	0.04	0.05	0.02	0.03	0.03	0.04	0.04
SO _x	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.005	0.005	0.005
PM10	0.02	0.03	0.05	0.03	0.03	0.02	0.02	0.02	0.02	0.03
CO	1.26	1.38	1.03	0.60	0.61	0.23	0.18	0.16	0.26	0.30

Source: ISPRA elaborations

3.9 CIVIL SECTOR: SMALL COMBUSTION AND OFF-ROAD VEHICLES (NFR SUBSECTOR 1.A.4 - 1.A.5)

3.9.1 Overview

Emissions from energy use in the civil sector cover combustion in small-scale combustion units, with thermal capacity < 50 MWth, and off road vehicles in the commercial, residential and agriculture sectors.

The emissions refer to the following categories:

- 1 A 4 a i Commercial / Institutional: Stationary
- 1 A 4 a ii Commercial / Institutional: Mobile
- 1 A 4 b i Residential: Stationary plants
- 1 A 4 b ii Residential: Household and gardening (mobile)
- 1 A 4 c i Agriculture/Forestry/Fishing: Stationary
- 1 A 4 c ii Agriculture/Forestry/Fishing: Off-road Vehicles and Other Machinery
- 1A 4 c iii Agriculture/Forestry/Fishing: National Fishing
- 1 A 5 a Other, Stationary (including military)
- 1 A 5 b Other, Mobile (Including military, land based and recreational boats)

In Table 3.35 the list of categories for small combustion and off-road vehicles identified as key categories by pollutant for 2022, 1990 and in the trend is reported.

Table 3.35 List of key categories by pollutant in the civil sector in 2022, 1990 and trend

	Key categories in 2022	Key categories in 1990	Key categories in trend
SO _x	1 A 4 b i	1 A 4 b i	1 A 4 a i
NO _x	1 A 4 b i 1 A 4 a i 1 A 4 c ii	1 A 4 c ii	1 A 4 a i 1 A 4 b i 1 A 4 c i
NMVOC	1 A 4 b i 1 A 4 a i	1 A 4 b i 1 A 4 c ii	1 A 4 b i 1 A 4 a i 1 A 4 c ii
CO	1 A 4 b i	1 A 4 b i 1 A 4 c ii	1 A 4 b i
PM ₁₀	1 A 4 b i	1 A 4 b i 1 A 4 c ii	1 A 4 b i 1 A 4 c ii
PM _{2.5}	1 A 4 b i	1 A 4 b i 1 A 4 c ii	1 A 4 b i 1 A 4 c ii
BC	1 A 4 b i	1 A 4 c ii 1 A 4 b i	1 A 4 b i 1 A 4 c ii
Cd	1 A 4 b i	1 A 4 b i 1 A 4 a i	
Hg	1 A 4 b i		
PAH	1 A 4 b i	1 A 4 b i	1 A 4 b i
DIOX	1 A 4 b i	1 A 4 a i 1 A 4 b i	1 A 4 b i 1 A 4 a i
HCB	1 A 4 b i 1 A 4 a i		1 A 4 a i 1 A 4 b i
PCB	1 A 4 b i		1 A 4 b i

3.9.2 Activity data

The Commercial / Institutional emissions arise from the energy used in the institutional, service and commercial buildings, mainly for heating. Additionally, this category includes all emissions due to wastes

used in electricity generation as well as biogas recovered in landfills and wastewater treatment plant. In the residential sector the emissions arise from the energy used in residential buildings, mainly for heating and the sector includes emissions from household and gardening machinery. The Agriculture/ Forestry/ Fishing sector includes all emissions due to the fuel, including biogas from biodigestors, used in agriculture, mainly to produce mechanical energy, the fuel use in fishing and for machinery used in the forestry sector. Emissions from military aircraft and naval vessels are reported under 1A.5.b Mobile.

Emissions from 1.A.4.a ii are reported as IE, included elsewhere, because of they refer to road transport emissions of institutional and commercial vehicles. These emissions are estimated, and reported in 1.A.3.b, with a model (COPERT 5) which consider the vehicle fleet subdivided by technology and fuel and not by user. Emissions from 1.A.5.a are also reported as IE because they refer to stationary combustion in commercial and residential of military which are included and reported in 1.A.4.a i and 1.A.4.b i; also in this case the relevant energy statistics are not available by user. The estimation procedure follows that of the basic combustion data sheet. Emissions are estimated from the energy consumption data that are reported in the national energy balance (MASE, several years (a)) and separating energy consumption between commercial/institutional, residential, agriculture and fishing, according to the information available in the Joint Questionnaire OECD/IEA/EUROSTAT prepared by the Ministry of Environment and officially sent to the international organizations. Emissions from 1.A.4.b Residential and 1.A.4.c Agriculture/Forestry/Fishing are disaggregated into those arising from stationary combustion and those from off-road vehicles and other machinery. Currently, a review process of the national energy balance is underway with Eurostat. In a first step, those responsible for the National Energy Balance implemented methodological changes for 2022 and subsequently applied these changes to 2021 to make it consistent. The review process, carried out in agreement with the relevant international institutions, regards definitions of "main activity producer" and "autoproducer" and the consequent adoption by Italy of these definitions in place of those previously used and linked to Italian legislation. This change in methodology has caused a fairly evident break in the series for some fuels and it is necessary further work to improve the time series consistency.

The time series of fuel consumption for the civil sector are reported in Table 3.36.

Table 3.36 Time series of fuel consumption for the civil sector

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
TJ										
1 A 4 a i Commercial / Institutional: Stationary plants	206,427	247,440	306,088	419,507	489,018	400,595	422,663	402,497	370,464	326,781
1 A 4 b i Residential: Stationary plants	1,002,131	1,003,620	1,036,801	1,172,245	1,222,522	1,078,829	1,019,739	997,306	1,091,916	955,359
1 A 4 b ii Residential: Household and gardening (mobile)	466	571	374	154	66	57	35	35	29	22
1 A 4 c i Agriculture/Forestry/ Fishing: Stationary	9,688	9,584	8,234	10,599	10,980	19,645	20,528	20,852	29,672	30,834
1 A 4 c ii Agriculture/Forestry/ Fishing: Off-road Vehicles and Other Machinery	96,536	101,928	94,668	95,869	84,461	81,263	83,276	84,000	82,878	78,924
1 A 4 c iii Agriculture/Forestry/ Fishing: National Fishing	8,413	9,651	8,584	10,464	7,731	6,194	6,630	6,688	6,233	5,937
1 A 5 b Other, Mobile (Including military, land based and recreational boats)	14,840	20,814	11,595	16,947	9,001	6,388	6,317	8,733	4,385	7,139

3.9.3 Methodological issues

The Tier 2 methodology is applied to the whole category. Emissions are estimated for each fuel and category at detailed level and country specific emission factors are used for the key fuel and categories drivers of total emission trend. More in detail, 1.A.4.a i, is key category in 2022 and in trend for NO_x, NMVOC and HCB emissions as well as for cadmium and Dioxin in 1990, and SO_x and dioxins in trend analysis. Most of these pollutants are due prevalently to emissions from waste incineration with energy recovery (about 99% for HCB, and more than 90% for PCB and HMs, except for Hg (78%) and Ni (83%)). Emissions from waste combustion in incinerator with energy recovery have been calculated with a Tier 3 methodology from the database of incinerator plants which includes plant specific emission factors on the basis of their technology and measurements data (ENEA-federAmbiente, 2012). The methodology used to estimate emissions from incinerators is reported in the relevant paragraph on waste incineration in the waste sector, and in particular EFs are reported in Table 7.3. Up to 2009 emission factors have been estimated on the basis of a study conducted by ENEA (De Stefanis P., 1999), based on emission data from a large sample of Italian incinerators (FEDERAMBIENTE, 1998; AMA-Comune di Roma, 1996), legal thresholds (Ministerial Decree 19 November 1997, n. 503 of the Ministry of Environment; Ministerial Decree 12 July 1990) and expert judgements. Waste management with incinerators is a commercial activity with recover of the energy auto-produced and emissions from these plants are allocated in the commercial / institutional category because of the final use of heat and electricity production. In fact, until the early 2000s, electricity and heat produced by incinerators have been prevalently used to satisfy the energy demand from connected activities: heating of buildings, domestic hot water and electricity for offices. This is still true in particular for industrial and hospital incinerators, meanwhile municipal solid waste incinerators have increased the amount of energy provided to the grid from the early 2000s until now, although only a small percentage of energy produced goes to the electricity grid (around 10%); the energy recovered by these plants is mainly used for district heating of commercial buildings or used to satisfy the internal energy demand of the plants. Since 2010, emission factors for urban waste incinerators have been updated on the basis of data provided by plants (ENEA-federAmbiente, 2012; De Stefanis P., 2012) concerning the annual stack flow, the amount of waste burned and the average concentrations of the pollutants at the stack and taking in account the abatement technologies in place. As the emission factors are considerably lower than the old ones due to the application of very efficient abatement systems it was necessary to apply a linear smoothing methodology assuming a progressive application of the abatement systems between 2005 and 2010. In a similar way, emission factors for industrial waste incinerators have been updated from 2010 onwards on the basis of the 2019 EMEP/EEA Guidebook. Similarly, to municipal waste smoothing has been applied between 2005 and 2010 supposing a linear application of the abatement systems. The other fuels driving emissions from this category are wood combustion, especially for NMVOC, and natural gas for NO_x while the trend of SO_x is driven by the decrease both of liquid fuel, as gasoil, fuel oil and kerosene, consumptions and their sulphur content which is also decreased according to European Union and national legislation. For what concerns wood combustion, the NMVOC average emission factor, as well as all the other pollutants, takes into account the different technologies used and is calculated on the basis of country specific emission factors and the ranges reported in the 2019 EMEP/EEA Guidebook; see paragraph 3.9.3.2 for details on methodology and emission factors. For natural gas and NO_x emissions a Tier 2 methodology is used and country specific emission factors, as described in the following paragraph 3.9.3.1. For the other fuels the default emission factors of EMEP/CORINAIR 2007 Guidebook have been used; it is planned to update these emission factors with those reported in the 2019 EMEP/EEA Guidebook and, according to the last review process, the update of EFs started from the 2023 report with COVNM emissions. For gasoil, biogas and gasoline different emission factors are used for stationary engines and boilers. Concerning the other pollutants, PM_{2.5} emissions from wood account for around 83% of the total 1.A.4.a i category; the other main fuel used for this category is biogas from landfills and wastewater treatment energy recovery, which account for around 9% of PM_{2.5} emissions of this category; an emission factor equal to 10 g/GJ is used. The other fuels have been estimated with EMEP/CORINAIR2007 emission factors. For NO_x, in addition to waste fuel, see methodology in the waste chapter and in particular emission factors reported in Table 7.3, and natural gas, as described in the following paragraph 3.9.3.1, the other main fuel driving emission estimates is biogas from landfills and wastewater treatment energy recovery, accounting for 44% of NO_x emissions of this category in 2022, but for which no guidance is provided in the Guidebook. An emission factor equal

to 1 kg/GJ has been used taking into account that the gas is burnt in stationary engines. NO_x emissions from waste and biogas account for more than 70% of total NO_x category emissions. HM and POP emissions from the sector are prevalently from waste incineration, estimated with country specific EFs, at technology level, and from wood combustion estimated also with country specific EFs in the range of 2019 EMEP/EEA Guidebook values.

For 1.A.4.b i, the category is key category in 2022 for SO_x, NO_x, NMVOC, CO, PM10, PM2.5, BC, Cd, Hg, PAH, Dioxin, HCB and PCB emissions. More, in 1990 NO_x, NH₃, Pb, Hg, HCB and PCB are not key categories. At trend assessment NO_x, NMVOC, CO, PM10, PM2.5, BC, PAH, dioxins, HCB and PCB are key categories. Most of these pollutants are due prevalently to emissions from wood combustion: between 98% and 100% for NH₃, CO, NMVOC, PM, BC, PAH, dioxins, HMs (except Hg with 45%, Cd with 92%, Ni with 94% and As with 81%), HCB and PCB emissions, 71% for SO_x and 45% of the total for NO_x for which a Tier 2 is applied. Methodology and emission factors are described in paragraph 3.9.3.2. For SO_x country specific and updated emission factors are used for wood, gasoil, residual oil, natural gas and LPG calculated on the basis of the maximum content of sulphur in these fuels; emissions from these fuels account for about 99.9% of SO_x category emissions. A country specific methodology has been developed and applied to estimate NO_x emissions from gas powered plants and all emissions from wood combustion. About 50% of the total emissions are due to the combustion of natural gas; methodology and country specific emission factors are described in the following paragraph 3.9.3.1. Biomass combustion accounts for around 45% of the total and methodology and country specific emission factors are also available in paragraph 3.9.3.2. All these fuels cover more than 90% of total category emissions. According to the previous review processes, an update of EFs occurs for NO_x, NMVOC and PM2.5 since 2000.

For 1.A.4.bii, 1.A.4.cii, 1.A.4.ciii and 1.A.5b emission estimates are calculated taking into account the relevant changes in emission factors along the time series due to the introduction of the relevant European Union Directives for off-road engines. Regarding mobile machinery used in agriculture, forestry and household, these sectors were not governed by any legislation until the Directive 97/68/EC (EC, 1997 [a]), which provides for a reduction in NO_x limits from 1st January 1999, and Directive 2004/26/EC (EC, 2004) which provide further reduction stages with substantial effects from 2011, with a following decreasing trend particularly in recent years. For engines with lower power as those used in forestry, household and gardening, the European Directives introduce emissions limits only starting from 2019 and 2021 so they have not had effect up to now. Moreover, for the category 1.A.4.bii, 1.A.4.cii and 1.A.4.ciii, Pb emissions from 2002 are reported as 'NA', because of the introduction of unleaded liquid fuels in the market in 2002. In particular heavy metals contents values derive from the analysis about the physical - chemical characterization of fossil fuels used in Italy (Innovhub, Fuel Experimental Station, several years). According to the review (EEA, 2019), PCB, HCB and Dioxins emissions have been estimated and included in the inventory for 1.A.4.ciii category with the emission factors of Tier1 available in the 2019 EMEP/EEA Guidebook.

3.9.3.1 NO_x emissions from gas powered plants in the civil sector

A national methodology has been developed and applied to estimate NO_x emissions from gas powered plants in the civil sector, according to the EMEP/EEA Guidebook (EMEP/EEA, 2019). On the basis of the information and data reported in available national studies for the year 2003, a distribution of heating plants in the domestic sector by technology and typology has been assessed for that year together with their specific emissions factors. Data related to heating plants, both commercial and residential, have been supplied for 2003 by a national energy research institute (CESI, 2005). In this study, for the residential sector, the sharing of single and multifamily houses plants by technology and a quantitative estimation of the relevant gas powered ones are reported, including their related NO_x emission factors. Domestic final consumption by type of plant, single or multifamily plants, has been estimated on the basis of data supplied by ENEA on their distribution (ENEA, several years). Data reported by ASSOTERMICA (ASSOTERMICA, several years) on the number of heating plants sold are used for the years after 2003 to update the information related to the technologies. A linear regression, for the period 1995-2003, has been applied, while for the period 1990-1994, the technology with the highest emission factor has been assumed to be operating. In Table 3.37 the time series of NO_x average emission factors for the relevant categories is reported.

Table 3.37 Time series of NO_x emissions factor for the civil sector

NO _x EFs	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	g/Gj									
1 A 4 a i Commercial / Institutional: Stationary	50.0	48.5	40.2	35.2	32.4	30.3	28.8	28.5	28.1	27.7
1 A 4 b i Residential: Stationary plants	50.0	48.2	38.6	32.4	31.3	30.5	29.4	29.3	29.3	29.2

3.9.3.2 Emissions from wood combustion in the civil sector

A national methodology has been developed and applied to estimate emissions from wood combustion in the civil sector, according to the TIER 2 methodology reported in the EMEP/EEA Guidebook (EMEP/EEA, 2019). In the past years, several surveys have been carried out to estimate national wood consumption in the domestic heating and the related technologies used. In the estimation process, three surveys have been taken into account: the first survey (Gerardi and Perrella, 2001) has evaluated the technologies for wood combustion used in Italy for the year 1999, the second survey (ARPA, 2007) was related to the year 2006, while the third survey (SCENARI/ISPRA, 2013) was related to the year 2012. For 2015 and 2019 information on the use of pellet, as available in the national energy balance, and on the relevant technologies, as provided by the industrial association, has been used to take in account the increase of pellet used for heating; the update has been developed taking in account also the results of the surveys on wood consumption and combustion technologies carried out by ISPRA (SCENARI/ISPRA, 2013) and by ISTAT (ISTAT, 2014). The technologies assessed by the abovementioned surveys and the distribution of fuel combustion by technologies are reported in Table 3.38.

Table 3.38 Distribution of wood combustion by technologies

Distribution of wood combustion by technologies	1999	2006	2012	2015	2019	2020-2021	2022
	%						
Fireplaces	51.3	44.7	51.2	49.0	40.3	39.7	40.8
Stoves	28.4	27.6	22.9	21.0	18.4	17.9	18.3
Advanced fireplaces	15.4	20.2	15.8	15.0	19.7	19.5	19.0
Pellet stoves	0	3.1	4.0	9.0	13.6	15.1	14.0
Advanced stoves	4.8	4.4	6.0	6.0	7.9	7.8	8.0

Average emission factors for 1999, 2006, 2012, 2015, 2019, 2020-2021 and 2022 have been estimated at national level taking into account the technology distributions; for 1990 only old technologies (fireplaces and stoves) have been considered and linear regressions have been applied to reconstruct the time series from 1990 to 2006. For the years till 2011, emission factors from 2006 have been used in absence of further available information. The distribution of combustion technologies is updated, starting from this year, on an annual basis based on the sales data of the equipment by type. For NMVOC, PAH, PM₁₀ and PM_{2.5} emission factors the results of the experimental study funded by the Ministry of Environment and conducted by the research institute 'Stazione Sperimentale dei Combustibili' now Innovhub (SSC, 2012; INNOVHUB, 2021) have been used. This study measured and compared NO_x, CO, NMVOC, SO_x, TSP, PM₁₀, PM_{2.5}, PAH and Dioxin emissions for the combustion of different wood typically used in Italy as beech, hornbeam, oak, locust and spruce-fir, in open and closed fireplaces, traditional and innovative stoves, and pellet stoves. Emissions from certificated and not certificated pellets have been also measured and compared. In general, measured emission factors results in the ranges supplied by the EMEP/EEA

Guidebook but for some pollutants and technologies results are sensibly different. In particular NMVOC emissions for all the technologies are close or lower to the minimum value of the range reported in the Guidebook, as well as PM emissions with exception of emissions from pellet stoves which are higher of the values suggested in the case of the use of not certificated pellet. For these pollutants the minimum values of the range in the Guidebook have been used when appropriate. For that concern PAH, measured emissions from open fireplaces are much lower than the minimum value of the range in the Guidebook while those from the advanced stoves are close to the superior values of the range for all the PAH compounds. In this case, for open fireplaces, experimental values have been used while for the other technologies the minimum or maximum values of the range in the Guidebook have been used as appropriate. For the other pollutants where differences with the values suggested by the Guidebook are not sensible, a more in-depth analysis will be conducted with the aim to update the emission factors used if needed. During 2020, a new experimental study funded by the Ministry of Environment and managed by ISPRA has been completed (INNOVHUB, 2021). This study regards the analysis on advanced appliances burning solid biomass (beech, fir and hornbeam, pellet A1, pellet A2). The pollutants that have been monitored are: CO, NO_x, SO₂, PM, PAH. The study also contains an interesting comparison between standard methodology and the BeReal method. In Table 3.39 emission factors used for the Italian inventory are reported.

Table 3.39 Emission factors for wood combustion

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
NO _x	50	55	59	61	61	61	66	66	66	66
CO	6000	5791	5591	5427	5395	5010	4633	4564	4564	4609
NMVOC	762	715	672	643	638	597	535	526	526	533
SO ₂	10	11	12	13	13	13	14	14	14	14
NH ₃	9	7	6	6	6	6	5	5	5	5
PM10	507	465	428	409	406	392	352	348	348	352
PM2.5	503	461	424	405	402	388	348	344	344	348
BC	40	37	35	34	34	33	31	31	31	31
PAH	0.25	0.24	0.23	0.23	0.22	0.21	0.19	0.19	0.19	0.19
Dioxin (µg/GJ)	0.48	0.47	0.45	0.44	0.43	0.40	0.37	0.36	0.36	0.37
PCB	0.00006	0.00006	0.00006	0.00006	0.00006	0.00006	0.00006	0.00006	0.00006	0.00006
HCB	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001	0.00001
As	0.001	0.001	0.001	0.001	0.0005	0.0005	0.0005	0.0005	0.0005	0.0005
Cd	0.002	0.002	0.001	0.001	0.001	0.001	0.001	0.001	0.001	0.001
Cr	0.001	0.002	0.003	0.003	0.003	0.003	0.003	0.003	0.003	0.003
Cu	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Hg	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004	0.0004
Ni	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002	0.002
Pb	0.04	0.04	0.04	0.04	0.04	0.04	0.03	0.03	0.03	0.03
Se	0.001	0.001	0.001	0.001	0.0005	0.0005	0.0005	0.0005	0.0005	0.0005
Zn	0.10	0.10	0.10	0.09	0.09	0.09	0.09	0.09	0.09	0.09
B(a)P	0.07	0.07	0.07	0.07	0.07	0.06	0.06	0.06	0.06	0.06

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
B(b)F	0.09	0.08	0.08	0.08	0.08	0.07	0.07	0.07	0.07	0.07
B(k)F	0.04	0.04	0.04	0.04	0.03	0.03	0.03	0.03	0.03	0.03
IND	0.05	0.05	0.05	0.05	0.04	0.04	0.04	0.04	0.04	0.04

Finally, ISTAT – the national statistical institute - following different requests made by various subjects interested in the data, designed and implemented a periodic campaign on "Household energy consumption" including biomass consumption. The results of this survey are the basis of the wood consumption data reported in the energy balance. In 2014 the national Institute of Statistics (ISTAT) carried out a survey, funded by the Ministry of Economic Development (MSE), on the final energy consumption of households for residential heating which include the fuel consumption of solid biomass, as wood and pellets (ISTAT, 2014). In this regard the survey resulted in an official statistic for 2012 and 2013 of wood and pellet fuel consumption at national and regional level including the information on the relevant equipment. The resulting figure for 2013 doubled the value reported in the National Energy Balance for previous years which asked for the need to update the whole time series. An ad hoc working group has been established, involving ISPRA, MSE and the energy management system national operator (GSE), to reconstruct the complete time series of wood and pellet fuel consumption which has been recalculated and officially submitted to Eurostat in June 2015. The methodology to recalculate consumption figures has taken in account the amount of wood harvested for energy purposes, the amount of wood biomass from pruning, import and export official statistics to estimate total wood consumption. A model to estimate the annual amount of wood for heating has been developed on the basis of the annual energy total biomass demand of households estimated considering the degree days time series, the number of households, the energy efficiency of equipment and fuel consumption statistics for the other fuels. As a consequence, time series for residential heating have been completely recalculated affecting the relevant pollutants and resulting in important recalculations at national total levels.

3.9.4 Time series and key categories

The time series of emissions for civil sector shows an increasing trend for several pollutants, except for SO_x and NO_x, due to a gradual shift from diesel fuel to gas, concerning SO_x, and to a replacement of classic boilers with those with low NO_x emission. Many other pollutants have a growing trend, as a consequence of the increase of wood combustion. In particular the pollutants which are more affected by the increase of wood biomass in this category according to data available in the National Energy Balance are PM, PAH, NMVOC and CO. In particular for 1.A.4.c.i the increasing trend of PAH in the last years is due to the increase of wood combustion for this category. More in detail the decrease of SO_x emissions is the combination of the switch of fuel from gasoil and fuel oil to natural gas and LPG and the reduction in the average sulphur content of liquid fuels. The SO_x emission factors for 1990 and 2022 by fuels are shown in the following Table 3.40.

Table 3.40 SO_x emission factors by fuel

FUEL – SO _x EFs(kg/Gj)	1990	2022
steam coal	0.646	0.646
coke oven coke	0.682	0.682
wood and similar	0.010	0.014
municipal waste	0.069	0.048
Biodiesel	0.047	0.047
residual oil	1.462	0.146

FUEL – SO _x EFs(kg/Gj)	1990	2022
gas oil	0.140	0.047
Kerosene	0.018	0.018
natural gas	0.0003	0.0003
Biogas	-	-
LPG	0.0022	0.0022
gas works gas	0.011	0.011
motor gasoline	0.023	0.023

Time series of emissions is reported in Table 3.41.

Table 3.41 Time series of emissions in civil sector: small combustion and off-road vehicles

		1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
SO _x (Gg)	1A4	96.5	42.7	26.4	22.8	12.1	10.3	10.1	9.8	10.1	9.2
	1A5	1.2	0.8	0.2	0.2	0.1	0.1	0.1	0.2	0.1	0.2
NO _x (Gg)	1A4	175.0	187.2	171.8	163.0	144.0	127.0	119.1	113.9	115.2	105.9
	1A5	11.2	12.0	7.2	13.5	6.1	3.3	2.7	3.5	1.9	2.8
CO (Gg)	1A4	1093.2	1088.8	1028.0	997.4	1716.1	1440.7	1307.7	1244.9	1395.3	1279.7
	1A5	65.1	79.0	45.5	54.5	17.3	16.5	17.4	24.9	12.0	20.3
PM10 (Gg)	1A4	84.6	89.8	84.6	80.3	129.1	110.2	96.1	91.6	102.7	94.1
	1A5	1.3	1.5	0.9	1.6	0.8	0.5	0.4	0.6	0.3	0.4
PM2.5 (Gg)	1A4	83.8	89.0	83.9	79.6	127.8	109.0	94.9	90.5	101.5	92.9
	1A5	1.3	1.5	0.9	1.6	0.8	0.5	0.4	0.6	0.3	0.4
BC (Gg)	1A4	14.9	16.1	14.6	12.4	13.7	11.0	9.5	9.0	9.8	8.8
	1A5	0.7	0.8	0.5	0.9	0.5	0.2	0.2	0.3	0.1	0.2
Pb (Mg)	1A4	82.0	34.3	24.6	46.3	16.5	15.1	14.1	13.6	14.7	13.7
	1A5	16.3	4.2	1.2	0.0	NA	0.1	0.2	0.2	0.1	0.1
Cd (Mg)	1A4	1.5	1.2	1.7	2.6	0.7	0.6	0.5	0.5	0.5	0.5
	1A5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Hg (Mg)	1A4	0.6	0.7	1.0	2.0	0.5	0.5	0.5	0.5	0.5	0.4
	1A5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA
PAH (Mg)	1A4	32.1	35.4	35.9	39.1	67.4	56.1	51.4	48.7	55.0	50.5
	1A5	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
HCB (Kg)	1A4	2.6	3.6	7.0	7.1	5.1	3.8	4.0	4.0	4.1	3.9
	1A5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

		1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
PCB (Kg)	1A4	14.9	18.3	25.7	36.3	22.0	19.0	19.0	18.5	20.4	18.7
	1A5	NA	NA	NA	NA	NA	NA	NA	NA	NA	NA

3.9.5 QA/QC and Uncertainty

Basic data used in the estimation process are reported by Ministry of Environment in the National Energy Balance (MASE, several years (a)) and by Terna (National Independent System Operator), concerning the waste used to generate electricity. The energy data used to estimate emissions have different levels of accuracy:

- the overall sum of residential and institutional/service/commercial energy consumption is quite reliable and their uncertainty is comparable with data reported in the BEN; the amount of fuels used is periodically reported by main suppliers;
- the energy consumption for agriculture and fisheries is reported in energy statistics; data are quite reliable as they have special taxation regimes and they are accounted for separately;
- the energy use for military and off roads is reported in official statistics, but models are applied to estimate the energy use at a more disaggregated level.

3.9.6 Recalculation

Some recalculations affected 1A4 category in this submission.

Energy recovery from waste reported in the commercial heating (1A4ai) has been updated from 2013 because of the update of data single plant leading to minor recalculations. Thanks to the NEC review process an error has been identified in NMVOC EF from clinical waste and it has been updated. More, the ratio PM2.5/PM10 has been updated for industrial waste from 2006.

In 2021, some recalculations occur because of the update of consumptions reported in National Energy Balance and of NOx emission factor on the basis of updated data published by (Assotermica, several years).

In 2020, recalculations occur because of an error on biogas has been fixed on.

Recalculations occurred for the whole time series because of the update of the energy balance, in particular the update of kerosene time series which has been added.

3.9.7 Planned improvements

The updating of average emission factors will continue in future submissions on the basis of the surveys on wood consumption and combustion technologies planned by ISTAT on fuel consumptions as well as from the results of an emission factor measurements campaign realized in Italy (ALTROCONSUMO, 2018), and the measurements campaign on advanced stoves completed by Innovhub. An in depth analysis of emission factors resulting from this experimental studies and their comparison with the values suggested by the last version of the EMEP/EEA Guidebook (EMEP/EEA, 2023) will be carried out and emission factors will be updated as needed.

3.10 FUGITIVE EMISSIONS (NFR SUBSECTOR 1.B)

3.10.1 Overview

Fugitive emissions arise during the stages of fuel production, from extraction of fossil fuels to their final use. Emissions are mainly due to leaks or other irregular releases of gases from the production and transformation of solid fuels, the production of oil and gas, the transmission and distribution of gas and from oil refining, as well as from geothermal energy production. In Table 3.42 the list of categories for fugitive emissions identified as key categories by pollutant for 2022, 1990 and in the trend is reported.

Table 3.42 List of key categories by pollutant in the civil sector in 2022, 1990 and trend

	Key categories in 2022	Key categories in 1990	Key categories in trend
SO _x	1 B 2 a iv	1 B 2 a iv	1 B 2 a iv, 1 B 2 c
NMVOOC		1 B 2 a v	1 B 2 a v
NH ₃			1 B 2 d
Hg	1 B 2 d	1 B 2 d	1 B 2 d

3.10.2 Methodological issues

In the following methodological issues including activity data and emission factors used are reported for each category and pollutant estimated in this sub sector.

Coal mining and handling (1B1a)

NMVOOC emissions from coal mining have been estimated on the basis of activity data published on the national energy balance (MASE, several years [a]) which report the amount of coal production and emission factors provided by the EMEP/EEA Guidebook (EMEP/EEA, 2016). PM emissions from storage of solid fuels have been estimated and included in this category. Activity data is the annual consumption of solid fuels published on the national energy balance (MASE, several years [a]) and emission factor are from the US EPA Guidebook.

Solid fuel transformation (1B1b)

NMVOOC emissions from coke production have been estimated on the basis of activity data published in the national energy balance (MASE, several years [a]) and country specific emission factors calculated taking in account the information provided by the relevant operators in the framework of the EPRTTR registry and the ETS. NO_x, SO_x and NH₃ emissions from coke production are estimated on the basis of data communicated by the national plants in the framework of the EPRTTR and are reported under 1.A.1 c category. NH₃ emissions have been estimated on the basis of data communicated by the operators for the EPRTTR registry from 2002. According to the review (EEA, 2019; EEA, 2020), PAH emissions from coke production has been also estimated with emission factor from the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019) and allocated in 1.B.1b.

Oil exploration and production (1B2a i)

NMVOOC emissions have been calculated according with activity data published on national energy balance (MASE, several years [a]), data by oil industry association (UNEM, several years), data and emission factors provided by the relevant operators.

Oil transport and storage and refining (1B2a iv)

Fugitive emissions from oil refining are estimated starting from the total crude oil losses as reported in the national energy balance (MASE, several years [a]) and occur prevalently from processes in refineries. This category is key for SO_x in 2022, in the base year and for the trend. Emissions in refineries have been estimated on the basis of activity data published in the national energy balance (MASE, several years [a]) or supplied by oil industry association (UNEM, several years) and operators especially in the framework

of the European Emissions Trading Scheme (EU-ETS). Fugitive emissions in refineries are mainly due to catalytic cracking production processes, sulphur recovery plants, flaring and emissions by other production processes including transport of crude oil and oil products. These emissions are then distributed among the different processes on the basis of average emission factors agreed and verified with the association of industrial operators, Unione Energia per la Mobilità, and yearly updated, from 2000, on the basis of data supplied by the plants in the framework of the European Emissions Trading Scheme, Large Combustion Plant Directive and EPRTR. SO_x, NO_x and PM emissions communicated by the plants in the framework of Large Combustion Plants directive are assumed to refer to combustion and are reported under 1.A.1b while the difference with the totals, communicated to the EPRTR, are considered as fugitive emissions and reported in 1.B.2a iv. NMVOC are communicated by the operators for the EPRTR registry as a total and the amount to be reported as fugitive is calculated subtracting by the total emission estimates for combustion activities and reduced for the implementation of losses control technology especially for transportation and storage of liquid fuels. ETS data are used to integrate and check emission data provided. Moreover, fugitive emissions are also checked with the average emission factors provided by the relevant industrial association for each relevant process, as fluid catalytic cracking, sulphur recovery plant, and storage and handling of petroleum products. NH₃ emissions from refineries have been estimated on the basis of data communicated by the operators for the EPRTR registry and distributed between combustion and fugitive emissions according to the emission factors available in the 2016 EMEP/EEA Guidebook. Emissions from refineries of HM and POPs are all reported in 1.A.1b on the basis of data submitted in the PRTR framework at plant level; it is not possible at the moment distinguish combustion by fugitive emissions of HM and POPs.

Distribution of oil products (1B2a v)

This category is key for NMVOC in 1990 and for the trend. The category includes fugitive emissions from oil transport which have been calculated according with the amount of transported oil (MIT, several years) and emission factors published on the IPCC guidelines (IPCC, 2006). Most of the crude oil is imported in Italy by shipment and delivered at the refineries by pipelines as offshore national production of crude oil. The category includes also NMVOC fugitive emissions for gasoline distribution, storage and at service stations. Emission factors are estimated starting from the emissions communicated in the nineties by the operators and applying the implementation of the abatement technologies as regulated by the relevant European Union legislation. Emissions from distribution of gasoline have been reduced as a result of the application of the DM 16th May 1996 (Ministerial Decree 16 May 1996), concerning the adoption of devices for the recovery of vapours and of the applications of measures on deposits of gasoline provided by the DM 21st January 2000 (Ministerial Decree 21 January 2000).

Flaring in refineries (1B2c)

This category is key for SO_x for the trend. For what concern emissions from flaring in refineries, the emission factors for SO_x, NO_x, NMVOC and CO have been provided by the relevant industrial association and are assumed constant since 1990 with the exception of SO_x that are yearly estimated on the basis of the amount of sulphur not recovered by the operators and flared. Activity data, in terms of gas flared, is from 2005 derived by the ETS data at plant level.

Fugitive emissions from geothermal production (1B2d)

According to the review process NH₃, Hg and other heavy metals from geothermal production has been estimated and included in the emission inventory in the 2018 submission with a Tier 2 methodology. Hg from this category is key category for 2022, for the base year and the trend while NH₃ is key only for the trend. Emissions are monitored by the Regional relevant environmental agency, ARPAT, where all the geothermal fields are located. Activity data, geothermal energy production, are published in the national energy balance (MASE, several years [a]) while emission data resulting by the monitoring are issued by ARPAT and reported from 2000 on yearly basis (ARPAT, several years). For earlier years emission factors of 2000 have been used.

3.10.2.1 Fugitive emissions from natural gas distribution (1.B.2b)

NMVOC fugitive emissions from the transport, storage and distributions (including housing) of natural gas (both in pipelines and in the distribution network) are calculated every year on the basis of fugitive

natural gas emissions and the content of NMVOC in the gas distributed; NMVOC emissions due to transport and distribution are around 99% of the total. Emissions are calculated starting from methane emissions estimates, considering the annual average percentage of NMVOC in the natural gas distributed in Italy as in Table 3.43. The methodology and references are reported in detail in the NIR (ISPRA, 2024[a]). CH₄, CO₂ and NMVOC emissions have been estimated on the basis of activity data published by industry, the national authority, and information collected annually by the Italian gas operators. Emission estimates take into account the information on: the amount of natural gas distributed supplied by the main national company (SNAM); length of pipelines, distinct by low, medium and high pressure and by type, cast iron, grey cast iron, steel or polyethylene pipelines as supplied by the national authority for the gas distribution (AEEG); natural gas losses reported in the national energy balance; methane emissions reported by operators, in their environmental reports (EDISON, SNAM, ENEL, Italgas). NMVOC and CO₂ emissions have been calculated considering CO₂ content in the leaked natural gas. Regarding exploration and production, an average emission factor, equal to 0.04 g/m³ gas produced, has been estimated on the basis of emission data communicated by the relevant companies for some years and applied to the whole time series.

The average natural gas chemical composition has been calculated from the composition of natural gas produced and imported. Main parameters of mixed natural gas, as calorific value, molecular weight, and density have been calculated as well. Data on chemical composition and calorific value are supplied by the main national gas providers for domestic natural gas and for each country of origin.

The following table shows average data for national pipelines natural gas.

Table 3.43 Average composition for pipelines natural gas and main parameters

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
HCV (kcal/m ³)	9,156	9,193	9,215	9,261	9,325	9,303	9,336	9,340	9,377	9,428
NCV (kcal/m ³)	8,255	8,290	8,320	8,354	8,412	8,391	8,428	8,432	8,467	8,515
Molecular weight	17.03	17.19	17.37	17.45	17.46	17.34	17.29	17.33	17.48	17.56
Density (kg/Sm ³)	0.72	0.73	0.74	0.74	0.74	0.73	0.73	0.73	0.74	0.74
CH ₄ (molar %)	94	93	92	92	92	93	93	93	92	91
NMVOC (molar %)	3	4	5	5	6	5	5	6	6	7
CO ₂ (molar %)	0.22	0.20	0.18	0.49	0.75	0.70	0.64	0.88	0.91	0.83
Other no carbon gas (molar %)	2.03	2.34	2.76	2.24	1.48	1.32	0.95	0.70	0.91	0.94
CH ₄ (weight %)	89	87	85	85	85	86	86	86	84	84
NMVOC (weight %)	7	9	10	11	11	10	11	11	12	13
CO ₂ (weight %)	0.57	0.51	0.47	1.23	1.89	1.78	1.63	2.25	2.29	2.09
Other no carbon gas (weight %)	3.27	3.74	4.37	3.51	2.30	2.08	1.50	1.09	1.41	1.43

More in details, emissions are estimated separately for the different phases: transmission in primary pipelines and distribution in low, medium, and high-pressure network, losses in pumping stations and in reducing pressure stations (including venting and other accidental losses) with their relevant emission factors, considering also information regarding the length of the pipelines and their type.

Emissions from low pressure distribution also include the distribution of gas at industrial plants and in residential and commercial sector; data on gas distribution are only available at an aggregate level thus not allowing a separate reporting. In addition, emissions from the use of natural gas in housing are

estimated and included. Emissions calculated are compared and balanced with emissions reported by the main distribution operators. Finally, the emission estimates for the different phases are summed and reported in the most appropriate category (transmission/distribution).

Table 3.44 provides the trend of natural gas distribution network length for each pipeline material and the average CH₄ emission factor.

Table 3.44 Length of low and medium pressure distribution network (km) and network emission factors for CH₄ and NMVOC

Material	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Steel and cast iron (km)	102,061	131,271	141,848	154,886	198,706	203,116	206,855	208,044	208,935	209,740
Grey cast iron (km)	24,164	22,784	21,314	15,080	4,658	2,398	2,061	2,061	2,060	2,060
Polyethylene (km)	775	8,150	12,550	31,530	49,663	56,943	59,593	59,854	61,325	61,441
Total (km)	127,000	162,205	175,712	201,496	253,027	262,457	268,509	269,959	272,320	273,241
CH ₄ EF (kg/km)	1,958	1,417	1,228	1,000	703	540	371	350	333	278
NMVOC EF (kg/km)	162	140	144	127	94	65	46	44	47	43

3.10.3 Time series and key categories

The trend of fugitive emissions from solid fuels is related to the extraction of coal and lignite that in Italy is quite low. The decrease of NMVOC fugitive emissions from oil and natural gas is due to the reduction of losses for gas transportation and distribution, because of the gradual replacement of old grey cast iron pipelines with steel and polyethylene pipelines for low and medium pressure network as reported in the previous paragraph.

3.10.4 QA/QC and Uncertainty

Different data sources are used for fugitive emissions estimates: official statistics by Environmental Ministry (MASE, several years [a], [b]), by Transport of Infrastructure Ministry (MIT, several years); national authorities (AEEG, several years; ISTAT, several years [a]), gas operators (ENI, several years [a]; EDISON, several years; SNAM, several years), and industrial association for oil and gas (UP, several years).

CH₄ and NMVOC emissions from transmission and distribution of natural gas are verified considering emission factors reported in literature and detailed information supplied by the main operators (ENI, several years [b]; Riva, 1997).

3.10.5 Recalculation

Minor changes occur for geothermal NH₃, Pb, Cd and Hg emissions in 2016, 2018 and 2020 due to update of data provided by the plant operators.

For COVNM emissions recalculation occur in 2021 because of the update of activity data.

3.10.6 Planned improvements

No further improvements are planned for this category.

4 IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2)

4.1 OVERVIEW OF THE SECTOR

Emission estimates in this category include emissions from all industrial processes and also by-products or fugitive emissions, which originate from these processes. Where emissions are released simultaneously from the production process and from combustion, as in the cement industry, they are estimated separately and included in the appropriate categories, in sector 2 and in sector 1 category 1.A.2. This sector makes important contributions to the emissions of heavy metals, PAH, dioxins and PCB. Regarding emissions of the main pollutants, in 2022, industrial processes (without the use of solvent) account for 7.7% of SO₂ emissions, 0.8% of NO_x, 0.2% of NH₃, 5.2% of NMVOC and 3.3% of CO. About particulate matter, in 2022 this sector accounts for 20.2% of PM₁₀ emissions and 7.3% of PM_{2.5}. Industrial processes make a significant contribution to the total Italian emissions of heavy metals, despite significant reductions since 1990; particularly this sector accounts for 34.7% of Pb emissions, 32.0% of Cd and 49.2% of Hg. Regarding POPs emissions, 11.8% of PAH total emissions is emitted from industrial processes as well as 26.6% of dioxins and 71.3% of PCB.

In 2022, iron and steel sector (2C1) is a key category at level assessment for PM_{2.5}, Pb, Cd, Hg, PAH, PCDD/F and PCB; emissions from other chemical industry (2B10a) is a key category source for SO₂ emissions; emissions from 2A5a Quarrying and mining and 2A5b Construction and demolition are key categories for PM₁₀. Emissions from 2G is a key category source for Cd, too. Food and beverage industry (2H2) is a key category for NMVOC emissions. In 1990 emissions from cement production (2A1) is a key category source for PM₁₀ and PM_{2.5}; other chemical industry (2B10a) is a key category for Hg and Cd and iron and steel production (2C1) is a key category for PM₁₀, PM_{2.5}, Cd, Hg, PAH, PCDD/F and PCB; quarrying and mining (2A5a) and construction and demolition (2A5b) are a key sources for PM₁₀; emissions from 2G is a key source for Cd too. At trend assessment, iron and steel sector is key category for Pb, Cd, Hg, PAH PCDD/F and PCB while other chemical industry (2B10a) is a key category for Hg, 2H2 is a key category for NMVOC and 2G is a key category for Cd emissions. Emissions from 2A5a quarrying and mining and from 2A5b Construction and demolition are a key category for PM₁₀.

As requested by the TERT in the last review processes, information has been included in this chapter about:

- the estimates for category 2A5b Construction and Demolition;
- the estimation process for PM_{2.5} from 2A1 Cement production;
- the estimates for category 2A5a Quarrying and mining;
- explanation of estimation for PAH from category 2C1 Iron and Steel, methods and data used;
- method, AD, EF for PM₁₀, PM_{2.5}, BC, PCDD/F and HCB from 2C3 Aluminium production.

4.2 METHODOLOGICAL ISSUES

Methodologies used for estimating emissions from this sector are based on and comply with the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007) and EMEP/EEA guidebook (EMEP/EEA, 2019), the IPCC Guidelines (IPCC, 1997; IPCC, 2006) and the Good Practice Guidance (IPCC, 2000). Included also in this sector are by-products or fugitive emissions, which originate from industrial processes.

There are different sources relevant to estimate emissions from this sector; activity data are provided by national statistics and industrial associations, but a lot of information is supplied directly from industry. In fact, as for the energy sector, references derive from data collected in the framework of the national PRTR reporting obligation, the Large Combustion Plant directives and the European Emissions Trading Scheme. Other small plants communicate their emissions which are also considered individually. These processes have improved the efficiency in collecting data and the exchange of information. Whenever data cannot be straight used for the inventory compilation, they are taken into account as verification practice. Environmental Reports published by industrial associations are also considered in the verification process.

4.2.1 Mineral products (2A)

In this sector emissions from the following processes are estimated and reported: cement production, lime production, glass production, quarrying&mining and construction and demolition.

Cement production (2A1), is considerable for PM_{2.5} emissions and accounts for 1.4% of the total national emissions in 2022. SO₂ emissions from cement production results from fuel combustion during the manufacture of cement so since the last submission they have been included under the energy sector 1A2f.

During the last 15 years, in Italy, changes in cement production sector have occurred, leading to a more stable structure confirming the leadership for the production in Europe. The oldest plants closed, wet processes were abandoned in favour of dry processes so as to improve the implementation of more modern and efficient technologies. Since 2011 Italy has become the second cement producer country in the EU 28 but the reduction in clinker production seems to have stopped, since 2016 clinker production at national level has kept almost the same. In 2022, 17 companies (50 plants of which: 29 full cycle and 21 grinding plants) operate in this sector: multinational companies and small and medium size enterprises (operating at national or only at local level) are present in the country. As for the localization of the operating plants: 44% is in northern Italy, 18% is in the central regions of the country and 38% is in the southern regions and in the islands (Federbeton/AITEC, 2023). In Italy different types of cement are produced; as for 2022 Federbeton/AITEC, the national cement association, has characterised the national production as follows: 72% is CEM II (Portland composite cement); 12% is CEM I (Portland ordinary cement); 12% is CEM IV (pozzolanic cement) and 4% is CEM III (blast furnace cement). Clinker production has been decreasing since 2007, although from 2016 to 2019 the production values have kept very close to the amount manufactured in 2016, in 2022 clinker production shows +3.3% compared to 2021; clinker demand in cement production was about 83% in 2022 (production of clinker out of production of cement). To estimate emissions from cement production, activity data on clinker/cement production are used as provided by ISTAT (ISTAT, several years up to 2008), MSE (MSE, several years since 2009 up to 2018) and facility reports in the framework of the Emissions Trading Scheme national legislation.

In this category only PM₁₀ and PM_{2.5} emissions are reported separately from combustion while all the other pollutant emissions are included in the energy sector in 1.A.2f category.

Emission factor for PM₁₀ emissions is equal to 234 g/Mg of clinker for the whole time series and is calculated on the basis of plants emission data in the nineties. Emission factor for PM_{2.5} is equal to 130 g/Mg of clinker for the whole timeseries.

Regarding SO₂ emissions, according to the EMEP/EEA Guidebook, no sulphur oxides process emissions have been reported under 2A1 because it was clarified that SO₂ emissions originate from fuel combustion and, as far as energy aspects only are concerned, they have been allocated under 1A2f.

The remaining categories of mineral products (*lime production (2A2)*) industry represent less than 1% for each pollutant.

As regards 2A3 category *Glass production*, HM, PM and BC emissions are reported under 1A2f and emission factors have been provided by the research institute of the sectoral industrial association (Stazione Sperimentale del Vetro) distinguished by the different types of glass production. On the basis of the 2017 review process (EEA, 2017 [a]), the previous notation key has been replaced by the IE notation key.

About the 2A5 category, following the suggestions of the NECD review, more information has been added but different activities have to be dealt with separately.

As regards 2A5a *Quarrying and mining of minerals other than coal*, there is no evidence of active mines of the main minerals as those indicated in the Guidebook (bauxite, copper, manganese and zinc). All these mines in Italy closed before 1990 for market reasons. At the same time there is no available data to apply a Tier 1 on other mineral mines. The value-added for this activity was adopted as a proxy to develop the

other mineral ore production time series and emissions factors included in 2019 EMEP/EEA Guidebook for PM10 and PM2.5 were applied to estimate PM10 and PM2.5 emissions along the whole timeseries.

As for the category 2A5b *Construction and Demolition*, the National Institute of Statistics provides information about residential and non-residential buildings, for both categories information about volumes (for the year from 1990-2000) or surfaces (usable living areas) could be retrieved to work out a timeseries of activity data and to develop estimates for TSP, PM10 and PM2.5. Emission factors for TSP, PM10 and PM2.5 are those included in the 2019 EMEP/EEA Guidebook which provides specific values for residential and non-residential buildings. As suggested by the last review the timeseries now includes also the estimation of TSP, PM10 and PM2.5 emissions from construction and demolition of roads. The Italian Ministry of transportations provides information about changes in the length of roads by road types, so that emission factors included in 2019 EMEP/EEA Guidebook for TSP, PM10 and PM2.5 can be applied to changes in roads surfaces.

For the category 2A5c *Storage, Handling and Transport of Mineral Products*, PM2.5 emissions have been estimated and reported in the sectoral categories 2A1 Cement Production and 2A2 Lime Production. The emissions from storage, handling and transport for other minerals than the aforementioned ones might not be included in the inventory because this potential under-estimate is likely to be below the threshold of significance.

4.2.2 Chemical industry (2B)

Emissions of this sector derive from organic and inorganic chemicals processes and are usually not significant except for SO_x emissions from the production of sulphuric acid and Hg emissions from chlorine production. Emission factors derive from data collected in the framework of the national EPER/E-PRTR register as well as from EMEP/EEA and EPA Guidebook.

As already mentioned, other chemical industry (2B10a) was key category for Cd and Hg emissions in 1990 and for SO_x emissions in 2022. Hg emissions are released from chlorine production facility with mercury cells process (EUROCHLOR, 1998). Total chlorine production in Italy amounted, in 1990, to 1,042,921 tonnes and reduced in 2021 to 236,735 tonnes. Activity production data are supplied by the National Institute of Statistics (ISTAT) and published in the official national statistics and since 2002 data have also been collected at industrial facility level in the national EPER/E-PRTR register. To estimate emissions from 1990 to 2001, the average emission factor supplied by EUROCHLOR for western Europe chlor-alkali production plants (EUROCHLOR, 2001) has been used, while since 2002 emission data have been supplied directly by the production facilities in the framework of the national EPER/E-PRTR. The average emission factor decreased from 1.11 g Hg/t in 2002 to zero in 2018. The reduction observed in emissions and zero emissions since 2018 are a consequence of both the conversion of production plants from the mercury cells process to the membrane technology and to the suspension of production at the existing facilities. In 2007 seven facilities carried out the chlor-alkali production: one facility had the membrane process in place, one facility was replacing mercury cells with membrane process while in the other five facilities the production was still based on the mercury cell process (Legambiente, 2007). In 2015 five facilities carried out chlor-alkali production: in four of them the membrane process was in place while one facility still operated the mercury cell process. In 2018 the four chlor-alkali facilities have the membrane process in place while the one still with mercury cells was obliged to stop the production with this technology and it is still in operation although the manufacturing process has been relying on the purchase of the intermediate products since then.

SO_x emissions reported in other chemical industry (2B10a) include emissions from sulphuric acid production and account for 5.9% of total SO_x emissions in 2022. Activity production data are supplied by the National Institute of Statistics (ISTAT) and published in the official national statistics and since 2004 data have also been collected at facility level in the national EPER/E-PRTR register. Emission factors from 1990 to 1994 and from 2002 are derived from emission data supplied directly by the production facilities in the framework of the CORINAIR inventory project and of the national EPER/E-PRTR, respectively.

On the basis of the 2017 review process NO_x, SO_x, CO, PM and BC emissions from 2B7 Soda ash production have been estimated. In Italy there is only one plant producing soda ash and it is in the framework of the EPRTR reporting. In particular, as regards PM emissions, the operator has never reported

PM10 emissions which implies that emissions are below the reporting threshold (50 t/year). As reported in the Guidebook measurements made in some plants indicate that more than 75% of the dust emitted is made of particle size > 10 µm and that the contribution of PM10 is relatively low. Moreover, the operator in its annual environmental report estimates TSP emissions (around 200 t/y) reporting explicitly that no PM10 emissions occur. The PM10 estimated using the EMEP/EEA EFs, is about 20 Mg which is anyway below the E-PRTR reporting thresholds.

4.2.3 Metal production (2C)

The main activities in this sector are those regarding the iron and steel production. The main processes involved in iron and steel production are those related to sinter and blast furnace plants, to basic oxygen and electric furnaces and to rolling mills.

The sintering process is a pre-treatment step in the production of iron where fine particles of metal ores are agglomerated. Agglomeration of the fine particles is necessary to increase the passageway for the gases during the blast furnace process and to improve physical features of the blast furnace burden. Coke and a mixture of sinter, lump ore and fluxes are introduced into the blast furnace. In the furnace the iron ore is increasingly reduced and liquid iron and slag are collected at the bottom of the furnace, from where they are tapped. The combustion of coke provides both the carbon monoxide (CO) needed for the reduction of iron oxide into iron and the additional heat needed to melt the iron and impurities. The resulting material, pig iron (and also scrap), is transformed into steel in subsequent furnaces which may be a basic oxygen furnace (BOF) or electric arc furnace (EAF). Oxygen steelmaking allows the oxidation of undesirable impurities contained in the metallic feedstock by blowing pure oxygen. The main elements thus converted into oxides are carbon, silicon, manganese, phosphorus and sulphur.

In an electric arc furnace steel is produced from polluted scrap. The scrap is mainly produced by cars shredding and does not have a constant quality, even if, thanks to the selection procedures, the scrap quality becomes better year by year. The iron and steel cycle is closed by rolling mills with production of long products, flat products and pipes.

In 1990 there were four integrated iron and steel plants in Italy. Since 2020, there is only one of the above-mentioned plants remaining; oxygen steel production represents about 15.8% of the total production and the arc furnace steel the remaining 84.2% (FEDERACCIAI, several years). Currently, long products represent about 49.0% of steel production in Italy, flat products about 38.9%, and pipe the remaining 12.0%. Most of the flat production derives from the only one integrated iron and steel plant while, in steel plants equipped with electric ovens almost all located in the northern regions, long products are predominantly produced (e.g carbon steel, stainless steels) and seamless pipes (only one plant) (FEDERACCIAI, several years).

Basic information for Iron and steel production derives from different sources in the period 1990-2021. Activity data are supplied by official statistics published in the national statistics yearbook (ISTAT, several years) and by the sectoral industrial association (FEDERACCIAI, several years). For the integrated plants, emission and production data have been communicated by the two largest plants for the years 1990-1995 in the framework of the CORINAIR emission inventory, distinguished by sinter, blast furnace and BOF, and by combustion and process emissions. From 2000 production data have been supplied by all the plants in the framework of the ETS scheme, for the years 2000-2004 disaggregated for sinter, blast furnace and BOF plants, from 2005 specifying carbonates and fuels consumption. For 2002-2015 data have also been supplied by all the four integrated iron and steel plants in the framework of the EPER/E-PRTR registry but not distinguished between combustion and process. National experts have also been involved in the process of elaboration of the "monitoring and control plan" for the largest integrated plant in Italy in the framework of the IPPC permit. Qualitative information and documentation available on the plants allowed reconstructing their history including closures or modifications of part of the plants; additional qualitative information regarding the plants, collected and checked for other environmental issues or directly asked to the plant, permitted to individuate the main driving of the emission trends for pig iron and steel productions. Emissions from lime production in steel making industries are reported in 1A2f Manufacturing Industries and Construction category. In 2022, iron and steel sector (2C1) is key category for PM2.5, Pb, Cd, Hg, PAH, PCDD/F and PCB. In Table 4.1 relevant emission factors are reported.

Table 4.1 Emission factors for iron and steel for the year 2022

		PM10 [g/Mg]	PM2.5 [g/Mg]	Cd [mg/Mg]	Hg [mg/Mg]	Pb [mg/Mg]	PCB [mg/Mg]	PAH [mg/Mg]	PCDD/F [µg T- eq/Mg]
Blast furnace charging	Areal	60	37.5						
	Point	2.8	1.8						
Pig iron tapping	Areal	41.4	25.9	0.3	0.3	15		950	
	Point	2.0	1.2	0.3	0.3				
Basic oxygen furnace	Areal	62	54.3	25	3	850	3.6		
	Point	10.7	9.4	2.3	1.7	49	3.6		
Electric arc furnace		124	108.5	50	150	3450	3.6	1.9	4.45
Rolling mills	Areal	59	45.9					125	
	Point	28.2	21.9					125	
Sinter plant (except combustion)	Areal	16	12.8						
	Point	3.5	2.8						

PM10 emission factors for integrated plants derive from personal communication of the largest Italian producer of pig iron and steel (ILVA, 1997) while PM10 emission factor for electric arc furnace derives from a sectoral study (APAT, 2003). The Emission factors manual PARCOM-ATMOS (TNO, 1992), the EMEP/Corinair Guidebook (EMEP/CORINAIR, 2006) and the IPPC BRef Report (IPPC, 2001) provide emission factors for heavy metals while a sectoral study (APAT, 2003) provides Cd emission factors for electric arc furnace. Recently, as reported above, work has begun to revise the EFs from the largest integrated plant (point sources) thanks to the analysis of the monitoring and control plans linked to the data collected in the framework of the IPPC permits. Emission factors derived from this survey on the IPPC permits have been applied since 2016 and relate both to combustion and process categories. Further information about combustion emissions is given in the combustion chapter. As regards the category 2C1, at the moment, EFs for PM emissions from blast furnace charging, pig iron tapping, basic oxygen furnace and sinter plant have been updated in the 2022 submission as for BOF emissions where Cd, Pb and Hg EFs have been updated too.

Regarding PAH emissions, for blast furnaces, results from measurements tell us that emission factor used for pig iron tapping covers also blast furnace charging while emissions from BOF are negligible. To complete the category analysis, for 1A1c category and, in particular, for coke production according to the review (EEA, 2019) PAH emission factor has been disaggregated into those deriving from the combustion process and the fugitive ones and estimated with the emission factors in the Guidebook (EMEP/EEA, 2019). Regarding POPs emissions, emission factors usually originate from EMEP/CORINAIR (EMEP/CORINAIR, 2007, EMEP/CORINAIR, 2006) except those relating to PAH and PCDD/PCDF from electric arc furnace that derive from direct measurements in some Italian production plants (ENEA-AIB-MATT, 2002). Dioxin emissions for sinter plant, and other sources within steelworks manufacturing oxygen steel occur during the combustion process and they are measured at the stack; emissions are therefore reported in the energy sector in 1.A.2a category. In 2022 the average emission factor is equal to 0.22 micrograms TEQ per Mg of sinter produced. EF is calculated yearly on the basis of measurements done in the two existing sinter plant in Italy. As regards HCB emissions, Italy reports HCB emissions from sintering production calculated with the 2006 Guidebook ("Sources of HCB emissions.pdf" does not distinguish between combustion and process. The 2016 Guidebook provides reference to the 2006 version) EF=0.032mg/Mg in 1A2a because in this case HCB emissions are clearly linked to the combustion activities. As for other iron and steel activities, a series of technical meetings with the most important

Italian manufacturers was held in the framework of the national PRTR in order to clarify methodologies for estimating POPs emissions. In the last years, a strict cooperation with some local environmental agencies allowed the acquisition of new data, the assessment of these data is still ongoing and improvements in emission estimates are expected for the next years. Thanks to the last review process in the framework of the NEC Directive (EEA, 2019) fugitive PAH emissions from coke oven (door leakage and extinction) have been estimated on the basis of 2019 EMEP/EEA Guidebook emission factors. As a consequence of the review process these emissions have been reported in 1B1b adopting the right allocation. Emission factors used in 1990 estimates generally derive from Guidebook EMEP/CORINAIR. The remaining categories of metal production industry represent about 0% for each pollutant because of the shutdown of several plants, in particular those linked to the non-ferrous production.

No plants for aluminium production by electrolysis have been working in Italy since 2012 and pollutants time series are reported, obviously, from 1990 to 2012, consequently no primary aluminium production has been occurring since 2012. Secondary aluminium production still occurs in Italy, activity data are provided annually by the statistical office of Assomet, the trade association dealing with non-ferrous metals. Emissions from secondary aluminium production are reported under the energy sector (1A2b). For that, from 2012 notation key "IE" has been reported in this category the review process. HCB emissions from secondary aluminium production are not reported and are expected to be null because these emissions derive from the degassing of aluminium when hexachloroethane is used, but this compound is banned in Italy from '90s. Further information about emissions from secondary aluminium production are included in the paragraph devoted to category 1A2b included in Chapter 3, paragraph 3.3.1.2 and 3.3.2.2.

As for the production of lead, zinc and copper (2C5, 2C6 and 2C7a categories), at the moment SO_x, HM and PM emissions are reported in the energy sector because up to now there was no information to distinguish between energy and process emissions and, above all, these processes are considered combustion processes with contact, consequently, emissions are dependent on the combustion process. In the last year, thanks to the ETS data, it has been possible to separate CO₂ emissions in these two components and Italy is investigating the possibility of extension to other pollutants for the next submissions. In particular, in Italy no production of primary copper has ever occurred while, as regards lead and zinc, there is a sole integrated plant for the primary productions, and this makes it difficult to ensure a good breakdown. Consequently, the issue related to the allocation of emissions is not only about combustion and process but also about the different productions of different metals in the same factory. To resolve this issue, an in-depth investigation has been started with the aim to better specify the technology used on the basis of E-PRTR and IPPC permits. Thanks to the analysis of these monitoring data, it was possible to recalculate the emissions from the production of zinc and lead starting from the measurements at the chimneys since 2014, the first year of useful data (in 2013 the monitoring system covered only 3 months). The first result of this investigation has been the update of EFs since 2014 (ISPRA, 2023). For the next submission the reconstruction of the historical series in the previous years is foreseen and, finally, the distinction between process and combustion emissions. More information on the upgrade done can be found in the paragraph on the combustion of non-ferrous metals (1A2b). Anyway, for Pb, Cd and PCB the notation key IE has been added in the NFR because of the relevant emissions are reported in the energy sector.

Moreover, in response to the review process Italy explained that the Hg emission factor for copper production in the 2019 EMEP/EEA Guidebook is not applicable because it refers to primary copper production while in Italy copper production between 1990 and 1998 was derived only from secondary technologies.

4.2.4 Other production (2G – 2H – 2I – 2L)

2G sector includes NMVOC emissions due to the use of lubricants as well as all potential emissions from the use of tobacco and explosives/fireworks. In 2H sector, non-energy emissions from pulp and paper as well as food and drink production, especially wine and bread, are reported. TSP emissions from wood processing are included and reported in 2I, Lead emissions from batteries manufacturing can be found in 2L sector.

Emissions from these categories are usually negligible except for NMVOC emissions from food and drink (2H2) accounting for 3.3% of the national total in 2022 and Cd from fireworks (2G) accounting for 8.5% of the national total, both are key categories in 2022. Emissions from food and drink (2H2) refer to the processes in the production of bread, wine, beer and spirits. Activity data are derived from official statistics supplied by the National Institute of Statistics (ISTAT) and relevant industrial associations. Time series of bread production is reconstructed for the '90 years on the basis of family surveys from the national Institute of statistics (ISTAT) while from 1998 data are those reported in the PRODCOM statistics officially communicated by ISTAT to EUROSTAT. PRODCOM data collection has improved along the years producing more reliable figures. In the '00 years, bread production has changed from fresh artisanal production to a more industrial oriented production, without any impact on the total. For wine, beer and spirits the statistical information on activity data is much more reliable and their trends are driven by the seasonal variation (for wine) or market demand (for beer) while for spirits it is mostly driven by a change in the personal habits and corresponding consumptions. Emission factors are those reported in the EMEP/CORINAIR Guidebook and, in lack of national information, they are assumed constant for the whole time series (CORINAIR, 1994; EMEP/CORINAIR, 2006).

Regarding 2G category (other product use) all potential emissions have been estimated both for the use of tobacco and fireworks; NMVOC, SO_x, NO_x, CO, NH₃, Cd, Pb, PM₁₀, PM_{2.5}, PCDD, Benzo(a)pyrene and PAH are estimated. For activity data, as regards fireworks, Eurostat data on import, export and production of fireworks have been used, while data on consumption of tobacco were collected from the Ministry of Health, observatory of tobacco smoking. Emission factors are those reported in the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019). NMVOC emissions from use of lubricants are also estimated. Activity data are from the energy statistics (MASE, several years [b]) while emission factor is from the EMEP/EEA Guidebook (EMEP/EEA, 2019).

Pulp and paper industry (2H1) referred to the acid sulphite and neutral sulphite semi-chemical processes up to 2007 and only to the neutral sulphite semi-chemical process for 2008 and 2009, while the kraft process was not present in Italy. Emissions of NO_x, NMVOC, SO_x and PM were estimated for those years on the basis of activity data provided by the two Italian production plants. In 2008 the bleached sulphite pulp production stopped and in 2009 the neutral sulphite semi-chemical pulp process plant also closed. So, for the IPPU inventory purposes, there was no production of pulp and paper after 2009 and consequently no emissions have been estimated. Acid sulphite process emissions are calculated for SO_x, NMVOC and NO_x on the basis of EFs available in the Best Available Techniques Reference Documents report (BRef report), for PM₁₀ on the basis of EF in the USEPA Guidebook (54% PST) while for PM_{2.5} and BC emission profiles reported in the EMEP/EEA 2016 Guidebook (Table 3.3) have been used. For neutral sulphite semi-chemical process the emission factors used through the time period referred for SO_x, NMVOC and NO_x to CORINAIR 1992, EMEP/CORINAIR Guidebook, and for NO_x from 1996, data were communicated by the operator of the plant.

NMVOC emissions include emissions from chipboard production where activity data are those in the FAOSTAT database for particle board and the emission factor 500 g/Mg product is from "Corinair 1992 Default Emission Factors Handbook".

In 2I category TSP emissions from wood processing are reported. Considering that in Italy wood furniture production start from wood panels and sawnwood, emissions are estimated on the basis of statistics from the FAOSTAT database for that kind of wood production and the emission factor in the 2019 EMEP/EEA Guidebook, which is equal to 1 kg/t of wood product.

In 2L category lead emissions from batteries manufacturing are reported. Activity data are provided by the non-ferrous metal industrial association (ASSOMET) and refer to the amount of lead used for the batteries production; the emission factor has been provided by the relevant industrial association (ANIE) calculated on the basis of average lead concentration to the chimney, equal to 0.2 mg/Nmc, the average flow (equal to 15 Nmc/h/tonnes Pb) and the annual number of hours.

4.3 TIME SERIES AND KEY CATEGORIES

The following sections present an outline of the main key categories, and relevant trends, in the industrial process sector. Table 4.2 reports the key categories identified in the sector.

Table 4.2 Key categories in the industrial processes sector in 2022.

	2A1	2A2	2A5a	2A5b	2B1	2B2	2B3	2B6	2B7	2B10a	2C1	2C2	2C3	2C7c	2G	2H1	2H2	2L	
	%																		
SO _x					0.00			0.16	0.17	5.90	1.40				0.03				
NO _x					0.01	0.04	0.00	0.00	0.02	0.35	0.37				0.02				
NH ₃					0.00	0.00			0.07	0.01					0.08				
NMV OC					0.00					0.19	0.36				1.15	0.16	3.31		
CO					0.00				0.37	0.62	2.10				0.21				
PM10	1.78	0.51	8.42	6.36				0.00	0.01	0.14	1.78				1.19		0.01		
PM2.5	1.41	0.14	1.20	0.91				0.00	0.01	0.07	2.13				1.43				
BC	0.34	0.01						0.00	0.00	0.01	0.06								
Pb											30.76				2.96				0.97
Cd										1.05	22.52				8.45				
Hg											49.17								
PAH											11.81				0.01				
Dioxin											26.63				0.00				
HCB																			
PCB											71.27								

Note: key categories are shaded in blue

There is a general reduction of emissions in the period 1990 - 2022 for most of the pollutants due to the implementation of different directives at European and national level. A strong decrease is observed especially in the chemical industry due to the introduction of relevant technological improvements and the complicated international situation which led to the considerable reduction of some productions.

4.3.1 Mineral products (2A)

As mentioned above, TSP, PM10 and PM2.5 emission factors for cement production are set constant from 1990 to 2020 while since the last submission SO₂ emissions have been set as not occurring along the whole timeseries as they originate from fuel combustion and consequently, they are allocated under 1A2f. The trends of TSP, PM10 and PM2.5 emissions follow that of the activity data. In Table 4.3, activity data, TSP, PM10 and PM2.5 emissions from cement production (2A1) are reported.

Table 4.3 Activity data and PM10 emissions from cement production, 1990 – 2022 (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Cement production [Gg]	42,414	35,432	41,119	47,291	34,283	20,825	19,241	18,060	20,621	18,797
Clinker production [Gg]	29,785	28,778	29,816	33,122	25,239	15,527	15,119	13,389	15,162	15,665
TSP emissions (Gg)	0.008	0.007	0.008	0.009	0.007	0.006	0.004	0.003	0.004	0.004
PM10 emissions [Gg]	0.007	0.007	0.007	0.008	0.006	0.006	0.004	0.003	0.004	0.004
PM2.5 emissions [Gg]	0.004	0.004	0.004	0.004	0.003	0.003	0.002	0.002	0.002	0.002

In Table 4.4, activity data, TSP, PM10 and PM2.5 emissions from quarrying and mining of other mineral than coal (2A5a) are reported.

Table 4.4 AD, estimates and significance on the national totals for PM10 and PM2.5 from 2A5a (1990-2022)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Annual production [Mt]	187	187	213	193	162	156	139	103	177	346
TSP emissions [Gg]	19	19	22	20	16	16	14	10	18	35
PM10 emissions [Gg]	9	9	11	10	8	8	7	5	9	17
PM2.5 emissions [Gg]	1	1	1	1	1	1	1	1	1	2

In Table 4.5 activity data, TSP, PM10 and PM2.5 emissions from construction and demolition of buildings and roads (2A5b) are reported.

Table 4.5 AD, estimates for TSP, PM10 and PM2.5 from 2A5b (1990-2022)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Buildings+roads [1000 m2]	88,813	89,037	77,421	100,685	110,213	53,571	44,350	54,167	44,116	30,086
TSP emissions (Gg)	0.125	0.128	0.110	0.139	0.174	0.085	0.068	0.086	0.067	0.043
PM10 emissions [Gg]	0.038	0.039	0.033	0.042	0.052	0.026	0.020	0.026	0.020	0.013
PM2.5 emissions [Gg]	0.004	0.004	0.003	0.004	0.005	0.003	0.002	0.003	0.002	0.001

4.3.2 Chemical industry (2B)

Other chemical industry (2B10a) was a key category for Cd and Hg emissions in 1990 and for SO_x in 2022 and for Hg at trend assessment. Hg emissions refer to chlorine production with mercury cells process; in Table 4.6, activity data and Hg emissions from chlorine production are reported. As reported in paragraph 4.1, to estimate emissions from 1990 to 2001, the average emission factor supplied by EUROCHLOR for western Europe chlor-alkali production plants has been used, while from 2002 emission data have been supplied directly from the production plants in the framework of the national EPER/E-PRTR reporting obligation. The average emission factor decreased from 1.11 g Hg/t in 2002 to zero in 2018. The reduction observed and zero Hg emissions since 2018 is a consequence of the conversion of production plants from the mercury cells process to the membrane technology but it depends also on suspensions of production processes at some facilities.

Table 4.6 Activity data and Hg emissions from chlorine production, 1990 – 2022

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Activity data [Gg]	1,043	869	786	535	258	218	249	241	236	237
Hg emissions [Mg]	3	2	1	0.48	0.12	0.04	-	-	-	-

SO_x emissions are prevalently from carbon black production. Sulphuric acid production, titanium oxide, other sulphate and phthalic anhydride productions are other sources reported in 2B10a and emitting SO_x. Activity data and emission factors for these sources are collected at plant level on annual basis.

4.3.3 Metal production (2C)

Emission trend of HMs, PCB and PCDD/PCDF is driven mainly by the electric arc furnaces iron and steel production which increased from 15.1 Mt in 1990 to 19.6 Mt in 2008; in 2009, because of the economic crisis, steel production from electric arc has decreased substantially and since 2010 the production has increased again up to 20.0 Mt in 2018 and 20.4 Mt in 2021, with a dip in 2020 to 17.01 Mt because of the pandemic. In 2022 EAF production is equal to 18.2 Mt. In Table 4.7, activity data and HM, PCB and PCDD/PCDF emissions from electric arc furnace (EAF) and from the whole sector 2C1 are reported, but dioxins emissions from sinter plant are reported in the energy sector in 1.A.2f category. In 2022 average emission factor is equal to 0.22 micrograms TEQ per Mg of sinter produced.

Table 4.7 Activity data and HMs, PCB and PCDD/PCDF emissions from electric arc furnace and 2C1, 1990 – 2022

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Steel production EAF [kt]	15,102	16,107	15,879	17,661	17,115	17,255	18,980	17,007	20,416	18,190
Cd emissions EAF [Mg]	1.1	1.1	0.8	0.9	0.9	0.9	0.9	0.9	1.0	0.9
Cd emissions 2C1 [Mg]	1.3	1.4	1.1	1.2	1.1	1.0	1.0	0.9	1.0	0.9
Hg emissions EAF [Mg]	2.3	2.4	2.4	2.6	2.6	2.6	2.8	2.6	3.1	2.7
Hg emissions 2C1 [Mg]	2.3	2.5	2.4	2.7	2.6	2.6	2.9	2.6	3.1	2.7
Pb emissions EAF [Mg]	52.1	55.6	54.8	60.9	59.0	59.5	65.5	58.7	70.4	62.8
Pb emissions 2C1 [Mg]	61.1	65.7	64.1	71.0	66.5	63.7	65.8	58.9	70.7	63.0
PCB emissions EAF [kg]	54.4	58.0	57.2	63.6	61.6	62.1	68.3	61.2	73.5	65.5
PCB emissions 2C1 [kg]	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
PCDD/F emissions EAF [g T-eq]	67.2	71.7	70.7	78.6	76.2	76.8	84.5	75.7	90.8	80.9
PCDD/F emissions 2C1 [g T-eq]	67.2	71.7	70.7	78.6	76.2	76.8	84.5	75.7	90.8	80.9

For Pb and Hg, the same EFs have been used for the whole time series (derived by the EMEP/CORINAIR Guidebook), while for Cd a national emission factor, equal to 50 mg/t, was available thanks to a sectoral study (APAT, 2003) and refers to the years after 1997. This study shows range < 1-54 mg/t and the value set to 50 mg/t was chosen for conservative reason being more consistent with the old one; this value should include technology progresses occurred in the iron and steel production activities in those years. Lacking information for the years backwards, the default CORINAIR EF was used. For PCB and PCDD/Fs, emission factors are constant from 1990 to 2022 and emission trends are ruled by the activity data. For SO₂ and PM emissions from lead, zinc and copper production they are included and reported in the

energy relevant sector. In Italy there is a sole integrated plant for the primary production of zinc and lead and this makes it difficult to ensure a good breakdown between the energy and the process sectors and the activities. During the latest year more information about the plant has been supplied taking advantage of a direct contact with the facility through the E-PRTR registry but it was not sufficient to split the emissions. Thanks to a deep survey on this category based on the analysis of IPPC permits and data available from the monitoring and control plan, the estimates for Pb, Cd, Zn, PM, NO_x e SO₂ have been revised since 2014. The analysis of this documentation will carry out further improvement such as, for example, a better allocation of emissions between combustion and process but also between the zinc and the lead production. Further information in the paragraph 4.2.3 and in the non-ferrous combustion paragraph.

Following the decision 2012/17 of the Executive Body of the Convention on Long Range Transboundary Air Pollution, that requests Italy to submit information concerning the status and details of its work to improve the emission inventory of PAH, Italy in recent years has reviewed the estimates regarding PAH major sources. In the 2013 submission different recalculations have been performed in the energy and waste sector, emissions from iron and steel production have been revised in the 2014 submission. The most important update regards pig iron tapping emission factor that considers, from 2000 onwards, the abatement due to fabric filters and the relevant EF derived from the Guidebook EMEP/CORINAIR 2006 (0.95 g/Mg). Investigations on the largest integrated plant in Italy confirmed the installation of fabric filters on each point of emission related to pig iron tapping (MATTM, 2011). As regards EAF too, EF was update on the basis of a sectoral study (APAT, 2003) which reports the development of abatement technologies in the '90s in Italy and the consequent evolution in the plants with the installation of fabric filters; but in this case the update is referred to 1990-1999 because the EF used in previous submissions concerned already abated emissions. In Table 4.8, activity data and PAH emissions from integrated plants and from the whole sector 2C1 are reported.

Table 4.8 Steel production data and PAH emissions from integrated plants, 1990 – 2022

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Pig iron production [Gg]	11,852	11,678	11,209	11,424	8,555	5,051	4,619	3,406	4,111	3,390
Steel production BOF [Gg]	10,365	11,664	10,744	11,688	8,635	4,763	4,211	3,371	3,996	3,409
PAH emissions i.p.* [Mg]	41.9	41.3	11.7	12.1	9.2	5.8	5.4	4.1	5.0	4.2
PAH emissions 2C1 [Mg]	44.9	44.5	14.3	15.1	11.9	8.2	8.0	6.4	7.7	6.6

4.3.4 Other production (2G – 2H – 2I – 2L)

These categories include use of fireworks, lubricants and tobacco (2G), Pulp & paper and food & drink (2H), wood processing (2I) and batteries manufacturing (2L). NMVOC emissions from these categories derive from 2G and 2H and it is equal to 38.1 Gg accounting for 4.6% of the national total, in particular emissions from food and drink (2H2) accounting for 3.3% of the national total. Emissions from this last category refer to the processes in the production of bread, wine, beer and spirits. Emission factors are assumed constant for the whole time series. In Table 4.9, activity data and NMVOC emissions from sector are reported.

Table 4.9 Activity data and total NMVOC emissions, 1990 – 2022


	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
2G Explosives/fireworks – [Mg]	3,122	3,122	10,686	13,108	10,748	5,480	4,289	4,762	9,165	7,741
2G Use of tobacco – [Mg]	92,573	92,573	101,973	94,674	90,721	79,823	72,619	70,030	69,354	70,048
2G Lubricants – [Gg]	673	630	638	580	429	404	396	338	372	327
2H Pulp and paper – [Gg]	140.4	82.2	80.0	80.4	NO	NO	NO	NO	NO	NO

		1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
2H	Activity data - Bread [Gg]	4,153	3,882	3,565	4,109	4,161	3,877	3,796	4,666	4,114	4,250
2H	Activity data – Wine [10 ⁶ dm ³]	5,521	5,620	5,409	5,057	4,673	5,073	5,219	5,433	5,319	5,636
2H	Activity data – Beer [10 ⁶ dm ³]	1,215	1,199	1,258	1,280	1,281	1,429	1,725	1,583	1,764	1,836
2H	Activity data – Spirits [10 ⁶ dm ³]	268	232	206	161	115	98	108	111	106	122
TOT	NMVOE emissions [Gg]	52.5	48.6	46.8	45.9	39.9	37.2	37.4	39.7	38.0	38.1

4.4 QA/QC AND VERIFICATION

Activity data and emissions reported under EU-ETS and the national EPER/EPRTTR register are compared to the information provided by the industrial associations. The general outcome of this verification step shows consistency among the information collected under different legislative frameworks and information provided by the relevant industrial associations. Every five years emissions referring to 1990-2015 are disaggregated at regional and provincial level and figures are compared with results obtained by regional bottom-up inventories. From 2015 onwards the disaggregation at local level takes place every four years, so up to now the disaggregation of the national inventory covers estimates for the years: 1990, 1995, 2000, 2005, 2010, 2015 and 2019. PM10 emissions disaggregated at local level are also used as input for air quality modelling. The distribution of PM10 emissions from the industrial processes sector at NUTS3 level for 2019 is reported in Figure 4.1; methodologies are described in the relevant publication (ISPRA, 2022).

Figure 4.1 PM10 emissions from industrial processes in 2019 (t)



4.5 RECALCULATIONS

4.5.1 Mineral industry (2A)

Recalculations occur for PM2.5 from category 2.A.5.a quarrying and mining, along the whole time series; due to a reporting mistake in the previous submission.

4.5.2 Chemical industry (2B)

Recalculations occur in 2017 because of a mistake in BC emission factor in Polyvinylchloride and Acrylonitrile Butadiene Styrene (ABS) resins production and in 2021 because of the update of activity data.

4.5.3 Metal industry (2C)

No recalculations occur for this source category.

4.5.4 Other product use (fireworks and tobacco) (2G)

Recalculation occurred for NMVOC emissions from lubricants use because of the update of the activity data for 2012-2021 and from fireworks for 2020-2021.

4.5.5 Other industrial processes (2H)

No recalculations occur for this source category.

4.6 PLANNED IMPROVEMENTS

Activities 2C3, 2C5, 2C6 and 2C7 are under investigations to allocate emissions between combustion and process.

5 IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2)

5.1 OVERVIEW OF THE SECTOR

In this sector all non-combustion emissions from other industrial sectors than manufacturing and energy industry are reported. Emissions are related to the use of solvent in paint application, degreasing and dry cleaning, chemical products, manufacture and processing and other solvent use, including emissions from road paving with asphalt and asphalt roofing activities. NMVOC emissions are estimated from all the categories of the sector as well as PM for polyester and polyvinylchloride processing, in the chemical product category, and for asphalt processes and PAH emissions from the preservation of wood in the other solvent use. The categories included in the sector are specified in the following.


- 2D3a Domestic solvent use includes emissions from the use of solvent in household cleaning and car care products as well as cosmetics.
- 2D3b Road paving with asphalt includes emissions from the production and use of asphalt for road paving.
- 2D3c Asphalt roofing includes emissions from the manufacturing of roofing products and the blowing of asphalt.
- 2D3d1 Decorative coating includes emissions from paint application for construction and buildings, domestic use and wood products.
- 2D3d2 Industrial coating includes emissions from paint application for manufacture of automobiles, car repairing, coil coating, boat building and other industrial paint application.
- 2D3e Degreasing includes emissions from the use of solvents for metal degreasing and cleaning.
- 2D3f Dry cleaning includes emissions from the use of solvent in cleaning machines.
- 2D3g Chemical products, manufacture and processing covers the emissions from the use of chemical products such as polyurethane and polystyrene foam processing, manufacture of paints, inks and glues, textile finishing and leather tanning.
- 2D3h Printing includes emissions from the use of solvent in the printing industry
- 2D3i Other product use addresses emissions from glass and mineral wool enduction, fat edible and non-edible oil extraction, preservation of wood, application of glues and adhesives, vehicles dewaxing.

NMVOC emissions from 2D3a, 2D3d, 2D3g, 2D3i and 2D3h are key categories in 2022; the same categories, except 2D3h, were also key categories in 1990. For the trend 1990-2022, 2D3a, 2D3g and 2D3h result as key categories.

The sector accounts, in 2022, for 40.3% of total national NMVOC emissions, whereas in 1990 the weight out of the total was equal to 30.3%. Total sectoral NMVOC emissions decreased by 44.5%, between 1990 and 2022. PM10 and PM2.5 account for 2.6% and 1.8%, respectively, in 2022. PAH emissions are also estimated but they account for less than 1%.

In Figure 5.1 the share of NMVOC emissions of the sector is reported for the years 1990 and 2022.

Figure 5.1 Share of NMVOC emissions for the solvent use sector in 1990 and 2022.



5.2 METHODOLOGICAL ISSUES

The sector is characterized by a multitude of activities which implies that the collection of activity data and emission factors is laborious. A lot of contacts have been established in different sectors with industrial associations and documentation has been collected even though improvements are still needed especially in some areas. Emissions of NMVOC from solvent use have been estimated according to the methodology reported in the EMEP/EEA guidebook, applying both national and international emission factors (Vetrella, 1994; EMEP/CORINAIR, 2007; EMEP/EEA, 2016). Country specific emission factors provided by several accredited sources have been used extensively, together with data from the national EPER/PRTR registry; in particular, for paint application (Offredi, several years; FIAT, several years), solvent use in dry cleaning (ENEA/USLRMA, 1995), solvent use in textile finishing and in the tanning industries (Techne, 1998; Regione Toscana, 2001; Regione Campania, 2005; GIADA, 2006). Basic information from industry on percentage reduction of solvent content in paints and other products has been applied to EMEP/EEA emission factors in order to evaluate the reduction in emissions during the considered period.

A more detailed description is reported for the 2022 key categories of NMVOC emissions in the following sections. A description of CO emissions estimation process for asphalt roofing is also included.

5.2.1 Domestic solvent use (2D3a)

The category comprises a lot of subcategories whose emissions, specifically NMVOC, originate from the use of solvent in household cleaning and car care products as well as cosmetics. Emissions from this category have been calculated using a detailed methodology, based on VOC content per type of consumer product. Emissions from domestic solvent use comprise emissions from the use of products for household and cleaning and for cosmetics which are derived as described in the following.

Activity data

Activity data are expressed as the sum, in tonnes, of household and cleaning products and cosmetics.

Household and cleaning products: data are communicated by the National Association of Detergents and Specialties for industry and home care (Assocasa, several years) either by personal communications or Association Reports and refer to the consumption of soaps and detergents and cleaning and maintenance products.

Cosmetics: data are the sum of cosmetics products in aerosol form and other cosmetics. Figures of cosmetics in aerosol form are provided by the Italian Aerosol Association (AIA, several years (a) and (b)) and refer to the number of pieces of products sold for personal care (spray deodorants, hair styling foams and other hair care products, shaving foams, and other products). These figures are then converted in tonnes by means of the capacity of the different cosmetics containers.

Figures for other cosmetics products are derived by the Production Statistics Database (Prodcom) supplied by the National Institute of Statistics (ISTAT, several years (a) and (b)) by difference with the previous aerosol data.

Time series of cosmetics production is reconstructed by means of the annual production index, considering the year 2000 as the base year because this is the year where production national statistics and Prodcom data coincide. The next step is the calculation of apparent consumption taking into account import-export data derived by the National Association of Cosmetic Companies (UNIPRO, several years). Since these figures also include aerosol cosmetics, the amount of aerosol cosmetics is subtracted.

Final consumption is therefore estimated.

Emission factors

NM VOC emission factors are expressed in percentage of solvent contained in products.

Household and cleaning products: figures are communicated by the relevant industrial association, ASSOCASA, by personal communications. For leather, shoes, wood etc. and car maintenance products, figures are taken from BiPro Association. For insecticides and disinfectants, emission factors derive from national studies at local level.

Cosmetics: for aerosol cosmetics, the emission factor is communicated by the Italian Aerosol Association for the year 2004 and supposed constant from 1995. For other cosmetics, information from BiPro has been considered (EC report 'Screening study to identify reductions in VOC emissions due to the restrictions in the VOC content of products', year 2002 (EC, 2002)), and supposed constant from 1996.

5.2.2 Road paving with asphalt (2D3b)

The category includes NM VOC and PM emissions from the application of asphalt to road. The emission factors for NM VOC emissions from road paving are set constant along the whole timeseries and are equal to 16 g NM VOC/Mg_{asphalt}. The activity data are yearly provided by the national association of producers while the EF is taken from the EMEP/EEA Guidebook.

5.2.3 Asphalt Roofing (2D3c)

The category includes CO and NM VOC emissions from the application of asphalt to insulate (waterproofing) the roof of buildings. The emission factors for CO and NM VOC emissions from asphalt roofing are set constant along the whole timeseries and are equal to 38 g CO/10³ m² and 0.085 g NM VOC/10³ m². The activity data are yearly provided by the national association of producers while the EF is taken from the EMEP/EEA Guidebook with the assumption that 1000 m² of asphalt material corresponds to 4 Mg. The following table describes AD, EF and emissions for CO and NM VOC between 1990 and 2022.

Table 5.1 AD, EF and emissions for CO and NM VOC from asphalt roofing, 1990-2022.

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Activity data – Asphalt roofing [10 ⁶ m ²]	200.0	200.0	200.0	240.0	204.0	152.2	122.9	117.6	134.4	130.6
CO emissions [Gg]	0.008	0.008	0.008	0.009	0.008	0.006	0.005	0.004	0.005	0.005
NM VOC emissions [Gg]	0.017	0.017	0.017	0.020	0.017	0.013	0.010	0.010	0.011	0.011

5.2.4 Decorative coating (2D3d1)

The category includes NMVOC emissions from the application of paint for construction and buildings, domestic use and wood products. Activity data on the consumption of paint for construction and buildings and related domestic use are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated on the basis of production figures provided annually by the National Institute of Statistics (ISTAT, several years [a] and [b]). From 2007 onwards, data are also provided by SSOG (Stazione Sperimentale per le industrie degli Oli e dei Grassi, Experimental Station for Oils and Fats Industries), which collects information and data regarding national production and imports for paint categories set out in the directive 2004/42/EC on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products. The purpose of this directive is to limit the total content of VOCs in certain paints and varnishes and vehicle refinishing products in order to prevent or reduce air pollution resulting from the contribution of VOCs to the formation of tropospheric ozone. The directive sets maximum VOCs content limit values for some paints and varnishes. As for emission factors, those for construction and buildings are taken from the EMEP/EEA guidebook and are considered constant till 2009, whereas the default values for domestic use vary in consideration of the different share between solvent and water content in paint throughout the years. In particular, the variation of emission factor from 1990 to 2000 is equal to 35% - 65% up to 25% - 75% in 2000, on the basis of qualitative information supplied by industry on the increase of water based paints products in the market. From 2010, emission factors are calculated taking into account maximum VOC content limit values for paint and varnishes set out in Annex II A of Directive 2004/42/EC and data collected by SSOG. The comparison of national emission estimates for this category with those produced by IIASA for 2010 resulted in similar values. On the other hand, information on activity data and emission factors for emissions from wood products are provided by the national association of wood finishing (Offredi, several years). Emission factors have been calculated for 1990, 1998 and 2003 on the basis of information provided by the industrial association distinguishing the different type of products which contain different solvent percentages. Data have been supplied also for the years 2005 and 2006. Actually, we are keeping constant the 2006 value unless the association provides us with updated information. For previous years, values have been interpolated.

In this category, emissions from paint application in construction and buildings is the largest contributor to national NMVOC emissions and the relevant share has grown considerably in recent years. NMVOC emissions due to the use of paint and other products except from industrial coating could not be controlled properly in the past since the EU Directive 2004/42/EC entered into force. This directive, transposed into the Italian legislation in 2004, sets out maximum VOC content for many paint, varnishes and vehicle refinishing products that had to be achieved in two steps. The early limit values, to be respected from 2007 till 2009, did not lead to a significant reduction of NMVOC emissions, while the latest values, that had to be respected from 2010 onwards, brought to a significant decrease.

5.2.5 Industrial coating (2D3d2)

The category includes emissions from paint application for manufacture of automobiles, car repairing, coil coating, boat building and other industrial paint application. Activity data on the number of vehicles are provided by the National Automobile Association (ACI, several years) in the Annual Statistical Report and the emission factors are those reported by the main automobile producers on the relevant activity in their environmental reports and communicated from 2003 in the framework of E-PRTR. For the paint used in car repairing, activity data are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated on the basis of production figures provided annually by the National Institute of Statistics (ISTAT, several years [a] and [b]). The default emission factor (provided by the EMEP guidebook) used from 1990 to 1995 equal to 700 g/kg paint is also confirmed by the European guidelines for car repairing provided by the Conseil Europeen de l'Industrie des Peintures (CEPE, 1999). The reduction of the emission factor in 1999 (13% of 1995) is applied on the basis of information on different shares between solvent and water based paint throughout the years provided by the national study PINTA, Piano nazionale di tutela della qualità dell'aria (ENEA, 1997). From 1996 to 1999 the reduction is linear. From 1999 to 2006 the value is kept constant. From 2007 onwards emission factors have been calculated taking into account the maximum VOC content limit values for paint and varnishes set out in Annex II B of

Directive 2004/42/EC and data collected by SSOG. The Italian implied emission factor is the weighted average of the different products used in this activity where data are collected at detailed level and communicated within the European Directive. The trend is driven by the increase in the last years of the use of primers and special finishes. Similar trend is noted for the construction and building and domestic paints where the variability is mainly due to the percentage of solvent based paint product used out of the total paints.

Concerning coil coating, boat building and other industrial paint application, activity data are provided by the Ministry of Productive Activities for 1990 and 1991 (MICA, 1999) and updated annually by the National Institute of Statistics (ISTAT, several years [a] and [b]). Emission factors are taken from the EMEP guidebook considering the national legislation where relevant. Emission factors of the other industrial paint application from 1990 to 1995 are constant and derive from the 1999 EMEP/CORINAIR guidebook. The reduction of the emission factor from 1996 to 2004 is applied on the basis of information on different share of paints throughout the years provided by the national study PINTA. From 2010, the value of the 1999 Guidebook has been chosen considering the further reduction of the sector (in PINTA, the reduction for 2005 with respect to 1995 is equal to 37%, and for 2010 64%; considering the default emission factor 250 g/kg of paint, the reduction is equal to 53%). NMVOC emissions from this category have been decreasing constantly since the nineties, when all industrial installations have been subjected to permits from local authorities. Since then, most of the installations have to comply with emission limit values and technological requirements imposed at regional level, taking into account the EU directives on industrial emissions (i.e. Directive 99/13/EC on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations (EC, 1999)) and often going beyond the European legislation. With regard to car repairing the emission cut from 2007 onwards is mainly due to the maximum contents of VOC set by EU Directive 2004/42/EC (EC, 2004).

5.2.6 Degreasing (2D3e)

NMVOC emissions have been estimated for this category. The emission factor used is constant along the time series and equal to 700 g NMVOC/kg solvent used, as specified in the EMEP/EEA Guidebook. According to the information provided by the National Industrial Association, due to technological improvements, the amount of solvent used in the products decreased during the period whereas it has been assumed that the percentage of NMVOC emissions remains constant. Activity data, solvent used, are also provided by the relevant industrial association (Federchimica, several years).

Table 5.2 AD, EF and emissions for NMVOC from metal degreasing 2D3e, 1990-2022.

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Activity data – Degreasing [Gg]	52.8	32.8	25.9	22.2	19.1	16.4	14.5	14.1	13.7	13.2
NMVOC emissions [Gg]	36.9	22.9	18.1	15.6	13.4	11.5	10.2	9.9	9.6	9.3

5.2.7 Dry cleaning (2D3f)

Concerning dry cleaning, activity data, equal to 30,000 machines, remain unchanged throughout the time series and the emission factor is calculated based on the allocation of machines to closed-circuit (CCM) and open-circuit (OCM). Different amounts of solvent are used in these machines and have different emission factors. The emission factors are calculated assuming that in 1990 the closed-circuit machines were 60%, 90% in 1995 and in up to 100% in 1999.

The average consumption of solvent per machine is equal to 258 kg/year for CCM and 763 kg/year for OCM, as derived from a national study by ENEA/USL-RMA (ENEA/USL-RMA, 1995). It is assumed that only perchlorethylene is used. These values are multiplied by the emission factors of the Guidebook EMEP, referred to the amount of solvent consumed (equal to 0.4 and 0.8 kg/kg of solvent, for CCM and OCM, respectively) and then the average annual emission factor was calculated based on the percentage distribution of closed and open circuit machines.

5.2.8 Chemical products, manufacture and processing (2D3g)

The category comprises emissions from the use of chemical products such as polyester, polyurethane, polyvinylchloride and polystyrene foam processing, manufacture of paints, inks and glues, textile finishing and leather tanning.

Activity data for polystyrene and polyurethane are derived from the relevant industrial associations, and ISTAT (ISTAT, several years [a] and [b]), whereas emission factors are from the EMEP/CORINAIR guidebook. For what concerns polyurethane, the relevant national industrial association has communicated that the phase out of CFC gases occurred in the second half of nineties and the blowing agent currently used is penthane. Because of manufacturing plant have abatement system in place PM emissions could all be considered as PM2.5.

As for polyvinylchloride (PVC), activity data and emission factors are supplied in the framework of the national PRTR. NMVOC emissions are entirely attributed to the phase of PVC production; no use of solvents occurs in the PVC processing. This information has been provided by the relevant industrial plant, EVC Italy, in 2001. Because of manufacturing plant have abatement system in place PM emissions could all be considered as PM2.5.

For the other categories, activity data are provided by the relevant industrial associations and by ISTAT, while emission factors are taken from the EMEP/CORINAIR guidebook considering national information on the solvent content in products supplied by the specific industrial associations.

As regard rubber processing, emission factors for the first years of nineties have been provided by the industrial association. The use of the Swedish emission factor from 1997 was justified in lack of other updated data.

For the glues manufacturing category, emission factors for 1990 are derived from the 1992 EMEP/CORINAIR guidebook. The trend of emission factor is estimated on the basis of the trend of the emission factor for consumption of glue (as indicated by the industrial association). From 1995 to 2004, the industrial association communicated data on consumption and solvent content by product. The reductions from 2000 are based on the assumptions of PINTA. From 2004 the emission factor has been assumed constant in lack of updated information. For previous years, values have been interpolated.

As regards leather tanning, emission factor for 1990 is from Legislative Decree 152/2006, equal to the maximum VOC content limit value (150 g/m²). For 2000 and 2003, emission factors have been calculated on the basis of emission figures derived by the national studies on the major leather tanning industries and statistical production.

As regards asphalt blowing and possible PAH and Benzo(a)pyrene emissions as suggested by the 2016 EMEP/EEA Guidebook, according to the relevant industrial association PAH emissions are negligible because all the asphalt blowing plants have abatement filter system of PM and afterburners of gas. Moreover, these plants should respect national environmental legislation not exceeding at the stack more than 0.1mg/Nm³ for total PAH. For this pollutant the relevant notation key NE has been used.

No PM emission factors are provided in the EMEP EEA 2019 GB for these activities, so after checking with the relevant industries the values of the EFs have been differentiated for the activities concerned with PM10 assumed as 80% of TSP and PM2.5 set equal to 60% of TSP (as suggested in the 2019 EMEP/EEA GB, 2B Chemical industry, paragraph 3.2.2.1, pag.14).

Table 5.3 EF and emissions for PM2.5 from metal Chemical products 2D3g, 1990-2022.

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
EF PM10 polyester proc. [g PM10/Mg]	92	92	92	92	92	92	92	92	92	92
EF PM10 polyvinylchloride proc. [g PM10/Mg]	24.0	20.8	9.2	8.8	8.8	8.8	8.8	8.8	8.8	8.8
EF PM2.5 polyester proc. [g PM2.5/Mg]	69	69	69	69	69	69	69	69	69	69

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
EF PM2.5 polyvinylchloride proc. [g PM2.5/Mg]	18	15.6	6.9	6.6	6.6	6.6	6.6	6.6	6.6	6.6
Total PM10 emissions [Gg]	0.03	0.03	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Total PM2.5 emissions [Gg]	0.02	0.02	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01

5.2.9 Other product use (2D3i)

The category includes NMVOC emissions from the application of glues and adhesives, which account for most of emissions from the category, emissions from fat, edible and non-edible oil extraction and minor emissions from glass wool enduction.

Activity data and emission factors for the application of glues and adhesives had been provided by the relevant industrial association up to 2004. After that period, activity data have been updated on the basis of information by ISTAT (ISTAT, several years [a] and [b]) whereas the emission factor is considered constant in absence of further information.

For fat, edible and non-edible oil extraction activity data derive from the FAOSTAT database (<http://faostat.fao.org>) whereas default NMVOC emission factors do not change over the period and are equal to 1.57 g/kg seeds. PM emissions are also estimated in this category.

5.3 TIME SERIES AND KEY CATEGORIES

The sector accounts, in 2022, for about 40.3% of total national NMVOC emissions. PM10 and PM2.5 account for 2.6% and 1.8%, respectively. PAH emissions are also estimated in this sector but they account for less than 1%.

NMVOC emissions from the use of solvent decreased from 1990 to 2022 of about 44.5%, from 596.5 Gg in 1990 to 331.3 Gg in 2022, mainly due to the reduction of emissions in paint application, in other use of solvent and in degreasing and dry cleaning activities. The general reduction observed in the emission trend of the sector is due to the implementation of the European Directive 1999/13/EC (EC, 1999) on the limitation of emissions of volatile organic compounds due to the use of organic solvents, entered into force in Italy in January 2004, and the European Directive 2004/42/EC (EC, 2004), entered into force in Italy in March 2006, which establishes a reduction of the solvent content in products. Moreover, the reduction of emissions from paint application, is also due to the implementation of the Italian Legislative Decree 161/2006.

Figure 5.2 shows emission trends from 1991 to 2022 with respect to 1990 by sub-sector.

The main source of emissions is paint application (2D3d) where NMVOC emissions derive mainly from the categories other industrial paint application and construction and buildings. The second source of emissions is domestic solvent use (2D3a), mostly for the consumption of cosmetics and cleaning products, followed by chemical products and other product use (2D3g), especially for emissions deriving from polyurethane processing, paints manufacturing and leather tanning.

Figure 5.2 Trend of NMVOC emissions from 1991 to 2022 as compared to 1990




Table 5.4 represents the pollutants estimated in the sector and the key categories identified.

Table 5.4 Key categories in the IPPU - Solvent and other product use sector in 2022

	2D3a	2D3b	2D3c	2D3d	2D3e	2D3f	2D3g	2D3h	2D3i
SO _x									
NO _x									
NH ₃									
NMVOC	11.86	0.06	0.00	15.00	1.13	0.38	6.33	2.24	3.27
CO			0.00						
PM10		0.91	0.03				0.004		1.69
PM2.5		0.17	0.01				0.005		1.61
BC		0.08	0.00						
Pb									
Cd									
Hg									
PAH									0.02
Dioxin									
HCB									

2D3a	2D3b	2D3c	2D3d	2D3e	2D3f	2D3g	2D3h	2D3i
PCB								

Note: key categories are shaded in blue

In Table 5.5 and Table 5.6 activity data and emission factors used to estimate emissions from the sector are reported at SNAP code level.

A strong decrease in the content of solvents in the products in the nineties is observed.

Table 5.5 Activity data in the IPPU - Solvent and other product use sector

			1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
06 01	Paint application											
06 01 01	Paint application : manufacture of automobiles	<i>vehicles</i>	2,865,857	2,521,355	2,770,104	1,766,930	1,310,425	1,325,327	1,271,513	1,098,012	1,170,208	1,209,905
06 01 02	Paint application : car repairing	<i>Mg paint</i>	22,250	17,850	24,276	23,475	19,479	25,395	37,416	16,899	9,873	13,492
06 01 03	Paint application : construction and buildings (except item 06.01.07)	<i>Mg paint</i>	111,644	120,736	125,928	163,455	168,358	158,661	196,105	177,383	178,811	180,873
06 01 04	Paint application : domestic use (except 06.01.07)	<i>Mg paint</i>	420,000	420,000	420,000	420,000	420,000	420,000	420,000	420,000	420,000	420,000
06 01 05	Paint application : coil coating	<i>Mg paint</i>	14,500	14,500	14,500	14,500	14,500	14,500	14,500	14,500	14,500	14,500
06 01 06	Paint application : boat building	<i>Mg paint</i>	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000
06 01 07	Paint application : wood	<i>Mg paint</i>	150,000	150,000	140,000	140,000	123,250	80,000	75,000	70,000	65,000	67,000
	Other industrial paint application	<i>Mg paint</i>	125,000	125,000	125,000	125,000	125,000	125,000	125,000	125,000	125,000	125,000
06 02	Degreasing, dry cleaning and electronics											
06 02 01	Metal degreasing	<i>Mg solvents</i>	52,758	32,775	25,895	22,237	19,095	16,398	14,517	14,081	13,659	13,249
06 02 02	Dry cleaning	<i>machines</i>	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000
06 03	Chemical products manufacturing or processing											
06 03 01	Polyester processing	<i>Mg product</i>	179,852	197,882	168,704	112,188	89,638	94,389	101,045	101,571	102,925	100,580
06 03 02	Polyvinylchloride processing	<i>Mg product</i>	617,600	575,600	405,285	348,497	0	0	0	0	0	0
06 03 03	Polyurethane processing	<i>Mg product</i>	145,700	230,633	350,187	175,278	247,350	245,460	231,545	240,631	235,007	229,757
06 03 04	Polystyrene foam processing (c)	<i>Mg product</i>	85,004	80,400	90,200	35,200	33,692	46,800	51,600	42,600	57,400	47,600
06 03 05	Rubber processing	<i>Mg product</i>	671,706	700,859	810,124	831,187	607,667	545,989	603,318	622,427	609,324	545,989
06 03 06	Pharmaceutical products manufacturing	<i>Mg product</i>	80,068	88,094	104,468	106,861	110,183	120,907	136,202	135,174	134,933	146,298
06 03 07	Paints manufacturing	<i>Mg product</i>	697,129	747,417	900,683	964,631	891,882	851,450	885,576	943,407	959,584	970,653
06 03 08	Inks manufacturing	<i>Mg product</i>	87,527	110,667	132,256	132,521	133,979	108,600	112,953	120,329	122,393	123,805
06 03 09	Glues manufacturing	<i>Mg product</i>	111,683	266,169	302,087	331,770	317,560	249,152	286,629	276,060	280,794	249,152
06 03 10	Asphalt blowing	<i>Mg product</i>	77,248	70,336	77,408	88,896	65,000	25,000	18,000	16,000	17,000	12,000
06 03 12	Textile finishing	<i>1000 m2</i>	1,332,679	1,301,105	1,173,047	987,705	831,236	631,573	543,469	518,522	569,048	517,890
06 03 13	Leather tanning	<i>1000 m2</i>	173,700	183,839	200,115	157,891	186,824	162,500	139,831	129,350	146,412	177,124
06 04	Other use of solvents and related activities											
06 04 01	Glass wool enduction	<i>Mg product</i>	105,029	119,120	139,421	129,958	115,332	86,929	99,552	88,319	105,574	104,898
06 04 02	Mineral wool enduction	<i>Mg product</i>	0	11,000	18,000	20,500	0	0	0	0	0	0
06 04 03	Printing industry	<i>Mg ink</i>	73,754	91,667	100,690	111,550	98,206	79,604	96,424	102,721	104,483	105,688
06 04 04	Fat, edible and non edible oil extraction	<i>Mg product</i>	3,476,760	3,655,506	3,100,397	3,669,261	3,469,017	3,088,122	4,436,953	4,487,458	4,365,373	3,868,424
06 04 05	Application of glues and adhesives	<i>Mg product</i>	98,500	234,751	266,996	292,687	280,150	219,801	252,863	243,540	247,716	219,801
06 04 06	Preservation of wood	<i>Mg product</i>	9,000	9,000	9,000	9,000	9,000	9,000	9,000	9,000	9,000	9,000
06 04 08	Domestic solvent use (other than paint application)(k)	<i>Mg product</i>	1,938,779	2,282,020	2,410,338	2,767,759	2,614,274	2,265,605	2,376,579	2,423,461	2,640,959	2,773,958
06 04 09	Vehicles dewaxing	<i>vehicles</i>	2,540,597	1,740,212	2,361,075	2,238,344	1,972,070	1,594,259	1,949,554	1,441,385	1,519,936	1,336,310

Table 5.6 Emission factors in the IPPU - Solvent and other product use sector

			1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
06 01	Paint application											
06 01 01	Paint application : manufacture of automobiles	<i>g/vehicles</i>	8,676	6,296	4,833	4,065	2,854	3,037	2,614	2,685	2,455	2,116
06 01 02	Paint application : car repairing	<i>g/Mg paint</i>	700,000	700,000	605,500	605,500	497,810	617,377	527,187	434,596	526,266	479,153
06 01 03	Paint application : construction and buildings (except item 06.01.07)	<i>g/Mg paint</i>	300,000	300,000	300,000	300,000	200,000	152,412	209,310	184,193	202,612	170,194
06 01 04	Paint application : domestic use (except 06.01.07)	<i>g/Mg paint</i>	126,450	113,100	99,750	99,750	67,710	54,360	72,516	62,370	66,375	66,375
06 01 05	Paint application : coil coating	<i>g/Mg paint</i>	200,000	200,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000	10,000
06 01 06	Paint application : boat building	<i>g/Mg paint</i>	750,000	750,000	622,500	475,417	340,000	340,000	340,000	340,000	340,000	340,000
06 01 07	Paint application : wood	<i>g/Mg paint</i>	446,500	425,000	406,300	390,750	377,250	354,000	333,250	313,750	313,750	313,750
	Other industrial paint application	<i>g/Mg paint</i>	530,000	530,000	439,900	337,583	250,000	250,000	250,000	250,000	250,000	250,000
06 02	Degreasing, dry cleaning and electronics											
06 02 01	Metal degreasing	<i>g/Mg solvents</i>	700,000	700,000	700,000	700,000	700,000	700,000	700,000	700,000	700,000	700,000
06 02 02	Dry cleaning	<i>g/machines</i>	306,000	154,000	103,000	103,000	103,000	103,000	103,000	103,000	103,000	103,000
06 03	Chemical products manufacturing or processing											
06 03 01	Polyester processing	<i>g/Mg product</i>	325	325	325	325	325	325	325	325	325	325
06 03 02	Polyvinylchloride processing	<i>g/Mg product</i>	0	0	0	0	0	0	0	0	0	0
06 03 03	Polyurethane processing	<i>g/Mg product</i>	120,000	110,000	60,000	60,000	60,000	60,000	60,000	60,000	60,000	60,000
06 03 04	Polystyrene foam processing (c)	<i>g/Mg product</i>	60,000	60,000	60,000	60,000	60,000	60,000	60,000	60,000	60,000	60,000
06 03 05	Rubber processing	<i>g/Mg product</i>	12,500	10,000	8,000	8,000	8,000	8,000	8,000	8,000	8,000	8,000
06 03 06	Pharmaceutical products manufacturing	<i>g/Mg product</i>	55,000	55,000	55,000	55,000	55,000	55,000	55,000	55,000	55,000	55,000
06 03 07	Paints manufacturing	<i>g/Mg product</i>	15,000	15,000	15,000	13,110	10,863	9,524	10,769	9,686	9,797	9,500
06 03 08	Inks manufacturing	<i>g/Mg product</i>	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000	30,000
06 03 09	Glues manufacturing	<i>g/Mg product</i>	20,000	5,041	3,603	2,806	2,806	2,806	2,806	2,806	2,806	2,806
06 03 10	Asphalt blowing	<i>g/Mg product</i>	544	544	544	544	544	544	544	544	544	544
06 03 12	Textile finishing	<i>g/1000 m2</i>	296	296	296	296	296	296	296	296	296	296
06 03 13	Leather tanning	<i>g/1000 m2</i>	150,000	150,000	125,000	105,378	82,267	71,000	54,100	44,000	52,000	52,000
06 04	Other use of solvents and related activities											
06 04 01	Glass wool enduction	<i>g/Mg product</i>	800	800	800	800	800	800	800	800	800	800
06 04 02	Mineral wool enduction	<i>g/Mg product</i>	300	300	300	300	300	300	300	300	300	300
06 04 03	Printing industry	<i>g/Mg ink</i>	234,649	228,190	184,332	174,227	174,227	174,227	174,227	174,227	174,227	174,227
06 04 04	Fat, edible and non edible oil extraction	<i>g/Mg product</i>	1,570	1,570	1,570	1,570	1,570	1,570	1,570	1,570	1,570	1,570
06 04 05	Application of glues and adhesives	<i>g/Mg product</i>	600,000	151,230	108,086	84,190	84,190	84,190	84,190	84,190	84,190	84,190
06 04 06	Preservation of wood	<i>g/Mg product</i>	105,000	105,000	105,000	105,000	105,000	105,000	105,000	105,000	105,000	105,000
06 04 08	Domestic solvent use (other than paint application)(k)	<i>g/Mg product</i>	60,117	52,262	42,356	46,153	42,172	34,483	38,263	34,979	36,429	35,199
06 04 09	Vehicles dewaxing	<i>g/vehicles</i>	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000	1,000

5.4 QA/QC AND VERIFICATION

Data production and consumption time series for some activities (paint application in constructions and buildings, polyester processing, polyurethane processing, pharmaceutical products, paints manufacturing, glues manufacturing, textile finishing, leather tanning, fat edible and non-edible oil extraction, application of glues and adhesives) are checked with data acquired by the National Statistics Institute (ISTAT, several years (a), (b) and (c)), the Sectoral Association of the Italian Federation of the Chemical Industry (AVISA, several years) and the Food and Agriculture Organization of the United Nations (FAO, several years). For specific categories, emission factors and emissions are also shared with the relevant industrial associations; this is particularly the case of paint application for wood, some chemical processes and anaesthesia and aerosol cans.

In the framework of the MeditAIRaneo project, ISPRA commissioned to Techne Consulting S.r.l. a survey to collect national information on emission factors in the solvent sector. The results, published in the report "Rassegna dei fattori di emissione nazionali ed internazionali relativamente al settore solventi" (TECHNE, 2004), have been used to verify and validate emission estimates. In 2008, ISPRA commissioned to Techne Consulting S.r.l. another survey to compare emission factors with the last update figures published in the EMEP/CORINAIR guidebook (EMEP/CORINAIR, 2007). The results are reported in "Fattori di emissione per l'utilizzo di solventi" (TECHNE, 2008) and have been used to update emission factors for polyurethane and polystyrene foam processing activities.


In addition, for paint application, data communicated from the industries in the framework of the EU Directive 2004/42, implemented by the Italian Legislative Decree 161/2006, on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products have been used as a verification of emission estimates. These data refer to the composition of the total amount of paints and varnishes (water and solvent contents) in different subcategories for interior and exterior use and the total amount of products used for vehicle refinishing and they are available from the year 2007.

Verifications of the emissions from the sector occurred in 2012, on account of the bilateral independent review between Italy and Spain and the revision of national estimates and projections in the context of the National emission ceilings Directive for the EU Member States and the Gothenburg Protocol of the Convention on Long-Range Transboundary Air Pollution (CLRTAP). The analysis by category did not highlight the need of major methodological revisions of the sector; an additional source of emissions was added affecting only NMVOC emissions.

Furthermore, every five years (for inventory years from 1990 to 2015) and every four years for inventory years from 2015 onwards ISPRA carries out emission estimates at NUTS level which is the occasion of an additional check with local environmental agencies.

The distribution of NMVOC emissions from the solvent and other product use sector at NUTS3 level for 2019 is reported in Figure 5.3; methodologies are described in the relevant publication (ISPRA, 2022).

Figure 5.3 NMVOC emissions from solvent and other product use in 2019 (t)



5.5 RECALCULATIONS

Recalculations occurred for the update of EFs in paint application in construction and buildings for the years 2019-2021.

In chemical product processing and manufacturing, a recalculation occurred on account of a revision in activity data for polyester, from 2018, and polyurethane processing, from 2008, and a change in EFs for leather tanning from 2019 due to information available from the industrial association reports.

NMVOC emissions from Road paving with asphalt (2D3d) have been revised on account of the last review process (EEA, 2023).

Other recalculations concerned the update of activity data in domestic solvent use of cosmetics, for 2021.

5.6 PLANNED IMPROVEMENTS

Specific developments will regard the improvement of emission factors for some relevant categories. In particular, several improvements are planned with the aim to update the status of technologies in this sector where main challenges regard the availability of data collected from the industry. Main focus will be on metal degreasing.

6 AGRICULTURE (NFR SECTOR 3)

6.1 OVERVIEW OF THE SECTOR


The agriculture sector is responsible for the largest part of NH₃ emissions, and contributes also to PM₁₀, PM_{2.5}, BC, TSP, NO_x, NMVOC, CO, SO₂, heavy metals (As, Cr, Cu, Ni, Se, Zn, Pb, Cd, Hg), Dioxins, PAH and HCB emissions. Italy estimates agricultural emissions for manure management (3B), agricultural soils (3D) including the use of pesticides, and field burning of agricultural wastes (3F). NO_x emissions are reported as NO₂.

In 2022, key categories level was identified for NH₃ emissions (3B1a, 3B1b, 3B3, 3B4gii, 3Da1, 3Da2a and 3Da2c), for NMVOC emissions (3B1a and 3B1b) and for PM₁₀ emissions (3Dc). In 1990 similar figures were obtained except for NH₃ emissions 3B4gii and 3Da2c and PM₁₀ emissions 3B4gii which were not key categories and HCB emissions 3Df which was key category. For the trend analysis, key categories were related to NH₃ emissions (3B1a, 3B3, 3B4a, 3B4gii, 3Da1 and 3Da2c), NMVOC emissions (3B1a and 3B1b), PM₁₀ emissions (3Dc) and HCB (3Df).

In 2022, NH₃ emissions from the agriculture sector were 317.3 Gg (90.4% of national emissions) where 3B, 3D and 3F categories represent 51.8%, 38.5% and 0.1% of total national emissions. The trend of NH₃ from 1990 to 2022 shows a 36.5% decrease due to the reduction in the number of animals, the diffusion of best environmental practices in manure management in relation to housing, storage and land spreading systems, the decrease of cultivated surface/crop production and use of N-fertilisers.

A representation of the contribution by source of agriculture NH₃ emissions for 1990 and 2022 is shown in Figure 6.1.

Figure 6.1 Share of NH₃ emissions in the agriculture sector for 1990 and 2022.



Agricultural official statistics are mainly collected from the National Institute of Statistics, ISTAT. Most important activity data (number of animals, N-fertilizers, agricultural surface and production, milk production) are available on-line: <http://dati.istat.it/>. ISTAT has a major role in the comprehensive collection of data through structural (such as the Farm Structure Survey, FSS) and conjunctural surveys, and the general agricultural census¹. For consistency reasons the same agricultural official statistics are used for UNFCCC and UNECE/CLRTAP emission inventory. ISPRA participates to the Agriculture, Forestry, and Fishing Quality Panel, which has been established to monitor and improve national statistics. This is the opportunity to get in touch with experts from the Agriculture Service from ISTAT in charge for main agricultural surveys. In this way, data used for the inventory is continuously updated according to the latest information available. Agricultural statistics reported by ISTAT are also published in the European statistics database² (EUROSTAT). The verification of statistics is part of the QA/QC procedures; therefore,

¹ The last census was conducted in 2021 (which surveyed data from the 2019/2020 crop year and livestock raised as of December 1, 2020) and the first results are available at the link <https://www.istat.it/it/archivio/273753>. Micro-data were released in January 2024.

² <http://ec.europa.eu/eurostat/data/database>

as soon as outliers are identified ISTAT and category associations are contacted. In Table 6.1 the time series of animal categories is shown.

Table 6.1 Time series of animals

Year	Dairy cattle	Non-dairy cattle	Buffalo	Sheep	Goats	Horses	Mules/asses	Swine	Rabbits	Poultry	Fur animals
1990	2,641,755	5,110,397	94,500	8,739,253	1,258,962	287,847	83,853	8,406,521	14,893,771	173,341,562	325,121
1995	2,079,783	5,189,304	148,404	10,667,971	1,372,937	314,778	37,844	8,060,676	17,110,587	184,202,416	220,000
2000	2,065,000	4,988,000	192,000	11,089,000	1,375,000	280,000	33,000	8,307,000	17,873,993	176,722,211	230,000
2005	1,842,004	4,409,921	205,093	7,954,167	945,895	278,471	30,254	9,200,270	20,504,282	174,667,361	200,000
2010	1,746,140	4,086,317	365,086	7,900,016	982,918	373,324	46,475	9,321,119	17,957,421	175,912,339	125,000
2015	1,826,484	3,954,864	374,458	7,148,534	961,676	384,767	70,872	8,674,793	15,760,502	177,391,671	180,000
2019	1,643,117	4,331,830	402,286	7,000,880	1,058,720	367,561	72,455	8,510,268	11,755,922	175,520,313	135,000
2020	1,638,382	4,354,633	407,027	7,034,164	1,065,712	367,561	72,455	8,543,029	11,010,203	178,906,532	50,000
2021	1,609,948	4,260,926	409,408	6,728,351	1,060,748	367,561	72,455	8,407,968	10,945,940	178,243,638	20,000
2022	1,631,128	4,001,608	416,053	6,567,546	1,010,143	365,414	75,978	8,739,384	12,674,189	165,277,922	20,000

As for poultry, since no annual statistics on the number of animals are available, the following methodology was followed. For 1990 the ISTAT data from the Agricultural Census have been used; for the years 1991-1999, the number of heads was estimated on the basis of the annual decreases/increases in the production of heads and meat supplied by UNA (National Union of Poultry, which later became UNAITALIA); for 2000 and 2010 ISTAT data from the Censuses of Agriculture for laying hens and broilers were used; for the period 2001-2009 and since 2011 data on the number of broilers and laying hens have been updated, as described below; for the other poultry category, since 1998 the data have been estimated on the basis of UNAITALIA data. Data on turkeys derive from the ISTAT statistics on the Census and the FSS survey.

As stated, data on the number of broilers and laying hens in the period 2001-2009 and since 2011 have been updated. The estimation methodology involved successive steps. Firstly, ISTAT data from the Census and FSS surveys (available for the years 2000, 2005, 2007, 2010, 2013 and 2016) were taken into account; on the basis of these data the number of heads was estimated for the missing years from 2001, assuming a linear trend. The second step involved estimating the number of animals since 2001 on the basis of production data provided by UNAITALIA. The annual variation in production was multiplied by the number of animals in the 2000 Census. The third step was to calculate the average of the two time series calculated in the previous steps.

In Table 6.2 the nitrogen content of N-fertilisers by type applied to soils is shown together with the differentiated EFs. Detailed figures for "other nitrogenous fertilizers" are reported from 1998 because disaggregated official statistics from ISTAT were available only from that year.

Table 6.2 Time series of N content by fertilisers and relevant emission factors

Type of fertilizers	Emission factor		1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
	% of N applied	g NH ₃ /kg N applied	Nitrogen content (t N yr ⁻¹)									
Ammonium sulphate	12.55 %	152.42	50,762	61,059	36,698	27,855	32,568	16,986	10,267	17,830	22,963	14,261
Calcium cyanamide	3.51 %	42.60	3,310	507	3,003	2,357	4,958	3,046	2,655	2,845	2,845	2,628
Nitrate (*)	6.3 0%	76.45	46,657	52,769	48,701	58,427	32,964	40,157	22,920	28,097	23,786	15,543
CAN	3.51 %	42.60	112,565	139,253	112,541	109,445	71,261	51,200	43,514	55,339	57,134	31,797
Urea	16.66 %	202.31	291,581	321,196	329,496	317,814	209,829	266,154	247,199	295,134	240,692	135,156
Other nitric nitrogen	3.51 %	42.60	-	-	3,204	5,219	3,332	1,189	1,784	2,155	1,012	2,224
Other ammoniacal nitrogen	3.51 %	42.60	-	-	6,278	18,069	12,412	7,035	7,335	9,036	10,480	6,415
Other amidic nitrogenous	16.66 %	202.31	-	-	6,988	17,420	15,366	11,796	24,331	29,366	43,075	13,394
Phosphate nitrogen	12.55 %	152.42	112,237	99,468	77,916	69,758	45,837	35,054	34,648	44,533	43,410	19,983
Potassium nitrogen	4.28 %	52.00	3,937	2,876	5,291	12,289	15,955	9,077	7,922	11,353	11,169	7,181
NPK nitrogen	12.55 %	152.42	138,018	101,528	113,897	106,384	64,462	50,174	36,686	45,730	53,190	28,166
Organic mineral	3.51 %	42.60	444	20,960	38,688	34,809	19,085	25,986	27,581	36,033	29,137	23,040
Total			759,510	799,614	782,701	779,846	528,029	517,854	466,842	577,451	538,893	299,788

(*) includes ammonium nitrate < 27% and ammonium nitrate > 27% and calcium nitrate. The emission factor g NH₃/kg N applied is a weighted average of factors 42.60 and 152.42 g NH₃/kg N applied of ammonium nitrate and calcium nitrate.

6.2 METHODOLOGICAL ISSUES

Methodologies used for estimating national emissions from this sector are based on and conform to the EMEP/EEA Guidebook (EMEP/EEA, 2023) and the IPCC Guidelines (IPCC, 1997; IPCC, 2006). Consistency among methodologies for the preparation of the agricultural emission inventory under the UNFCCC and UNECE/CLRTAP is guaranteed through an operational synergy for activity data collection, inventory preparation and reporting to international conventions and European Directives. Information reported in the National Inventory Report/Common Reporting Format (NIR/CRF) for the GHG inventory is coherent and consistent with information reported in the Informative Inventory Report/Nomenclature for Reporting (IIR/NFR).

Factor 1.214 (= 17/14) was used to convert ammonia nitrogen (N-NH₃) to ammonia (NH₃) and factor 3.286 (= 46/14) was used to convert the nitrogen form (N-NO) to nitrogen dioxide (NO₂).

6.2.1 Manure management (3B)

For 3B category, Italy has estimated emissions for pollutants recommended in the 2023 EMEP/EEA Guidebook (NH₃, NO_x, NMVOC, PM₁₀ and PM_{2.5}). A detailed and updated description of the methodologies for the estimation of NH₃ emissions, as well as of national specific circumstances and reference material, is provided in sectoral reports (APAT, 2005; C3ndor et al., 2008; C3ndor, 2011), and in the NIR (ISPRA, several years [a]). Detailed information on activity data sources, methods and EFs by pollutant for 3B category is shown in Table 6.3.

Table 6.3 Activity data sources, methods and emission factors by pollutant for manure management.

NFR code	Animal category	Method	Activity data	Emission Factor
3B1a, 3B1b	Cattle	T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5)	NS	CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC)
3B4a, 3B2, 3B4d, 3B4e, 3B4f	Buffalo, Sheep, Goats, Horses, Mules and Asses	T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5)	NS, IS	CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC)
3B3	Swine	T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5)	NS	CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC)
3B4gi, 3B4gii, 3B4giii, 3B4giv	Poultry	T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5)	AS	CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC)
3B4h	Other	T2 (NH ₃ , NO _x , NMVOC), T1 (PM10, PM2.5)	NS	CS (NH ₃ , NO _x), D (PM10, PM2.5), T2 (NMVOC)

NS=national statistics; IS= International statistics (FAO); AS= category association statistics (UNAITALIA); CS=country-specific; D=Default (from EMEP/EEA Guidebook)

Concerning the 3B category, the estimation procedure for NH₃ emissions consists in successive subtractions from the quantification of nitrogen excreted annually for each livestock category. This quantity can be divided in two different fluxes, depending on whether animals are inside (housing, storage and manure application) or outside the stable (grazing). More in detail, the proportion of nitrogen excreted within the shelter undergoes losses through volatilization already during the course of the manure's stay within the breeding premises. These losses are calculated with the emission factors for emissions from housing for different livestock categories; this amount of nitrogen lost is therefore subtracted from the total nitrogen excreted to obtain the amount of nitrogen for storage. During storage another fraction of nitrogen is lost (calculated with the relevant emission factor for storage), which is then subtracted to obtain the amount of nitrogen available for the agronomic spreading. Losses occurring during the spreading are finally calculated with the specific emission factor for spreading. For the nitrogen excreted in the pasture losses due to volatilization calculated with the relevant emission factor for grazing by livestock only occur at this stage (CRPA, 2006[a]).

The manure application source is reported in 3Da2a Animal manure applied to soils and the animal grazing source is reported in 3Da3 Urine and dung deposited by grazing animals. As regards the animal grazing, the percentage of grazing animals is equal to (CRPA, 1997, CRPA, 2006[a]): 5% for dairy cattle, 2.2% for other cattle, 2.9% for buffalo, 60% for equines, 90% for sheep and goats. The excretion rates, slurry/solid manure production and average weights are derived from country specific information (CRPA, 2018; CRPA, 2006[a]; GU, 2006; Xiccato et al., 2005; Regione Emilia Romagna, 2004). Other improvements of country specific EFs were obtained with research studies (CRPA, 2010[b]; CRPA, 2006[a], [b]; CRPA, 2000). Average weight and N excretion rate for NH₃ estimations are reported in Table 6.4.

Table 6.4 Average weight and nitrogen excretion rates from livestock categories in 2022

Category	Weight	Housing	Grazing	Total
	kg	kg N head-1 yr-1		
Non-dairy cattle	393.6	50.29	1.39	51.67
Dairy cattle	602.7	106.22	5.59	111.81
Buffalo	493.9	57.20	1.71	58.91
Other swine (*)	88.1	13.45	-	13.45
Sow (*)	172.1	28.39	-	28.39
Sheep	48.2	1.62	14.58	16.20
Goats	49.1	1.62	14.58	16.20

Category	Weight kg	Housing	Grazing kg N head-1 yr-1	Total
Horses	550.0	20.00	30.00	50.00
Mules and asses	300.0	20.00	30.00	50.00
Poultry	1.7	0.48	-	0.48
Rabbit	1.6	1.02	-	1.02
Fur animals	1.0	4.10	-	4.10

(*) Other swine and sows are sources that represent the 'swine' category

6.2.1.1 Dairy cattle (3B1a)

Following the update of the gross energy intake (GE), based on the estimation of the parameter digestibility (DE) of diet, the excreted nitrogen value of dairy cows was updated for the whole time series. Excreted nitrogen is in fact calculated from GE using equations 10.31-10.33 of the 2006 IPCC Guidelines. In addition, the percentage for protein in diet has been updated from the previous submission. This parameter is used with GE in the estimation of excreted nitrogen.

As regards the DE parameter, the estimation methodology, set out in the 2019 IPCC Guidelines, allows, from data on average annual milk production per cow and per production level (low "<5000 kg/head/year", medium "5000-8500 kg/head/year" and high ">8500 kg/head/year") and information on animal diets, to calculate an average value of DE and Y_m for dairy cows. Based on data from the Italian Livestock Breeders' Association (AIA) on average annual milk production and the number of dairy cows in production, by region and breed, the distribution of animals was calculated according to the three productivity levels identified by the 2019 IPCC Guidelines, for the years 2004-2019. The AIA carries out milk productivity checks on behalf of the Ministry of Agriculture and each year the sample of animals checked is about 50% of the number of animals reared. The difference in dairy cow numbers between the AIA total and the ISTAT total (used for emission estimates) was attributed to the low production level. The DE values assigned to the three production levels (low, medium, high) are 62, 65 and 70.11 respectively and were identified in collaboration with the CRPA dairy cow feeding experts. The value 62 is the minimum value of the range indicated in Table 10.12 for low producing cows. The value 65 is lower than the average value of the range indicated in Table 10.12 for medium producing cows. The value 70.11 for high-producing cows is a weighted average of two values: the first is 65 (corresponding to diets with DE \geq 70 and NDF \geq 35) and was attributed to 27% of the high-producing cows fed without silage fodder; the second is 72 (corresponding to diets with DE \geq 70 and NDF \leq 35) and was attributed to 73% (= 100 - 27%) of the high-producing cows fed silage fodder. With reference to the 27% of cows, this value includes cows whose milk is intended for the production of Parmigiano Reggiano (17% of total cows), and cows fed with good quality dry and green fodder (e.g. for the production of Trentingrana PDO (Protected Designation of Origin), Latte Fieno STG (Traditional Speciality Guaranteed) and other mountain cheeses; 10% of total cows). In support of the choices made for high productivity values, mention is made of a study published in 2020 (Gislon et al, 2020) carried out on eight Italian Friesian cows in multiparous lactation, with high productivity, using a 4 × 4 replicated Latin square pattern. The experimental design of the square involves all cows receiving all diets (with adaptation periods between each), so we have 2 groups of 4 cows that rotated 4 times on as many diets. The number of observations for each diet is 32. The cow effect is nullified because they all receive all diets and the results obtained are therefore irrefutable and highly representative, according to CRPA experts. Four diets, based on the following forages (% of dry matter, DM; neutral detergent fiber content, NDF, expressed as % of DM), were tested: corn silage (CS, 49.3; 32.8 NDF), alfalfa silage (AS, 26.8; 27.1 NDF), wheat silage (WS, 20.0; 33.7 NDF), and a typical hay-based Parmigiano Reggiano cheese production diet (PR, 25.3 of both alfalfa and Italian ryegrass hay; 36.7 NDF). The lowest DM digestibility was observed for the PR diet (64.5%) and the highest for the CS diet (73.3%); AS and WS diets showed intermediate values (71.4 and 70.3% respectively). PR diet is associated with diets with DE \geq 70 and NDF \geq 35 in table 10.12 of the 2019 IPCC Guidelines and the other three diets are associated with diets with DE \geq 70 and NDF \leq 35 in the same table. For the year 2022, the percentages of

dairy cows according to the three productivity levels (high, medium and low) are 78.2%, 11.2% and 10.7%. The digestibility values associated with these productivity levels are, as mentioned earlier, 70.11%, 65% and 62% respectively. With these data, the average digestibility value of the diets consumed by dairy cows was calculated as 68.67%. As regards the percentage for protein in diet, mentioned above, on the basis of data from around 500 samples of rations (unifeed) of lactating and dry dairy cows, analysed by the CRPA's zootechnical feed service for the three-year period 2017-2019, from all over Italy, the crude protein of the ration was updated. The data were obtained by weighing the values expressed as % of the dry matter of the ration with the average annual lactation period (equal to 305 days) and the dry period (equal to 60 days). The value obtained, 14.22, was used for the time series from 2010 onwards, as indicated by the CRPA experts. For the previous years, the previous figure of 15.32 was left until 2000, and an average value of 14.5 was used for the intermediate years between 2000 and 2010.

In Table 6.5 the animal waste management system (AWMS) distribution and EFs used for the dairy cattle category are reported. For 2014-2022 for dairy cattle emissions were used data reported in Table 6.5 and specifically for housing data from 2010 and for storage, data from 2013. EF was multiplied by the percentage of the nitrogen excreted in housing equal to 95% of the total, assuming that 5% is excreted in grazing. The EF is a weighted average based on country specific emission factors for different livestock housing and the distribution of animals in shelters which has been assumed in the following main housing systems reported in Table 6.5 (based on a 1998 CRPA survey carried out in Lombardy, Emilia Romagna and the centre of Italy and on ISTAT statistics of 2003 and on 2010 Agricultural Census). Between 2005 and 2010 a gradual transition to the updated distribution of housing systems has been assumed for the intermediate years taking in account the gradual penetration of systems to ensure animal welfare.

Table 6.5 AWMS distribution and EF by manure management system for the dairy cattle category

Emission factors by manure management system	1990	2003	2005	2010	2013
Housing					
cubicle house: 14.3 N-NH ₃ kg/head/year (Bonazzi et al, 2005)	14.7%	14.7%	14.7%	27.9%	27.9%
loose housing on bedding: 15.7 N-NH ₃ kg/head/year (Bonazzi et al, 2005)	9.2%	9.2%	9.2%	42.6%	42.6%
tied cows: 12.9 N-NH ₃ kg/head/year (Bonazzi et al, 2005)	76.1%	76.1%	76.1%	29.5%	29.5%
EF N-NH ₃ kg/head/year	13.4	13.4	13.4	14.5	14.5
Storage					
liquid manure	liquid manure = 36%	liquid manure = 36%	liquid manure = 36%	liquid manure = 48%	liquid manure = 48%
Tanks (for liquid manure): 23% of N at storage (Bonazzi et al, 2005)	40.0%	75.5%	75.5%	82.3%	70.1%
Lagoons (for liquid manure): 32.2% (multiplication factor equal to 1.4 respect to tanks)	50.0%	12.5%	12.5%	2.5%	1.7%
covered storage (for liquid manure):	10.0%	12.5%	12.5%	15.2%	28.3%
covered tanks high reduction: 4.6% (reduction of 80% compared to tanks)		1.0%	1.0%	3.0%	4.0%
covered tanks medium reduction: 9.2% (reduction of 60% compared to tanks)		1.0%	1.0%	3.0%	4.0%
covered tanks low reduction: 13.8% (reduction of 40% compared to tanks)	10.0%	10.5%	10.5%	8.9%	16.4%
biogas: no emission				0.32%(1)	3.9%(1)
solid storage: 14.2% of N at storage (Regione Emilia Romagna, 2001)	solid manure = 64%	solid manure = 64%	solid manure = 64%	solid manure = 52%	solid manure = 52%

(1) Data were calculated from the results of the CRPA study on assessing the emission effects of livestock processing (CRPA, 2018). The study estimated that in 2020 19% of cattle manure would be sent to anaerobic digestion and 48% of digestate tanks would be covered. Based on these results, it was assumed that: in 2010, 1% of manure went to digesters and 11% were covered digestate tanks; in 2013 these values become 10% and 27% respectively. This trend was assumed on the basis of the exponential growth in the last ten years of anaerobic digesters.

As regards the manure storage (see Table 6.5), emission factors are expressed as a percentage of the nitrogen contained in manure to storage. Emission factors used for tanks is derived from national literature (Bonazzi et al, 2005) and emission factors for lagoons and covered storage have been estimated applying an increase (for lagoons) and a reduction (for covered storage) to tanks EF (as referenced in CRPA, 2006[a] and CRPA, 2006[b]).

The proportion of liquid system (considering liquid system= liquid system + digesters) and solid storage (considering solid storage= solid storage + digesters), reported in the CRF (Common Reporting Format for the GHG inventory) refer to the nitrogen excreted and not to the amount of animal waste. The proportion reported in the Table 6.5 refer to the manure production according to the type of housing.

EFs for lagoons and covered storage have been provided by CRPA (CRPA, 2006[a]). For lagoons, they have a high exposure area relative to their capacity and represent a higher emission type than the tank. Considering the volumes of the two types of storage, an increase in the surface of slurry in the lagoons with respect to the tanks can be estimated equal to 40%. Since ammonia emissions are estimated to be proportional to the surface of slurry exposed to air, emissions from lagoons will be approximately 40% higher than those of the tanks (CRPA, 1997). For covered storage, the emission reduction has been assumed on the basis of the ILF-BREF document (EC, 2003) and the Ammonia Guidance Document (Bittman S. et al, 2014), as reported in CRPA (CRPA, 2006[a]; 2018).

A linear emission reduction in the period 1990-2003 has been estimated to assess the dynamics of evolution of storage systems from the values available in 1990 and 2003, as reported by CRPA (CRPA, 2006[a]). In 2003 respect to 1990 an increase of storage in tanks with respect to lagoons as well as a small increase of covered storage is observed as available in the Table 6.5. On the basis of ISTAT statistics on storage systems as 2010 Agricultural Census and 2013 Farm Structure Survey, an update of emission factors from manure storage for cattle category has been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010 and 2010-2013) taking in account the gradual penetration of the abatement technologies.

On the basis of the study for the evaluation of the effects on emissions of livestock management practices carried out by CRPA for the emission scenarios for 2020 and 2030 (CRPA, 2018), NH₃ emissions from storage for cattle have been modified considering the average distribution of the covered tanks related to the different ammonia emission reduction efficiencies.

EFs for manure storage reported in the Table 6.5 have been multiplied by the percentage of nitrogen remaining after housing emissions and the result has been multiplied by the nitrogen excreted in housing to obtain emissions from storage. Emissions have been divided by total heads to obtain the EF kg/head reported in the Table 6.8 for the year 2022.

Regarding emission factors for cattle, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios in 2003 and an ad hoc survey conduct in the 2005 by CRPA (CRPA, 2006 [a], [b]) and on ISTAT statistics such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey.

6.2.1.2 *Swine (3B3)*

Activity data of swine population (3B3) reported in the IIR/NFR are different from data reported in the NIR/CRF. In fact, piglets (swine less than 20 kg) are included in the swine population in the NIR/CRF for the estimation of CH₄ emission from enteric fermentation, while they are not included in the number of the NFR templates because the NH₃ EF used for sows takes into account the emissions from piglets, thus ensuring the comparability of the implied emission factors. For NH₃ estimations average weighted emission factors for each category (other swine and sows) are calculated taking in account the relevant emission factors of the abatement technologies for each manure system.

The implemented abatement technologies for the years 1990, 2003 and 2005 are reported in Table 6.6. For 2014-2022 for swine emissions were used data reported in Table 6.6 and specifically for housing, data from 2005, for storage, data from 2013 and for spreading 2014-2015, data from 2013 and, for 2016 onwards, data from 2016.

Table 6.6 Abatement technologies for the swine category

Livestock category	1990	2003	2005	2010	2013	
Housing						
fattening swine	55% Partly-slatted floor (PSF);20% Fully-slatted floor (FSF);25% solid floor	55% PSF;25% FSF;20% solid floor	26% FSF;39% PSF;12% FSF + vacuum system (VS);4% FSF + with flush canals;7% FSF + with flush tubes;5% PSF + VS;6% PSF + with flush canals;1% PSF + with flush tubes	Same distribution for the year 2005	Same distribution for the year 2005	
gestating sows (75% of the total sows)	65% FSF; 35% PSF	50% FSF; 50% PSF	26% FSF; 52% PSF; 5% FSF + vacuum system (VS); 5% FSF + with flush canals;7% FSF + with flush tubes; 2% PSF + VS; 2% PSF + with flush canals; 1% PSF + with flush tubes	Same distribution for the year 2005	Same distribution for the year 2005	
lactating sows (25% of the total sows)	75% FSF+ deep collection pit; 25% sloping floor	65% FSF+ deep collection pit; 35% sloping floor	52% FSF + deep collection pit; 39% sloping floor; 3% with flush; 6% mechanical removal	Same distribution for the year 2005	Same distribution for the year 2005	
weaners 6-20 kg	80% FSF + deep collection pit; 20% sloping floor	70% FSF+ deep collection pit; 30% sloping floor	63% FSF + deep collection pit; 14% sloping floor; 7% FSF + VS; 11% FSF with flush tubes; 2% FSF + scraper; 2% PSF + VS; 1% PSF + deep collection pit	Same distribution for the year 2005	Same distribution for the year 2005	
	1990	2003	2005	2010	2013	
Storage						
swine	61% lagoons;36% tanks;3% covered storage:	54% lagoons; 43% tanks; 3% covered storage:	46% lagoons; 51% tanks; 3% covered storage:	10% lagoons; 79% tanks; 11% covered storage:	7% lagoons; 67% tanks; 25% covered storage:	
covered tanks high reduction: reduction of 80% compared to tanks				1%	3%	
covered tanks medium reduction: reduction of 60% compared to tanks		1%	1%	4%	5%	
covered tanks low reduction: reduction of 40% compared to tanks	3%	2%	2%	6%	17%	
biogas: no emission				0.32%(1)	0.80%(1)	
	1990	2003	2005	2010	2013	
Land spreading						
swine	100%	80% broadcasting	78% broadcasting	70% broadcasting	48% broadcasting	36% broadcasting
		10% low efficiency	11% low efficiency	17% low efficiency	30% low efficiency	27% low efficiency
				6% medium efficiency	12% medium efficiency	26% medium efficiency
		10% high efficiency	11% high efficiency	7% high efficiency	11% high efficiency	11% high efficiency

(1) Data were calculated from the results of the CRPA study on assessing the emission effects of livestock processing (CRPA, 2018). The study estimated that in 2020 3% of swine manure would be sent to anaerobic digestion and 48% of digestate tanks would be covered. Based on these results, it was assumed that: in 2010 and 2013 the percentage of manure sent to digesters remains at 3%, while the percentage of covered digestate tanks changes, becoming 11% and 27% respectively.

Regarding emission factors for swine, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios in 2003 and an ad hoc survey conducted in the 2005 by CRPA (CRPA, 2006 [a], [b]). Furthermore, an update of emission factors from manure storage and land spreading for swine category has been estimated on the basis of ISTAT statistics on manure storage systems and

land spreading techniques such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010, 2010-2013 and 2014-2016) taking in account the gradual penetration of the abatement technologies.

On the basis of the study for the evaluation of the effects on emissions of livestock management practices carried out by CRPA for the emission scenarios for 2020 and 2030 (CRPA, 2018), NH₃ emissions from storage for swine have been modified considering the average distribution of the covered tanks related to the different ammonia emission reduction efficiencies.

6.2.1.3 Poultry (3B4g)

As regards 3B4gi (laying hens) and 3B4gii (Broilers) categories, NH₃ emissions show different trends. The different trend for the laying hens is due to the evolution of different abatement technologies along the period, that are considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Emission factors used for each of the different abatement technologies for laying hens (as referenced in CRPA, 2006[a] and CRPA, 2006[b]) are reported in Table 6.7.

Table 6.7 AWMS distribution, abatement technologies and EF by manure management system for the laying hens

Emission factors by manure management system	1990	2003	2005	2010
Housing				
open manure storage under cages (for liquid manure) (RS) = 0.220 kg NH ₃ /head/year (EC, 2003)	100%	20%	11%	4%
deep pit = 0.162 kg NH ₃ /head/year (ENEA, 2003)		24%		
vertical tiered cages with manure belts and forced air drying = 0.06 kg NH ₃ /head/year (ENEA, 2003) [reduction in ammonia emissions of 73% compared to RS]		56%	74%	50%
vertical tiered cages with manure belt and whisk-forced air drying = 0.088 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 60% compared to RS]			2%	
aerated open manure storage (deep-pit or high-risesystems and canal house) = 0.154 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 30% compared to RS]			10%	11%
vertical tiered cages with manure belt and drying tunnel over the cages = 0.044 kg NH ₃ /head/year (EC, 2003) [reduction in ammonia emissions of 80% compared to RS]			3%	
Loose housing with outdoor access (RS) = 0.3 kg NH ₃ /head/year (Bittman S. et al, 2014)				7%
Loose housing without outdoor access = 0.18 kg NH ₃ /head/year (Bittman S. et al, 2014; our assumptions)				28%
Storage				
liquid manure = 16% (percentage of nitrogen to storage) (Nicholson et al, 2004)	100%	20%	11%	4%
solid manure = 7.3% (ENEA, 2003)		80%	89%	96%
Land spreading				
liquid manure = 37.1% of TAN applied (TAN/TKN = 35%) (CRPA, 2006[a]) [broadcasting]	100%	5%	9%	9%
low efficiency = 7.8% (bandspreading and incorporation within 6 hours for liquid manure) [reduction of 40% compared to broadcasting]		50%	65%	65%
high efficiency = 2.6% (shallow and deep injection for liquid manure) [reduction of 80% compared to broadcasting]		45%	26%	26%
solid manure = 67% of TAN applied (TAN/TKN = 21%) (Nicholson et al, 2004; CRPA, 2006[a]) [broadcasting]		10%	9%	10%
low efficiency = 11.0% (incorporation within 12-24 hours for solid manure) [reduction of 20% compared to broadcasting]		40%	37%	43%
high efficiency = 2.8% (incorporation within 4 hours for solid manure) [reduction of 80% compared to broadcasting]		50%	54%	46%

Emission factors used for each of the different techniques for housing are derived from ILF BREF of IPPC (EC, 2003) and a study at national level on ammonia emissions from laying hens (ENEA, 2003). In 2010, on the basis of the housing distribution collected from the 2010 Agricultural Census and emission factors

and abatement systems data reported in the Guidance from the UNECE Task Force on Reactive Nitrogen (Bittman S. et al, 2014) emission factors have been updated. Between 2005 and 2010 a gradual transition to the updated distribution of housing systems has been assumed for the intermediate years taking in account the gradual penetration of systems to ensure animal welfare. From 1995 a chicken-dung drying process system has been introduced for laying hens and improved along the period.

As regards the manure storage, emission factors are expressed as a percentage of the nitrogen contained in manure to storage. Emission factors used for liquid manure is derived from Nicholson et al (Nicholson et al, 2004) and emission factors for solid manure is from ENEA (ENEA, 2003). On the basis of the 2010 Agricultural Census conducted by ISTAT, an update of emission factors from manure storage for laying hens category has been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010) taking in account the gradual penetration of the abatement technologies. EFs for manure storage reported in Table 6.7 have been multiplied by the amount of nitrogen remaining after housing emissions.

For land spreading, emissions have been estimated by CRPA (CRPA, 2006[a] and CRPA, 2006[b]). As regards the liquid manure, the amount of N-NH₄ emissions, in percentage of the applied ammoniacal nitrogen, have been assumed equal to those of the cattle slurry due to the lack of data (CRPA, 2006[a]). As regards the solid manure, the amount of N-NH₄ emissions, in percentage of the applied ammoniacal nitrogen, were equal to 67% (Nicholson et al, 2004; CRPA, 2006[a]). In 2003 and 2005 the evolution of different improvements technologies based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios and a survey conduct by CRPA, has been implemented in the EFs used. For the period 1900-2003, a linear emission reduction has been estimated and applied. The efficiency of reduction techniques has been estimated on the basis of the UNECE document Control techniques for preventing and abating emissions of ammonia (as referenced in CRPA, 2006[a] and CRPA, 2006[b]). EFs for land spreading reported in Table 6.7 have been multiplied by the amount of nitrogen remaining after storage emissions.

As regards broilers, only a slight improvement on spreading system has occurred.

As recommends by the 2019 and 2020 NECD review (EEA, 2019; EEA, 2020), emissions of NO_x, NH₃, PM and NMVOC from turkeys have been estimated and reported in category 3B4giii.

Regarding emission factors for poultry, the evolution of different abatement technologies along the period is considered in the EFs used for NH₃ estimation for housing, storage and land spreading systems. Improvements in the abatement technologies are based on the results of both the IIASA questionnaire for the implementation of RAINS scenarios in 2003 and an ad hoc survey conduct in the 2005 by CRPA (CRPA, 2006 [a], [b]) and on ISTAT statistics such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey.

Regarding equines, the ISTAT statistics used for estimation include equines intended and not intended for food production. From 2018 to 2021, ISTAT did not publish data on stocks, and to 2022, the ISTAT data are those from the National Livestock Registry Database (BDN), which includes all equines bred.

Average emission factors for NH₃ per head are reported in Table 6.8

Table 6.8 NH₃ emission factors for manure management for the year 2022

Category	Housing	Storage	Land spreading kg NH ₃ head-1 yr-1	Grazing	Total
Non-dairy cattle	7.92	6.38	6.04	0.13	20.48
Dairy cattle	16.73	13.23	1	0.54	42.48
Buffalo	9.01	9.42	7.86	0.17	26.45
Other swine (*)	2.38	1.63	1.12		5.13
Sow (*)	4.86	3.46	2.37		10.70

Category	Housing	Storage	Land spreading kg NH ₃ head-1 yr-1	Grazing	Total
Sheep	0.22	0.28	0.32	0.71	1.53
Goats	0.22	0.25	0.33	0.71	1.50
Horses	3.21	4.43	1.97	2.91	12.51
Mules and asses	3.21	4.43	1.97	2.91	12.51
Laying hens	0.13	0.04	0.04		0.21
Broilers	0.08	0.04	0.03		0.14
Turkeys	0.25	0.15	0.06		0.46
Other poultry	0.09	0.02	0.02		0.13
Rabbit	0.34	0.13	0.06		0.54
Fur animals	1.37		0.30		1.67

(*) Other swine and sows are sources that represent the 'swine' category

NH₃ emissions from digesters biogas facilities (in particular due to different phases of the process: during storage of feedstock on the premises of the biogas facility, during the liquid–solid separation of the digestate, during storage of the digestate) have been estimated on the basis of the amount of nitrogen in manure feeding anaerobic digesters and the tier 1 emission factor derived by the EMEP/EEA Guidebook (EMEP/EEA, 2023). On the basis of CRPA data on measurements of nitrogen quantities in livestock manure (downstream of releases to housing and storage) per animal category and type of manure, the nitrogen quantities in livestock manure sent to anaerobic digestion were estimated. The coefficients, expressed as g N/kg manure, were calculated gross of losses and then the losses to housing were deducted. The resulting coefficients were then multiplied by the quantities of manure sent for anaerobic digestion. The whole time series was updated. NH₃ emissions from digesters biogas facilities have been subtracted from manure management category (for cattle, swine and poultry categories) and allocated in the anaerobic digestion at biogas facilities (5B2 of the waste sector). As requested during the 2019 and 2020 NECD reviews (EEA, 2019; EEA, 2020), the amount of total feedstock, livestock manure and nitrogen in manure treated by biogas facilities are shown in Table 6.9.

Table 6.9 Total feedstock, animal manure and nitrogen in manure treated by biogas facilities.

Year	Total feedstock (t)	Animal manure in total feedstock (t)	Nitrogen in animal manure (kg)
1990	-	-	-
1995	n.a.	273,863	1,173,358
2000	n.a.	165,670	698,951
2005	n.a.	1,078,548	4,399,024
2010	6,513,271	1,766,348	7,023,590
2015	29,551,431	13,600,442	71,578,207
2019	33,263,320	15,266,757	88,788,793
2020	34,301,943	15,763,715	94,496,002
2021	35,754,088	16,405,322	98,516,252
2022	35,868,469	16,445,677	96,603,707

n.a. = not available

Because of multiple substrates fed to bio-digesters, the following average characteristics of the feedstock, reported in Table 6.10, as supplied by CRPA, are considered for the Italian bio-digesters to calculate the total amount of feed from animal manure anaerobic digestion (CRPA, 2018).

Table 6.10 Percentages of different substrates for anaerobic digestion feedstock

Type of feed	Units	animal manure	energy crops	agro-industrial by-products
Animal manure only	% in the feed	100	0	0
Animal manure + energy crops + agro-industrial by-products	% in the feed	28	52	20
Animal manure + energy crops	% in the feed	38	62	0
Animal manure + agro-industrial by-products	% in the feed	69	0	31
Energy crops + agro-industrial by-products	% in the feed	0	81	19

Source: CRPA

On the basis of the information reported above and in consideration of the typical feed of the bio-digesters the average parameters for animal manure, energy crops and agro-industrial by-products are those reported in Table 6.11. The biogas methane content is generally reported to range from 50% to 65%, for the inventory purposes and according to CRPA methane content is assumed to be 55%. As regards the average volatile solids content, values for animal manure and agro-industrial by-products have been changed based on the recent study of CRPA (CRPA, 2018).

Table 6.11 Average parameters of different substrates for anaerobic digestion feedstock

Parameters	Units	animal manure	energy crops	agro-industrial by-products
Average biogas producing potential	m ³ biogas/kg VS	0.4	0.6	0.6
Average CH ₄ content	%	55	55	55
Average volatile solids content	kg/t feed	139	280	237

Source: CRPA

For further information on the method of estimating the quantity of manure sent to digesters and the amount of nitrogen stored in digesters, see the information and data reported in the NIR (see paragraphs 5.3.2 Methodological issues in chapter 5 and A7.2 Manure management (3B) in annex 7).

The percentage of nitrogen lost through N-NH₃ emissions from anaerobic digesters was subtracted from the percentage of nitrogen left after emissions during housing and storage, reducing the amount of nitrogen used at the spreading. The amount of nitrogen used at the spreading also includes the digestate.

For NO_x emissions (during storage) tier 2 method reported in the EMEP/EEA Guidebook (EMEP/EEA, 2023) was used for calculations. EFs by livestock category and manure type derived from the EMEP/EEA Guidebook (EMEP/EEA, 2023) are based on nitrogen mass-flow approach built from country specific data on nitrogen excretion and solid/liquid distribution of manure.

For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2023). As requested during the 2020 NECD reviews (EEA, 2020), NMVOC emissions from turkeys are estimated and reported in the 3B4giii category.

In the Table 6.12 parameters used to estimate NMVOC emissions of categories 3B1a and 3B1b such as feed intake in MJ, silage fraction, proportion of housing, emission of NH₃ from housing, storage and spreading for 2022 are reported, as requested during the 2023 NECD reviews.

Table 6.12 Parameters of estimate NMVOC emissions of categories 3B1a and 3B1b for 2022

Parameters	Dairy cattle	Non-dairy cattle
GE MJ/capo/year	129,766.13	55,599.36
xhouse %	0.95	0.98
Fracsilage %	0.50	0.40
Fracsilage_store %	0.25	0.25
EF tier 2 silage_feeding kg NMVOC /MJ feed intake	0.0002002	0.0002002
EF tier 2 building kg NMVOC /MJ feed intake	0.0000353	0.0000353
EF tier 2 grazing kg NMVOC /MJ feed intake	0.0000069	0.0000069
ENMVOC, silage_store kg NMVOC/head	3.09	1.09
ENMVOC, silage_feeding kg NMVOC/head	12.34	4.35
ENMVOC, house kg NMVOC/head	4.35	1.92
ENH ₃ , building tons	22,468.79	26,096.49
ENH ₃ , storage tons	17,776.80	21,017.29
ENH ₃ , application tons	16,084.72	19,918.87
ENMVOC, manure_store kg NMVOC/head	3.44	1.55
ENMVOC, MM g NMVOC/head	23,219.82	8,906.60
Head number	1,631,128	4,001,608
ENMVOC, MM kt NMVOC	37.87	35.64

For particulate matter emissions a tier 1 method was used for calculations. EFs for PM₁₀ and PM_{2.5} are derived from the EMEP/EEA Guidebook (EMEP/EEA, 2023; EMEP/CORINAIR, 2006), modified based on the Italian animal breeding characteristics and weight parameters. For swine and poultry, up to 2004 the emission factors from the 2006 EMEP/CORINAIR Guidebook were used and from 2010 PM emission estimates are based on emission factors provided by the 2023 EMEP/EEA Guidebook. From 2005 to 2009 a gradual shift in emission factors was estimated, reflecting a gradual change in manure management systems, based on the studies underlying the emission factors in the two versions of the Guidebook and on data recorded by ISTAT surveys (FSS and the agricultural census). The 2023 EMEP/EEA emission factors are based on studies conducted between 2006 and 2016 which include scientific works conducted in Italy. These studies have suggested that Takai's emission factors suggested in the 2006 EMEP/CORINAIR Guidebook are too high and do not represent current particulate emission levels.

As requested during the 2020 NECD reviews (EEA, 2020), PM emissions from turkeys are estimated and reported in the 3B4giii category.

PM emissions from turkeys, sheep, goats, mules and asses and fur animals are also estimated. PM emission factors for sheep, goats, mules and asses and fur animals are from EMEP/EEA Guidebook 2023, Table A1.7 EFs for inhalable dust, respirable dust, PM₁₀ and PM_{2.5}. For turkeys were used the factors in

Table 3.5 Default Tier 1 estimates of EF for particle emissions from livestock husbandry (housing). The emission factors reported in Table 6.13 are the result of the product of the Guidebook emission factors and the ratio between the Guidebook average weights (reported in Table A1.5 Conventional livestock units and weights of livestock on which the N excretion estimates in Table 3.9 were based) and those of the national inventory by animal category. Average emission factors for PM per head are reported in Table 6.13.

Table 6.13 PM emission factors for manure management for the year 2022

Category	PM10	PM2.5
	kg PM head-1 yr-1	
Non-dairy cattle	0.324	0.214
Dairy cattle	0.657	0.428
Buffalo	0.499	0.326
Other swine (*)	0.190	0.008
Sow (*)	0.227	0.012
Sheep	0.054	0.016
Goats	0.055	0.017
Horses	0.242	0.154
Mules and asses	0.137	0.086
Laying hens	0.033	0.002
Broilers	0.024	0.002
Turkeys	0.109	0.020
Other poultry	0.051	0.008
Rabbit	-	-
Fur animals	0.003	0.002

(*) Other swine and sows are sources that represent the 'swine' category

6.3 AGRICULTURAL SOILS (3D)

For agricultural soils, estimations of NH₃ emissions account for the direct application of synthetic N-fertilizers (3Da1), animal manure applied to soils (3Da2a), sewage sludge applied to soils (3Da2b), other organic fertilisers applied to soil (3Da2c), animal grazing (3Da3), crop residues applied to soils (3Da4) and N fixed by leguminous crops (3De). For the same sources, emissions of NO_x were estimated (except for 3De cultivated crops and 3Da4 crop residues applied to soils). Indirect emissions from managed soils (3Db) and off-farm storage, handling and transport of bulk agricultural products (3Dd) have not been estimated as in the guidelines there is insufficient information. PM10 and PM2.5 emissions from the Farm-level agricultural operations including storage, handling and transport of agricultural products have been estimated and reported in 3Dc category. NMVOC emissions from animal manure applied to soils, animal grazing and cultivated crops have been estimated and reported in 3Da2a, 3Da3 and 3De categories respectively. HCB emissions from the use of pesticides have been estimated and reported in 3Df category.

NH₃ emissions from synthetic N-fertilizer (3Da1) are based on the guidebook methodology (EMEP/EEA, 2023), which provides different EFs by type of fertilizers and pH of the soil (EFs in Table 6.2). A tier 2

method has been implemented for 3Da1 source. NH₃ emissions from synthetic N-fertilizers are obtained with the amount of the N content by type of fertilizer multiplied by the specific EFs. NH₃ EFs from the use of synthetic fertilizers for normal and high pH factors (reported in the EMEP/EEA Guidebook (EMEP/EEA, 2023)) have been updated. To calculate a weighted average of emission factors distinguished according to soil pH type, CREA-AA³ processed the national agricultural areas distinguished according to eight soil pH classes, finally calculating the area percentages for normal and high pH (Rivieccio R., 2019).

Based on the comparison with ASSOFERTILIZZANTI and ISTAT experts on the time series of synthetic fertiliser use, the nitrate data in the years 2009-2011 were revised for the current submission. In addition, nitrate data (quantity and nitrogen content) were recalculated to include the estimated CAN fertiliser. The emission factor of NH₃ for the use of CAN from the EMEP/EEA Guidebook (EMEP/EEA, 2023) has been used. NO_x emission factor for synthetic N-fertilizer is equal to 0.04 kg NO₂/kg fertiliser N applied (EMEP/CORINAIR, 2023).

The method for estimating NH₃ emissions from animal manure applied to soils (3Da2a) is described in 3B (tier 2). On the basis of ISTAT statistics on spreading systems such as 2010 Agricultural Census, 2013 and 2016 Farm Structure Survey, an update of emission factors from land spreading for cattle, swine, laying hens and broilers categories have been estimated. A gradual transition to the updated emission factors has been assumed for the intermediate years (for the period 2005-2010, 2010-2013 and 2013-2016) taking in account the gradual penetration of the abatement technologies. For NO_x emissions (during spreading) a tier 2 method was used for calculations. EFs by livestock category and manure type derived from the EMEP/EEA Guidebook (EMEP/EEA, 2023) are based on nitrogen mass-flow approach. For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2023).

Concerning the sludge spreading (3Da2b), the total production of sludge from urban wastewater plants, as well as the total amount of sludge used in agriculture and some parameters such as N content, are communicated from 1995 by the Ministry for the Environment, Land and Sea from 1995 (MATTM, several years[a]) in the framework of the reporting commitments fixed by the European Sewage Sludge Directive (EC, 1986) transposed into the national Legislative Decree 27 January 1992, n. 99. From 1990 to 1994 activity data and parameters were reconstructed, as reported in detail in the Chapter 7 of the National Inventory Report on the Italian greenhouse gas inventory (ISPRA, several years [a]).

The amount of sewage N applied was calculated using the amount of sewage sludge (expressed in tons of dry matter) and the N content of sludge. The dry matter contained in sludge at national level is assumed to be 25% of total sludge. In Table 6.14, the total amount of sewage sludge production as well as sludge used in agriculture and nitrogen content in sludge is reported. The default NH₃ EF (0.13 kg NH₃/kg N applied) and NO_x EF (0.04 kg NO₂/kg N applied) are from EMEP/EEA Guidebook (EMEP/EEA, 2023).

Table 6.14 Sludge spreading activity data and parameters, 1990 – 2022.

Year	Sewage sludge production (t)	Sewage sludge used in agriculture (t)	Sewage sludge used in agriculture (t of dry matter)	N concentration in sludge (% dry matter)	Total N in sludge (t)
1990	3,272,148	392,658	98,164	5.2	5,071
1995	2,437,024	630,046	157,512	5.2	8,137
2000	3,402,017	869,696	217,424	5.0	10,954
2005	4,298,576	862,970	215,742	4.1	8,874
2010	3,697,625	992,859	248,215	4.0	10,040
2015	3,069,302	888,899	222,225	3.7	8,303

³ Council for Agricultural Research and Analysis of Agricultural Economics - Agriculture and Environment

Year	Sewage sludge production (t)	Sewage sludge used in agriculture (t)	Sewage sludge used in agriculture (t of dry matter)	N concentration in sludge (% dry matter)	Total N in sludge (t)
2019	2,954,886	685,371	171,343	4.0	6,791
2020	2,918,703	721,659	180,415	3.9	7,078
2021	2,929,994	686,838	171,710	3.9	6,745
2022	2,974,585	716,735	179,184	3.9	6,950

As regards the other organic fertilisers applied to soil (3Da2c) category, the use of other organic N fertilisers, including compost and organic amendments, and N content are provided by ISTAT (as reported in the paragraph 6.1). The default NH_3 EF (0.08 kg NH_3 /kg waste N applied) and NO_x EF (0.04 kg NO_2 /kg N waste applied) are from EMEP/EEA Guidebook (EMEP/EEA, 2023). For 3Da3 the time series of the quantity of N from animal grazing is the same as that reported in the NIR and in the relevant CRF tables. The method for estimating NH_3 emissions is described in 3B (tier 2). The default NO_x EF is from EMEP/EEA Guidebook (EMEP/EEA, 2023). For NMVOC emissions a tier 2 method was used for calculations. Tier 2 NMVOC EFs are those reported in the EMEP/EEA Guidebook (EMEP/EEA, 2023).

Nitrogen input from N-fixing crops (3De) has been estimated starting from data on surface and production for N-fixing crops and forage legumes; nitrogen input from N-fixing crops (kg N yr⁻¹) is calculated with a country-specific methodology. Peculiarities that are present in Italy were considered: N-fixing crops and legumes forage. Nitrogen input is calculated with two parameters: cultivated surface and nitrogen fixed per hectare (Erdamn 1959 in Giardini, 1983). Emissions are calculated using the default emission factor 1 kg N- NH_3 /ha (EMEP/CORINAIR, 2006). In Table 6.15, cultivated surface from N-fixing species (ha yr⁻¹) and N fixed by each species (kg N ha⁻¹ yr⁻¹) are shown.

Table 6.15 Cultivated surface (ha) and nitrogen fixed by each variety (kg N ha⁻¹ yr⁻¹)

N fixed kg N ha ⁻¹ yr ⁻¹	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022	
Bean, f.s.	40	29,096	23,943	23,448	23,146	19,027	17,059	18,253	17,915	18,378	15,748
Bean, d.s.	40	23,002	14,462	11,046	8,755	7,001	5,870	5,587	5,541	5,265	3,489
Broad bean, f.s.	40	16,564	14,180	11,998	9,484	8,487	7,914	7,624	7,372	7,310	14,424
Broad bean, d.s.	40	104,045	63,257	47,841	48,507	52,108	42,157	60,007	61,982	57,207	50,058
Pea, f.s.	50	28,192	21,582	11,403	11,636	8,691	14,940	16,197	16,154	15,730	16,428
Pea, d.s.	72	10,127	6,625	4,498	11,134	11,692	11,181	22,926	20,766	17,771	16,033
Chickpea	40	4,624	3,023	3,996	5,256	6,813	11,167	20,999	18,579	17,617	14,068
Lentil	40	1,048	1,038	1,016	1,786	2,458	3,099	5,861	5,612	5,710	4,925
Vetch	80	5,768	6,532	6,800	7,142	7,560	7,795	7,827	7,827	7,827	7,827
Lupin	40	3,303	3,070	3,300	2,500	3,401	3,358	3,337	3,337	3,337	3,337
Soya bean	58	521,169	195,191	256,647	152,331	159,511	308,979	273,332	256,134	285,464	342,532
Alfalfa	194	987,000	823,834	810,866	779,430	745,128	667,325	719,073	715,642	694,481	684,187
Clover grass	103	224,087	125,009	114,844	103,677	102,691	119,942	127,087	127,270	124,308	143,941
Total		1,958,025	1,301,746	1,307,702	1,164,784	1,134,567	1,220,786	1,288,110	1,264,131	1,260,404	1,316,997

s.=fresh seed; d.s.=dry seed

NM VOC emissions from cultivated crops have been estimated and reported in 3De category. The method (tier 1) for estimating NM VOC emissions from cultivated crops (3De) is described in 3D chapter of the EMEP/EEA Guidebook (EMEP/EEA, 2023). The default NM VOC EF is from EMEP/EEA Guidebook (EMEP/EEA, 2023). Hectares of wheat, rape, rye crops surface and total grass surface were considered as activity data according to the methodology EMEP/EEA Guidebook (EMEP/EEA, 2023).

NH₃ emissions from crop residues applied to soils have been estimated and reported in 3Da4 category. The method (tier 2) for estimating NH₃ emissions from crop residues (3Da4) is described in 3D chapter of the EMEP/EEA Guidebook (EMEP/EEA, 2023). Agricultural areas (cereals, legumes, tubers, vegetables, industrial crops, temporary and permanent fodder crops), above-ground residues, nitrogen content, and removed, burned and incorporated fractions of residues are country-specific data already used in estimating N₂O emissions from crop residues returned to soils. Regarding the incorporated fraction, incorporation is assumed to occur after 3 days, except in areas where green manure practice occurs. Green manure practice involves chopping cover crops and then incorporating them into the soil. The two operations are accomplished within 3 days. Therefore, for these residues, according to the 2023 EMEP/EEA Guidebook, there is no NH₃ emission. CREA-AA provided data on green manure areas by crop for the years 2019-2021, extracted from the FADN database (CREA, 2023).

PM₁₀ and PM_{2.5} emissions from the Farm-level agricultural operations including storage, handling and transport of agricultural products have been estimated and reported in 3Dc category. The method (tier 1) for estimating PM₁₀ and PM_{2.5} emissions is described in 3D chapter of the EMEP/EEA Guidebook (EMEP/EEA, 2023). The default PM₁₀ and PM_{2.5} EFs are from EMEP/EEA Guidebook (EMEP/EEA, 2023). Hectares of total arable crop surface have been used as activity data for PM emissions according to the methodology EMEP/EEA Guidebook (EMEP/EEA, 2023).

HCB emissions from the use of pesticides (3Df) have been estimated. HCB emissions result from the use of HCB as pesticide but also by the use of other pesticides which contain HCB as an impurity. For the period 1996-2001, data are from the database of pesticides contained in the National agricultural information system (*Sistema informativo agricolo nazionale* - SIAN⁴). For the period 2002-2008, SIAN data have been elaborated by Provincial Agency for the Protection of the Environment of the Autonomous Province of Trento⁵. From 2009 activity data have been processed by the Service for risks and environmental sustainability of technologies, chemical substances, production cycles and water services and for inspection activities of ISPRA on the basis of data provided by ISTAT related to substances chlorothalonil, picloram, lindane and chlortal-dimetile which are the active ingredients of pesticides containing HCB.

The availability of data allows estimating emissions from pesticides where HCB is found as an impurity, as in lindane, DCPA, chlorothalonil and Picloram. Emissions from the use of HCB as a pesticide were not estimated. As result from the 2020 NEC review, the HCB emissions from the use of pesticides have been revised. On the basis of the amount of HCB contained in these pesticides (lindane: 0.005%; DCPA: 0.004%; chlorothalonil: 0.004%; Picloram: 0.005%) and according to the EMEP/EEA Guidebook (EMEP/EEA, 2023), which states all the HCB present as a contaminant will be volatilised, HCB emissions result in 118.82 kg for 1990 and 0.00284 kg in 2021. An international research work at European level (Berdowski et al., 1997) estimated 400 kg of HCB emissions from pesticide use for Italy in 1990 while in the last years these emissions should be null.

Detailed information on activity data sources, methods and EFs by pollutant for 3D category is shown in Table 6.16.

⁴ <http://www.sian.it/portale-sian/attivaserivizio.jsp?sid=174&pid=6&servizio=Banca+Dati+Fitofarmaci&bottoni=no>

⁵ http://www.appa.provincia.tn.it/fitofarmaci/programmazione_dei_controlli_ambientali/-Criteri_vendita_prodotti_fitosanitari/pagina55.html

Table 6.16 Activity data sources, methods and emission factors by pollutant for agriculture soils

NFR code	Category	Method	Activity data	Emission Factor
3Da1	Inorganic N-fertilizers (includes also urea application)	T2 (NH ₃), T1 (NO _x)	NS	T2 (NH ₃), D (NO _x)
3Da2a	Animal manure applied to soils	T2 (NH ₃ , NO _x , NMVOC)	NS	CS (NH ₃), D (NO _x), T2 (NMVOC)
3Da2b	Sewage sludge applied to soils	T1 (NH ₃ , NO _x)	NS	D (NH ₃ , NO _x)
3Da2c	Other organic fertilisers applied to soils (including compost)	T1 (NH ₃ , NO _x)	NS	D (NH ₃ , NO _x)
3Da3	Urine and dung deposited by grazing animals	T2 (NH ₃ , NO _x , NMVOC)	NS	CS (NH ₃), D (NO _x), T2 (NMVOC)
3Da4	Crop residues applied to soils	T2 (NH ₃)	NS	T2 (NH ₃)
3Db	Indirect emissions from managed soils			
3Dc	Farm-level agricultural operations including storage, handling and transport of agricultural products	T1 (PM ₁₀ , PM _{2.5})	NS	D (PM ₁₀ , PM _{2.5})
3Dd	Off-farm storage, handling and transport of bulk agricultural products			
3De	Cultivated crops	CS (NH ₃), T1 (NMVOC)	NS	D (NH ₃ , NMVOC)
3Df	Use of pesticides	T1 (HCB)	NS	D (HCB)

6.4 FIELD BURNING OF AGRICULTURAL RESIDUES (3F)

NMVOC, CO, NO_x, NH₃, SO₂, PM₁₀, PM_{2.5}, BC, As, Cr, Cu, Ni, Se, Zn, Pb, Cd, Hg, Dioxin and PAH emissions have been estimated, applying the tier 1 and tier 2 approach (for heavy metals, PAH emissions and BC). A detailed description of the methodology and parameters used is shown in the NIR (ISPRA, several years [a]). The same methodology to estimate emissions from open burning of waste, as reported in paragraph 7.2 of the waste section (see Small scale waste burning (5C2) subparagraph), is used on the basis of the amount of fixed residues instead of removable residues.

Concerning NO_x and CO, IPCC emission factors have been used (IPCC, 2006), while for PM₁₀, PM_{2.5}, NMVOC, BC, NH₃, SO₂, heavy metals, dioxin and PAH emission factors from the EMEP/EEA Guidebook (EMEP/EEA, 2023) have been applied.

6.5 TIME SERIES AND KEY CATEGORIES

The following sections present an outline of the main key categories in the agriculture sector.

The agriculture sector is the main source of NH₃ emissions in Italy; for the main pollutants, in 2022 the sector accounts for:

- 90.4% of national total NH₃ emissions
- 14.6% of national total NMVOC emissions
- 10.2% of national total PM₁₀ emissions
- 7.1% of national total NO_x emissions
- 2.5% of national total PM_{2.5} emissions
- 2.2% of national total Cd emissions

Moreover, the sector comprises 0.4% of BC emissions, 0.5% of CO, 0.5% of PAH, 0.3% of Hg, 0.01% of As, 0.04% of Se, 0.05% of SO₂, 0.02% of Cr, 0.02% of Dioxins, 0.02% of Ni, 0.01% of Zn, 0.01% of Pb and 0.002% of Cu. There are no particular differences as compared to the sectoral share in 1990 when the agriculture sector accounted for 94.4% of NH₃ emissions, 9.4% of PM₁₀, 2.6% of PM_{2.5}, except for NMVOC emissions (7.7%), NO_x emissions (2.9%) and HCB emissions where agriculture accounted for 83.4% of total national emissions.

Table 6.17 reports the key categories identified in the agriculture sector while the time series of NH₃ emissions by sources is shown in Table 6.18.

Concerning NH₃ emissions, the category manure management (3B) represents in 2022 51.8% of national total ammonia emissions (51.3% in 1990). NH₃ emissions from cattle (3B1) stand for 58.4% of 3B emissions, while emissions from swine (3B3) and poultry (3B4g) represent 17.2% and 13.0%, respectively. The category agricultural soils (3D) represents in 2022 38.5% of national total ammonia emissions (43.0% in 1990). The animal manure applied to soils (3Da2a) and the use of synthetic N-fertilisers (3Da1) represent 48.3% and 32.5% of 3D emissions, respectively.

Regarding PM₁₀ emissions, the category manure management (3B) accounts for 4.8% in 2022 (5.4% in 1990) of national total PM₁₀ emissions. Poultry (3B4g), cattle (3B1) and swine (3B3) represent the major contributors to the total PM₁₀ emissions from category 3B with 55.2%, 23.8% and 13.7%, respectively. The category Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc) accounts for 5.0% in 2022 (3.7% in 1990) of national total PM₁₀ emissions. For PM_{2.5} emissions, the category manure management (3B) contributes for 1.8% in 2022 (1.9% in 1990) of national total PM_{2.5} emissions. Cattle (3B1) accounts for 60.2%, while poultry (3B4g) stands for 25.0% to the total PM_{2.5} emissions from category 3B. The category Farm-level agricultural operations including storage, handling and transport of agricultural products (3Dc) accounts for 0.3% in 2022 (0.2% in 1990) of national total PM_{2.5} emissions.

Concerning NO_x emissions, the category manure management (3B) represents in 2022 0.25% of national total NO_x emissions (0.09% in 1990). For NO_x emissions, the category agricultural soils (3D) contributes for 6.8% in 2022 (2.8% in 1990) of national total NO_x emissions. Inorganic N-fertilizers (3Da1) and Animal manure applied to soils (3Da2a) account for 43.4% and 28.4% of total 3D emissions, respectively.

Concerning NMVOC emissions, the category manure management (3B) represents in 2022 11.8% of national total NMVOC emissions (6.2% in 1990). For NMVOC emissions, the category agricultural soils (3D) contributes for 2.8% in 2022 (1.5% in 1990) of national total NMVOC emissions. For NMVOC emissions, the category manure management (3B) and agricultural soils (3D) contributes for 80.8% and 19.0% in 2022 of agricultural NMVOC emissions. Cattle (3B1), poultry (3B4g) and buffalo (3B4a) represent the major contributors to the total NMVOC emissions from category 3B with 75.8%, 12.6% and 5.3%, respectively. Most of the emissions in the 3D category (69.5%) derive from Animal manure applied to soils (3Da2a).

Table 6.17 Key categories in the agriculture sector in 2022.

	SO _x	NO _x	NH ₃	NMVOC	CO	PM ₁₀	PM _{2.5}	BC	Pb	Cd	Hg	PAH	DIOX	HCB	PCB
	%														
3B1a		0.06	13.93	4.60		0.52	0.48								
3B1b		0.08	16.31	4.33		0.63	0.59								
3B2		0.02	0.93	0.10		0.17	0.07								
3B3		0.00	8.91	0.39		0.66	0.04								
3B4a		0.01	2.19	0.62		0.10	0.09								
3B4d		0.00	0.13	0.01		0.03	0.01								
3B4e		0.01	0.80	0.12		0.04	0.04								
3B4f		0.00	0.17	0.01		0.01	0.00								

	SO _x	NO _x	NH ₃	NM VOC	CO	PM10	PM2.5 %	BC	Pb	Cd	Hg	PAH	DIOX	HCB	PCB
3B4gi		0.02	1.84	0.26		0.61	0.07								
3B4gii		0.02	3.39	0.60		1.20	0.17								
3B4giii		0.01	1.03	0.25		0.48	0.12								
3B4giv		0.00	0.48	0.37		0.38	0.09								
3B4h		0.01	1.72	0.13		0.00	0.00								
3Da1		1.93	12.53												
3Da2a		2.96	18.61	1.92											
3Da2b		0.04	0.26												
3Da2c		0.98	3.47												
3Da3		0.90	2.32	0.02											
3Da4			0.91												
3Dc						5.05	0.28								
3De			0.46	0.83											
3Df															
3F	0.05	0.04	0.07	0.03	0.52	0.32	0.43	0.36	0.01	2.24	0.28	0.48	0.02		

Note: key categories are shaded in blue

Table 6.18 Time series of ammonia emissions in agriculture (Gg)

NFR SECTOR 3	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
3B1a Manure management - Dairy cattle	93.01	75.33	70.66	62.20	59.92	55.00	48.95	49.18	48.23	48.87
3B1b Manure management - Non-dairy cattle	88.58	86.73	84.69	73.95	71.28	60.98	65.15	64.32	62.15	57.21
3B2 Manure management - Sheep	4.34	5.30	5.51	3.95	3.92	3.55	3.48	3.49	3.34	3.26
3B3 Manure management - Swine	36.65	34.98	35.54	36.81	34.22	31.71	30.88	31.01	30.46	31.27
3B4a Manure management - Buffalo	3.06	4.54	5.80	6.41	10.35	7.83	7.66	7.64	7.62	7.67
3B4d Manure management - Goats	0.58	0.63	0.63	0.44	0.45	0.44	0.49	0.49	0.49	0.47
3B4e Manure management - Horses	2.20	2.40	2.14	2.13	2.85	2.94	2.81	2.81	2.81	2.79
3B4f Manure management - Mules and asses	0.64	0.29	0.25	0.23	0.35	0.54	0.55	0.55	0.55	0.58
3B4gi Manure management - Laying hens	13.87	12.68	9.56	6.30	7.84	6.87	6.59	6.50	6.54	6.44
3B4gii Manure management - Broilers	12.37	12.96	12.17	11.77	12.00	12.45	12.04	12.28	12.36	11.91


NFR SECTOR 3	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
3B4giii Manure management - Turkeys	7.30	7.24	7.18	5.42	6.08	5.80	4.96	5.16	4.91	3.61
3B4giv Manure management - Other poultry	1.76	4.06	3.15	4.62	2.27	2.70	2.65	2.34	2.24	1.68
3B4h Manure management - Other animals (*)	7.51	8.41	8.79	10.00	8.69	7.72	5.76	5.29	5.22	6.04
3Da1 Inorganic N-fertilizers (includes also urea application)	112.92	114.72	113.17	110.54	74.69	78.45	72.97	89.31	82.41	43.94
3Da2a Animal manure applied to soils	95.14	86.90	79.01	70.17	68.08	66.10	66.50	67.28	66.74	65.27
3Da2b Sewage sludge applied to soils	0.66	1.06	1.42	1.15	1.31	1.08	0.88	0.92	0.88	0.90
3Da2c Other organic fertilisers applied to soils (including compost)	1.32	1.45	1.81	1.78	3.29	6.68	6.84	9.82	8.81	12.16
3Da3 Urine and dung deposited by grazing animals	10.09	11.33	11.48	8.74	8.97	8.60	8.51	8.55	8.33	8.15
3Da4 Crop residues applied to soils	5.07	5.17	4.75	4.56	3.89	3.29	3.69	3.75	3.37	3.18
3De Cultivated crops	2.38	1.58	1.59	1.41	1.38	1.48	1.56	1.54	1.53	1.60
3F Field burning of agricultural residues	0.48	0.47	0.47	0.42	0.30	0.29	0.27	0.27	0.28	0.26
Total	499.90	478.23	459.77	423.01	382.15	364.49	353.19	372.49	359.25	317.25

Note: (*) 3B4h includes rabbits and fur animals

The largest and most intensive agricultural area in Italy is the Po River catchment with the following characteristics: high crop yields due to climatic factors, double cropping system adopted by livestock farms, flooded rice fields, high livestock density and animal production that keep animals in stables all the year (Bassanino et al 2011, Bechini and Castoldi 2009). 64%, 78% and 88% of cattle, poultry and swine production are located in Piedmont, Lombardy, Emilia-Romagna, and Veneto Regions (Northern Italy/Po River Basin). At regional level, the presence of large cattle, poultry and swine farms in the Po basin assume a particular relevance for air quality issues, especially, for the specific meteorological conditions of this area.

In the Figure 6.2 NH₃ emissions from main categories of 3B are reported.

Figure 6.2 NH₃ emissions from 3B cattle, swine and poultry categories.



The reduction of NH₃ emissions from 3B is mainly related to the reduction in the number of animals. Between 1990 and 2022 total NH₃ emissions from 3B have reduced by 33.1%. Cattle livestock decreased by 27.3% (from 7,752,152 to 5,632,736 heads). Dairy cattle and non-dairy cattle have decreased by 38.3% and 21.7%, respectively. The so-called first pillar of the EU Common Agriculture Policy (CAP), dealing with market support, had a strong impact through the milk quota system by reducing animal numbers in the dairy sector to compensate for increasing animal productivity (EEA, 2016).

Swine have increased between 1990 and 2022 by 4.0% while poultry have decreased by 4.7% (see Table 6.1). Abatement technologies are considered in the EFs used for NH₃ estimations. Research studies funded by ISPRA, such as the MeditAiraneo project, or by the Ministry of Environment have allowed us to collect information on the inclusion of abatement technologies in Italy, especially those related to the swine and poultry recovery and treatment of manure and to land spreading (CRPA, 2006[b]; C3ndor et al., 2008; CRPA, 2010[b]).

With regard to the number of swine, the increase in 2001 and in the following years was due to an increase in demand for pork meat favoured by more competitive prices compared to other meats, but also to the diseases affecting beef (Bovine Spongiform Encephalopathy - BSE) and poultry (bird flu). The reduction in the number of swine in 2012, which affected all sub-categories (piglets, fattening pigs and breeding pigs), was affected by the implementation of Directive 2008/120, which set new standards in terms of the welfare of pigs and in particular sows in breeding wards.

Emission abatement techniques change the emission factors (which are a weighted average of the emission factors between the reference, unabated system, and systems that emit less). The change in emission factors combined with the change in the number of animals changes the emission trend.

NH₃ emissions of 3D category are driven by the animal manure applied to soils and the use of inorganic N-fertilizers. Between 1990-2022 emissions have respectively decreased by 31.4% and 61.1% mainly due to the reduction of the number of animals and the use of inorganic N-fertilizers, that are decreased overall by 60.5% (in particular urea decreased by 53.6%), in terms of nitrogen content.


The interannual variability of ammonia emissions from synthetic fertilisers depends entirely on the data provided by the National Institute of Statistics (ISTAT) on the distributed quantities of synthetic fertilisers and, in particular, on the quantities of nitrogen they contain. While the emission factors are constant throughout the historical series. For the years 2012 and 2016, the increase in distributed urea almost completely explains the overall increase in emissions from synthetic fertilisers. In 2020, 70% of the increase is explained by the increase in urea and 30% by the increase in other synthetic fertilisers.

According to Assofertilizzanti–Federchimica⁶, in relation to the 2020-2022 data, farmers tend to anticipate purchases (and thus stockpile) when expectations of rising market prices are present. The 2022 data are lower than annual average (for nitrogen, phosphorus pentoxide, and potassium oxide) because they follow a two-year period in which quantities purchased increased and refer to a year in which high market prices prompted operators to delay purchases in anticipation of falling prices, an event, which actually occurred during 2023.

Furthermore, the EU Nitrates Directive which aims at reducing and preventing water pollution caused by nitrates from agricultural sources has addressed the lower use of synthetic and nitrogen-based fertilisers (EEA, 2016).

Every 4 years (first every 5) the national emission inventory is disaggregated at NUTS3 level as requested by CLRTAP (Córdoba et al., 2008). A database with the time series for all sectors and pollutants has been published (ISPRA, 2022; ISPRA, 2021; ISPRA, 2009; ISPRA, several years[c]; ISPRA, several years[d]). The disaggregation of 2019 agricultural emissions has also been finalised and figures are available at the following web site: <https://emissioni.sina.isprambiente.it/inventari-locali/> The disaggregation (NUTS3) of the NH₃ agricultural emissions is shown in Figure 6.3. In 2019, four regions contributed with more than 60% of agricultural NH₃ emissions: Lombardia, Veneto, Emilia Romagna and Piemonte.

Figure 6.3 NH₃ emissions from Agriculture in 2019 (t)



⁶ Federchimica is the National Association of the Chemical Industry and Assofertilizzanti represents the production companies of the fertilizer industry.

6.6 QA/QC AND VERIFICATION

QA/QC procedures for the agriculture sector are in line with the 2006 IPCC Guidelines and consistent with the EMEP/EEA Guidebook. Italy has drawn up a QA/QC procedure manual and elaborates annually a QA/QC plan both for the UNFCCC and UNECE/CLRTAP inventories. In the QA/QC Agriculture section GHG and NH₃ emissions improvements are specified (ISPRA, several years [b]). Furthermore, feedback for the agricultural emission inventory derives also from communication of data to different institutions (ISTAT, UNAITALIA, CRPA etc.) and/or at local level (regional environmental institutions). In addition, ISPRA participates in a technical working group on agriculture within the National Statistical System, composed by producers and users of agricultural statistics.

In 2011 a validation of EFs and estimations was carried out considering the results of a research study that estimated at NUTS 2 level emissions for the use of synthetic N-fertilizers considering type of cultivation, altitude, and climatic conditions (CRPA, 2010[b]; C ndor and Valli, 2011).

Data used to estimate emissions were verified with census data. Differences in the livestock number are found comparing conjunctural surveys (used for emissions estimation) and the Agricultural census for the year 2020: +5% non-dairy cattle, +3% poultry, -2% buffalo and swine, +12% goats, while equines are almost three times as large.

Data on national sales of synthetic nitrogen fertilizers (by type of fertilizers) as provided by Assofertilizzanti (data are available online - <https://assofertilizzanti.federchimica.it/fertilizzanti/statistiche-fertilizzanti>) for the period 2011-2022 have been compared to official statistics provided by ISTAT. ISTAT simple mineral nitrogen fertilizers data are about 15% lower than those of Assofertilizzanti, for the years 2011-2022. These differences in recent years could be attributable to rising fertilizer prices. ISTAT statistics almost certainly do not take into account that farmers tend to anticipate purchases (and thus stockpile) when expectations of rising market prices are present. The 2022 data are lower than annual average (for nitrogen, phosphorus pentoxide, and potassium oxide) because they follow a two-year period in which quantities purchased increased and refer to a year in which high market prices prompted operators to delay purchases in anticipation of falling prices, an event, which actually occurred during 2023.

Ammonia emissions for swine and poultry manure management from housing and storage were compared with data reported in the E-PRTR registry for the year 2019 for swine and poultry, which represent 36.8% of national NH₃ emissions for the same categories (3B).

In 2021, a technical report (CREA, 2021) produced by CREA, in collaboration with ISPRA, on the assessment of emissions related to the use of nitrogen fertilizers was published, which contains several analyses of nitrogen fertilizer consumption data from different data sources: IFASTAT/Fertilizer Europe, FAO, EUROSTAT, Assofertilizzanti, National Integrated Production Specifications, Farm Accountancy Data Network (FADN). These analyses suggest further investigation to better understand the differences found between these data sources and the ISTAT data used for the estimates.

In 2021, NH₃ emissions for dairy cows were also calculated with the AgrEE tool - Agricultural Emission Estimation tool (made by the JRC) and differences with national estimates for the year 2015 were analysed. National estimates are slightly higher than the AgrEE tool results (16.5 kt N-NH₃ for spreading vs 16.1 kt N-NH₃) for several reasons. In particular two of them are: for N₂O emissions from storage, according to the Guidebook, it is necessary to remove emissions from housing, while IPCC considers the N excreted at the housing; for NH₃ emissions from spreading, the Guidebook does not remove N lost through leaching and run-off at storage, unlike the IPCC methodology. The verification with AgrEE tool revealed other differences that will be taken into account for improved emission estimates.

6.7 RECALCULATIONS

In 2024, recalculations were implemented for the agricultural emission inventory.

Changes related to NH₃ emissions are described below.

Major recalculations of NH₃ emissions are due to: update of NH₃ emission factors from synthetic fertilizers, reported in the EMEP/EEA Guidebook 2023; change in the coefficients of N contained in livestock manure

sent to anaerobic digestion of cattle, swine, poultry since 1991; correction of NH₃ EFs from storage for cattle in the years 2003-2005; correction of NH₃ EFs from storage for cattle and pigs since 2014; correction of N values of synthetics for the year 2016, N values of soil improvers for the years 2015 and 2016, N values of other organics for the year 2016.

The estimate of NH₃ from the residues was added.

In addition, emissions from spreading changed throughout the time series as a result of: change in emissions from housing and storage; change in emissions of NH₃ from anaerobic digesters; change in emissions of N₂O from storage; correction of the average weight of the subcategory other cattle < 1 year of 2021 (that affects the NO₂ and N₂ emissions, N leached by manure management and N bedding); change in the distribution of dairy cow housing in the years 2006-2009 goes to affect the allocation of effluent between liquid and solid, which therefore also affects the NO₂ emissions from dairy cow storage years 2006-2009 (and thus N₂ emissions and the estimate of N bedding were also changed); N correction to calves in the years 1990-2015 goes to affect NO₂ emissions storage by non-dairy cattle category (and thus N₂ emissions were also changed).

Minor recalculations are due to: correction of NH₃ EFs from housing for dairy cows for the years 2006-2009; correction of the average weight of the subcategory other cattle < 1 year of 2021; update of the sludge data for the years 2018-2021; correction of the N value of calves (included in 'less than 1 year for the slaughter' non-dairy cattle category) for the years 1990-2015.

The following updates resulted in a recalculation in emissions from biomass combustion: the estimate of sorghum residue since 1990 has been changed, now aligned with the estimate of emissions from agricultural soils; the percentage of rice straw burned since 2001 has been changed, and the production of the years 2019-2021 has been updated.

Changes related to NO₂ emissions are described below.

The recalculation of NO₂ emissions is due to: correction of N values of synthetics for the year 2016, N values of soil improvers for the years 2015 and 2016, N values of other organics for the year 2016; updating of emission factor for biomass combustion substituted with that in the 2023 EMEP/EEA Guidebook; changes in NH₃ emissions from spreading; update of the sludge data for the years 2018-2021; correction of the average weight of the subcategory other cattle < 1 year of 2021; correction of the N value of calves (included in 'less than 1 year for the slaughter' non-dairy cattle category) for the years 1990-2015; change in the distribution of dairy cow housing in the years 2006-2009 goes to affect the allocation of effluent between liquid and solid, which therefore also affects the NO₂ emissions from dairy cow storage years 2006-2009.

Changes related to PM emissions are described below.

The recalculation of PM emissions from livestock is due to: correction of the average weight of the subcategory other cattle < 1 year of 2021; correction of the N value of calves (included in 'less than 1 year for the slaughter' non-dairy cattle category) for the years 1990-2015; change in the distribution of dairy cow housing in the years 2006-2009 goes to affect the allocation of effluent between liquid and solid, which therefore also affects the PM emissions from dairy cow storage years 2006-2009.

The recalculation of PM emissions from farm-level agricultural operations is due to update of surface data of polyphitic grassland and waxy corn grassland for 2017 and of other vegetables (including in chard) of 2021.

Changes related to NMVOC emissions are described below.

The recalculation of NMVOC emissions from cultivated crops is due to the update of surface data of polyphitic grassland and waxy corn grassland for 2017. The recalculation of NMVOC emissions from manure management is due to changes in NH₃ emissions from housing and storage and the correction of the average weight of the subcategory other cattle < 1 year of 2021. The recalculation of NMVOC emissions from Animal manure applied to soils is due to changes in NH₃ emissions from housing and spreading and the correction of the average weight of the subcategory other cattle < 1 year of 2021. The

recalculation of NMVOC emissions from grazing animals is due to the correction of the average weight of the subcategory other cattle < 1 year of 2021.

As regards HCB emissions, the recalculation is due to the update of 2021 activity data.

As regards field burning of agricultural residues emissions, the recalculation of all emissions is due to: updating of emission factors for CO, NO_x, NMVOC, PM₁₀, PM_{2.5}, substituted with those in the 2023 EMEP/EEA Guidebook and in the 2006 IPCC guidelines (for CO, NO_x); the estimate of sorghum residue since 1990 has been changed, now aligned with the estimate of emissions from agricultural soils; the percentage of rice straw burned since 2001 has been changed, and the production of the years 2019-2021 has been updated.

6.8 PLANNED IMPROVEMENTS

In the coming years, the Permanent census of agriculture will provide valuable information on animal and agronomic production methods. The focus of the Permanent census is to provide a comprehensive information framework on the structure of the agricultural system and the livestock at national, regional and local level by integrating archive data and carrying out statistical support surveys. Statistical registers will be created with the aim of increasing the quantity and quality of information to reduce the response.

An update of the information on AWMS and abatement technologies will be carried out with the data of the 2020 Census of Agriculture as soon as data processing carried out on the micro-data released in January 2024 by the National Statistical Institute (ISTAT) is completed.

7 WASTE (NFR SECTOR 5)

7.1 OVERVIEW OF THE SECTOR

Italy estimates the categories of the waste sector, as reported in the following box. In the last submissions Cd, Hg and Pb emissions from 5E have been estimated too, as well as NH₃ emission from municipal waste incineration. In the current submission, Italy has reported NH₃ emissions from latrines (EEA, 2023) and PM from the disposal of mineral waste have been estimated. In the last available national census on wastewater treatment plants (ISTAT, 2015) the following data are reported: 99.4% of people are served by the sewage system, 17,897 wastewater system plants serve a total of 98,360,724 people equivalent; 8,377 are the Imhoff tanks present in Italy, 1,607 are the primary wastewater treatment plants, 5,604 are the secondary wastewater treatment plants and 2,309 are the advanced wastewater treatment plants. The biogas collected from the anaerobic digestion of wastewaters is burned with heat/energy recovery and relevant emissions are reported in Category 1 while emissions from the exceeding biogas which is flared are not estimated at the moment because emission factors are under investigation, but anyway it should be negligible.

NFR		SNAP	
5A	Solid waste disposal on land	09 04 01	Managed waste disposal on land
		09 04 02	Unmanaged waste disposal on land
5B	Biological treatment of waste	09 10 05	Compost production
		09 10 06	Anaerobic digestion at biogas facilities
5C1a	Municipal waste incineration	09 02 01	Incineration of municipal wastes
5C1b	Other waste incineration	09 02 02	Incineration of industrial wastes
		09 02 05	Incineration of sludge from wastewater treatment
		09 02 07	Incineration of hospital wastes
		09 02 08	Incineration of waste oil
5C1bv	Cremation	09 09 01	Cremation of corpses
5C2	Small scale waste burning	09 07 00	Open burning of agricultural wastes
5D	Wastewater handling	09 10 01	Waste water treatment in industry
		09 10 02	Waste water treatment in residential and commercial sector
		09 10 07	Dry toilets (including latrines)
5E	Other waste	09 10 XX	Car and building fires

Concerning air pollutants, emissions estimated for each sector are reported in Table 7.1.

Table 7.1 Air pollutant emissions estimated for each sector.

	5A	5B	5C1a	5C1bi	5C1bii	5C1bii i	5C1bi v	5C2	5C1bv	5D	5E
Main pollutants											
NO _x			x	x	x	x	x	x	x		
CO			x	x	x	x	x	x	x		
NMVO C	x	x	x	x	x	x	x	x	x	x	
SO _x			x	x	x	x	x	x	x		
NH ₃	x	x	x								x

	5A	5B	5C1a	5C1bi	5C1bii	5C1bii i	5C1bi v	5C2	5C1bv	5D	5E
Particulate matter											
TSP	x		x	x	x	x	x	x	x		x
PM10	x		x	x	x	x	x	x	x		x
PM2.5	x		x	x	x	x	x	x	x		x
Priority heavy metals											
Pb			x	x	x	x	x	x	x		x
Cd			x	x	x	x	x	x	x		x
Hg			x	x	x	x	x	x	x		x
POPs Annex II											
PCB			x	x		x	x		x		
POPs Annex III											
Dioxins			x	x	x	x	x	x	x		x
PAH			x	x	x	x	x	x	x		
HCB			x	x		x	x		x		
Other heavy metals											
As			x	x	x	x	x	x	x		
Cr			x	x	x	x	x	x	x		
Cu			x	x	x	x	x	x	x		
Ni			x	x	x	x	x			x	
Se			x	x		x		x	x		
Zn			x	x	x		x	x	x		

In 2022, open burning of waste (5C2) is key category for Cd and CO, car and building fires (5E) is a key category for PM and dioxins emissions while clinical waste incineration (5Cbiii) is key category for HCB. In 1990, municipal waste incineration (5C1a), industrial waste incineration (5C1 bi) and car and building fires (5E) are key categories for dioxins emissions. As regard the trend, municipal waste incineration (5C1a) and car and building fires (5E) are key categories for dioxins emissions, biological treatment of waste (5B2) is key category for NH₃ emissions, whereas open burning of agricultural waste (5C2) is key category for Cd emissions.

The waste sector, and in particular Waste incineration (5C), is a source of different pollutants; for the main pollutants, in 2022, the sector accounts for:

- 16.5 % in national total Dioxin emissions;
- 8.7 % in national total BC emissions;
- 8.6% in national total HCB emissions;
- 7.7 % in national total Cd emissions .

Moreover, the sector comprises 3.5% of total NH₃ emissions, 3.2% of CO, 1.8% of Hg, 1.0% of NMVOC, 1.5% of PAH, 3.9% and 2.8% in national total of PM_{2.5} and PM₁₀ emissions respectively and for what concerns all remaining pollutants are below 1%.

7.2 METHODOLOGICAL ISSUES

7.2.1 Solid waste disposal on land (5A)

Solid waste disposal on land is a major source concerning greenhouse gas emissions but not concerning air pollutants. Notwithstanding, NMVOC and NH₃ emissions are estimated, as a percentage of methane emitted, calculated using the IPCC Tier 2 methodology (IPCC, 1997; IPCC, 2000), through the application of the First Order Decay Model (FOD). As a consequence of the last review process also PM emissions have been estimated. A detailed description of the model and its application to Italian landfills is reported in the National Inventory Report on the Italian greenhouse gas inventory (ISPRA, 2024 [a]).

Following the suggestion of NEC review (EEA, 2017 [a]) more info about the extraction and use of biogas is provided below. The amount of biogas recovery in landfills has increased as a result of the implementation of the European Directive on the landfill of waste (EC, 1999); the amounts of biogas recovered and flared have been estimated taking into account the amount of energy produced, the energy efficiency of the methane recovered, the captation efficiency and the efficiency in recovering methane for energy purposes assuming that the rest of methane captured is flared. Emissions for all the relevant pollutants from biogas recovered from landfills and used for energy purposes are reported in the energy sector in "1A4a biomass" category together with wood, the biomass fraction of incinerated waste and biogas from wastewater plants. In Table 7.2 consumptions and low calorific values are reported for the year 2022.

Table 7.2 1A4a biomass detailed activity data. Year 2022

Fuels		Consumption (Gg)	LCV (TJ/Gg)
Wood and similar	Wood	317.99	10.47
	Steam Wood	0.00	30.80
Incinerated waste (biomass)		2065.54	11.26
Biogas from landfills		199.16	54.34
Biogas from wastewater plants		23.28	54.34

It is assumed that landfill gas composition is 50% VOC. The percentage by weight of CH₄ compared to the total VOC emitted is 98.7%. The remaining 1.3% (NMVOC) consists of paraffinic, aromatic and halogenated hydrocarbons (Gaudioso et al., 1993); this assumption refers to US EPA data (US EPA, 1990). As regard ammonia, emission factor has been assumed equal to 1 volume per cent of VOC too (Tchobanoglous et al., 1993). According with the discussion during the ESD review about CH₄ emissions from landfills and the consequent technical correction (EEA, 2017 [b]), Italy revised the half-life values considering the distribution of dry and wet regions in Italy. New data (CREA, 2017) regarding raining and evapotranspiration have been elaborated allowing to distinguish between dry and wet region and the estimates have been splitted in two components considering the location of SWDS. Methane, and consequently NMVOC and NH₃ air pollutants, is emitted from the degradation of waste occurring in municipal landfills, both managed and unmanaged (due to national legislation, from 2000 municipal solid wastes are disposed only into managed landfills). The main parameters that influence the estimation of emissions from landfills are, apart from the amount of waste disposed into managed landfill: the waste composition (which vary through the years in the model); the fraction of methane in the landfill gas (included in VOC, which has been assumed equal to 50%) and the amount of landfill gas collected and treated. These parameters are strictly dependent on the waste management policies throughout the waste streams which consist of: waste generation, collection and transportation, separation for resource

recovery, treatment for volume reduction, stabilisation, recycling and energy recovery and disposal at landfill sites. Basic data on waste production and landfills system are those provided by the national Waste Cadastre, basically built with data reported through the Uniform Statement Format (MUD). The Waste Cadastre is formed by a national branch, hosted by ISPRA, and by regional and provincial branches. These figures are elaborated and published by ISPRA yearly since 1999: the yearbooks report waste production data, as well as data concerning landfilling, incineration, composting, anaerobic digestion and generally waste life-cycle data (APAT-ONR, several years; ISPRA, several years [a]).

For inventory purposes, a database of waste production, waste disposal in managed and unmanaged landfills and sludge disposal in landfills was created and it has been assumed that waste landfilling started in 1950. For the year 2022, the non-hazardous landfills in Italy disposed 5,619 kt of MSW and 2,963 kt of industrial wastes, as well as 247 kt of sludge from urban wastewater treatment plants. In Table 7.3, the time series of AMSW and domestic sludge disposed into non-hazardous landfills from 1990 is reported.

Table 7.3 Trend of MSW production and MSW, AMSW and domestic sludge disposed in landfills (Gg)

ACTIVITY DATA (Gg)	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
MSW production	22,231	25,780	28,959	31,664	32,479	29,524	30,079	28,941	29,596	29,051
MSW disposed in landfills for non-hazardous waste	17,432	22,459	21,917	17,226	15,015	7,819	6,283	5,817	5,619	5,173
Assimilated MSW disposed in landfills for non-hazardous waste	2,828	2,978	2,825	2,914	3,508	3,222	3,256	2,910	2,963	2,469
Sludge disposed in managed landfills for non-hazardous waste	2,454	1,531	1,326	544	346	387	232	253	106	96
Total Waste to managed landfills for non-hazardous waste	16,363	21,897	26,069	20,684	18,870	11,428	9,771	8,980	8,687	7,738
Total Waste to unmanaged landfills for non-hazardous waste	6,351	5,071	-	-	-	-	-	-	-	-
Total Waste to landfills for non-hazardous waste	22,714	26,968	26,069	20,684	18,870	11,428	9,771	8,980	8,687	7,738

7.2.2 Biological treatment of waste (5B)

Under this category, NMVOC and NH₃ emissions from compost production and from anaerobic digestion are reported. The amount of waste treated in biological treatments has shown a great increase from 1990 to 2022 (from 283,879 Mg to 6,435,007 Mg for composting and from 0 to 96,604 Mg of N-excreted from manure management). Information on input waste to composting plants is published yearly by ISPRA since 1996, including data for 1993 and 1994 (ANPA, 1998; APAT-ONR, several years; ISPRA, several years [a]), while for 1987 and 1995 only data on compost production are available (MASE, several years [a]; AUSITRA-Assoambiente, 1995); on the basis of this information the whole time series has been reconstructed. The composting plants are classified in two different kinds: the plants that treat a selected waste (food, market, garden waste, sewage sludge and other organic waste, mainly from the agro-food industry); and the mechanical-biological treatment plants, that treat the unselected waste to produce compost, refuse derived fuel (RDF), and a waste with selected characteristics for landfilling or incinerating system. It is assumed that 100% of the input waste to the composting plants from selected waste is treated as compost, while in mechanical-biological treatment plants 30% of the input waste is treated as compost on the basis of national studies and references (Favoio and Cortellini, 2001; Favoio and Girò, 2001). NMVOC emission factor (51g NMVOC kg⁻¹ treated waste) is from international scientific literature

too (Finn and Spencer, 1997). Following the recent NEC review (EEA, 2022), more information is provided below. In Italy, almost all of the plants are industrial plants, with enclosed areas for rotting and decomposition served by biofilters, turning when needed to maintain the right porosity and, above all, forced ventilation or suction system at least since the 2000s also because requested by regional law. To ensure high quality compost the CIC (Consorzio Italiano Compostatori) introduced a few years ago a certification procedure that, in 2014, covers the 28% of national capacity regarding compost from selected waste (ISPRA, 2017). As reported in (ISPRA, 2017 Update of CH₄ emission factor from composting. Technical note n.1/2017), the plant engineering framework in Italy is very advanced being characterized by industrial scale plants and the technical requirements and specific air supplies normally required in environmental permits typically consider a high excess of air, as only 10% of the flow of air required corresponds to the stoichiometric quantity for biodegradation while the remaining 90% is required for heat drainage so that the system is maintained in mesophilic conditions; naturally, the excess oxygen in the atmosphere inside the compost heaps tends to keep the system fundamentally aerobic and to avoid specific methane production. This plant framework was made possible by the regulations that supported it since the early 2000s, while less information was available in the 90s. At the moment Italy is looking for further information to satisfy the request of the review process (EEA, 2023) and be able to introduce the penetration rate of abatement technologies.

NH₃ emissions from biogas facilities (anaerobic digesters) in the agriculture sector have been updated in the previous submission on the basis of the study carried out by CRPA (CRPA, 2018) and in particular data relative to the percentages of the different substrates that feed the anaerobic digesters and data relative to the average content of volatile solids by type of substrates have been changed. As a result of these changes, the amount of manure sent to the digesters decreases considerably and also the NH₃ emissions. These emissions have been subtracted from 3B manure management category (cattle, swine and poultry) and allocated in the anaerobic digestion at biogas facilities (5B2 of the waste sector).

7.2.3 Waste Incineration (5C1a – 5C1b)

Regarding waste incineration, methodology used for estimating emissions is based on and consistent with the EMEP/CORINAIR Guidebook (EMEP/EEA, 2019). In this sector only emissions from facilities without energy recovery are reported, whereas emissions from waste incineration facilities with energy recovery are reported in the Energy Sector 1A4a because energy produced in incinerators is still prevalently used to satisfy the internal energy demand of the plants (auto production) and in this sense it would be wrong, according to the guidelines, to report them under 1A1a Public Electricity and Heat Production instead of 1A4a. In 2022, about 99% of the total amount of urban waste incinerated is treated in plants with energy recovery system. Existing incinerators in Italy are used for the disposal of municipal waste, together with some industrial waste, sanitary waste and sewage sludge for which the incineration plant has been authorized by the competent authority. Other incineration plants are used exclusively for industrial and sanitary waste, both hazardous and not, and for the combustion of waste oils, whereas there are plants that treat residual waste from waste treatments, as well as sewage sludge.

A complete database of the incineration plants is now available, updated with the information reported in the yearly report on waste production and management published by ISPRA (APAT-ONR, several years; ISPRA, several years). For each plant a lot of information is reported, among which the year of the construction and possible upgrade, the typology of combustion chamber and gas treatment section, energy recovery section (thermal or electric), and the type and amount of waste incinerated (municipal, industrial, etc.). A specific emission factor is therefore used for each pollutant combined with plant specific waste activity data.

In Table 7.4, emission factors for each pollutant and waste typology are reported. Emission factors have been estimated on the basis of a study conducted by ENEA (De Stefanis P., 1999), based on emission data from a large sample of Italian incinerators (FEDERAMBIENTE, 1998; AMA-Comune di Roma, 1996), legal thresholds (Ministerial Decree 19 November 1997, n. 503 of the Ministry of Environment; Ministerial Decree 12 July 1990) and expert judgements. Since 2010, emission factors for urban waste incinerators have been updated on the basis of data provided by plants (ENEA-federAmbiente, 2012; De Stefanis P., 2012) concerning the annual stack flow, the amount of waste burned and the average concentrations of

the pollutants at the stack. As the emission factors are considerably lower than the old ones due to the application of very efficient abatement systems it was necessary to apply a linear smoothing methodology assuming a progressive application of the abatement systems between 2005 and 2010. In a similar way, emission factors for industrial waste incinerators have been updated from 2010 onwards on the basis of the 2019 EMEP/EEA Guidebook. Similarly to municipal waste smoothing has been applied between 2005 and 2010 supposing a linear application of the abatement systems. In particular, for 5C1bi the HCB emission factors comes from table 3-1 of the 2019 EMEP/EEA Guidebook considering an abatement efficiency of 90% while PCB emission factors derives from 2007 Guidebook but considering 90% abatement (no value in the 2019 EMEP/EEA Guidebook). For 5C1biii the emission factors from 2019 EMEP/EEA Guidebook have been used for PCB while HCB emission factors derives from 2007 Guidebook because of a lack of information in the last guidebook. For 5C1biv emission factors from 2019 EMEP/EEA Guidebook have been considered without abatement because information about this specific category of plant is under investigation.

Table 7.4 Emission factors for waste incineration

Air Pollutant	u.m.	Municipal 1990-2009	Municipal since 2010	Industrial 1990-2009	Industrial since 2010	Clinical 1990-2009	Clinical since 2010	Sludge 1990-2009	Sludge since 2010	Oil 1990-2009	Oil since 2010
NO _x	kg/t	1.15	0.62	2.00	2.00	0.60	0.60	3.00	3.00	2.00	2.00
CO	kg/t	0.07	0.07	0.56	0.56	0.08	0.08	0.60	0.60	0.08	0.08
NM VOC	kg/t	0.46	0.46	7.40	7.40	0.70	0.70	0.25	0.25	7.40	7.40
SO ₂	kg/t	0.39	0.02	1.28	1.28	0.03	0.03	1.80	1.80	1.28	1.28
PM ₁₀	g/t	46.00	6.06	240.00	7.00	25.68	25.68	180.00	41.00	240.00	0.07
PM _{2.5}	g/t	30.89	4.07	240.00	4.00	25.68	25.68	180.00	11.00	240.00	0.04
As	g/t	0.05	0.02	0.12	0.00	0.00	0.00	0.50	0.24	0.12	0.00
Cu	g/t	1.00	0.00	1.20	0.12	0.56	0.56	10.00	2.00	1.20	0.01
Se	g/t	0.01	0.01	0.01	0.00	0.04	0.04	-	0.01	0.01	0.00
Zn	g/t	0.02	0.02	12.60	1.26	-	-	10.00	3.30	12.60	0.13
Cd	g/t	0.25	0.01	0.80	0.01	0.00	0.00	1.20	0.80	0.80	0.00
Cr	g/t	0.45	0.00	1.60	0.16	0.01	0.01	3.00	0.70	1.60	0.02
Hg	g/t	0.15	0.03	0.80	0.01	0.04	0.04	1.20	1.15	0.80	0.01
Ni	g/t	16.35	0.00	0.80	0.01	0.03	0.03	3.00	0.40	0.80	0.00
Pb	g/t	1.35	1.04	24.00	0.13	0.02	0.02	3.00	0.40	24.00	0.01
PAH	g/t	0.05	0.00	0.48	0.00	0.00	0.00	0.60	0.00	0.48	0.00
PCB	g/t	0.005	0.00005	0.0050	0.0005	0.020	0.020	0.005	0.0045	-	-
HCB	g/t	0.001	0.00002	0.0001	0.0002	0.019	0.019	0.500	0.0047	-	-
NH ₃	g/t	3.00	3.00	-	-	-	-	-	-	-	-

Concerning dioxin emissions, clinical and industrial emission factors are also derived from data collected from a large sample of Italian incinerators and legal thresholds, as well as expert judgement; in particular for municipal solid waste, emission factors vary within the years and the facility on the basis of plant

technology (i.e. typology of combustion chamber and gas treatment section) and the year of the upgrade. This site specific evaluation has been possible thanks to a study conducted in the past for a sample of municipal waste incinerators located in Regione Lombardia in order to produce an assessment of field-based values applicable to other facilities with the same characteristics (Pastorelli et al., 2001) and, since 2010 urban waste data, thanks to the abovementioned survey (ENEA-federAmbiente, 2012). Moreover, for the incineration plants reported in the national EPER/PRTR register, verification of emissions has been carried out. In Table 7.5 dioxin emission factors for waste incineration are reported for 1990 and 2022.

Table 7.5 Dioxin emission factors for 1990 and 2022

Waste Typology	u.m	1990	2022
Municipal	g/t	115 - 1.6	0.116
Clinical	g/t	200	0.500
Industrial	g/t	80 - 135	0.500
Sludge	g/t	77	0.500
Oil	g/t	200	0.500

In Table 7.6 activity data are reported by type of waste.

Table 7.6 Amount of waste incinerated by type (Gg)

Waste incinerated	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Total waste	1,656.2	2,149.1	3,061.7	4,964.2	6,977.3	7,534.6	7,648.0	7,479.6	7,484.0	7,233.6
with energy recovery	911.2	1,557.8	2,749.7	4,720.6	6,795.9	7,431.3	7,558.5	7,409.7	7,373.4	7,123.6
without energy recovery	745.0	591.3	312.0	243.5	181.4	103.2	89.5	69.8	110.6	110.0
Municipal waste (5C1a)	1,025.6	1,436.6	2,324.9	3,219.9	4,336.9	4,698.4	4,358.9	4,227.2	4,304.9	4,131.1
with energy recovery	626.4	1,185.5	2,161.4	3,168.0	4,284.0	4,698.4	4,358.9	4,227.2	4,304.9	4,131.1
without energy recovery	399.2	251.1	163.5	51.9	52.9	0.0	0.0	0.0	0.0	0.0
Industrial waste (5C1b i-ii-iv)	496.1	560.7	626.5	1,618.1	2,505.3	2,734.6	3,183.5	3,145.0	3,065.0	2,989.0
with energy recovery	259.5	331.2	511.6	1,447.0	2,399.4	2,676.0	3,138.4	3,119.0	3,005.2	2,929.5
without energy recovery	236.6	229.6	114.8	171.1	105.9	58.6	45.2	25.9	59.8	59.5
Clinical waste (5C1biii)	134.5	151.7	110.3	126.2	135.1	101.6	105.6	107.4	114.1	113.5
with energy recovery	25.3	41.1	76.7	105.7	112.5	57.0	61.2	63.5	63.3	63.0
without energy recovery	109.2	110.6	33.6	20.5	22.6	44.6	44.4	43.9	50.8	50.5

7.2.4 Cremation of corpses (5C1bv)

Emissions from incineration of human bodies in crematoria have been carried out for the entire time series. The methodology used for estimating emissions is based on and conform to the EMEP/EEA Air Pollutant Emission Inventory Guidebook (EMEP/EEA, 2023).

Activity data have been supplied by a specific branch of Federutility, which is the federation of energy and water companies (SEFIT, several years), whereas emission factors derive from a survey carried out by the same subject in 2015 and in 2019. For some metal, such as Pb, Cd, As, Cr, Cu, Ni and Se EFs are those reported in the Guidebook 2023. Up to some years ago cremation was not so popular in Italy also because the Catholic Church encouraged burial. Partly because cemeteries are becoming overcrowded, in addition to Covid emergency, the number of cremations in Italy has risen from 5,809 in 1990 to 259,915 in 2022, higher than the peak value of 247,840 in 2020. Moreover, it is practice cremating also mortal remains: activity data have been supplied too by SEFIT, from 1999, whereas mortal remains from 1990 to 1998 have been reconstructed on the basis of an expert judgment (SEFIT, several years).

In Table 7.7 time series of number of cremations, mortal remains, as well as annual deaths and crematoria in Italy are reported. The major emissions from crematoria are nitrogen oxides, carbon monoxide, sulphur dioxide, particulate matter, mercury, hydrogen fluoride (HF), hydrogen chloride (HCl), NMVOCs, other heavy metals, and some POPs. As a consequence of the review process, mortal remains time series has been revised: mortal remains have been converted into corpses equivalents, multiplying data for 75/90 ratio, according with national experts (Sefit, 2022), based on the average time of cremation (90 minutes for a corpse and 75 minutes for a mortal remain. In Table 7.8 emission factors for cremation 2022 are reported.

Table 7.7 Cremation time series (activity data)

Cremation of corpses	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Cremations	5,809	15,436	30,167	48,196	77,379	137,168	194,669	247,840	244,186	259,915
Deaths	543,700	555,203	560,241	567,304	587,488	653,000	634,432	746,146	709,035	713,499
Mortal remains	1,000	1,750	1,779	9,880	18,899	34,178	38,305	29,266	45,959	45,986
% of cremation	1.07	2.78	5.38	8.50	13.17	21.01	30.68	33.22	34.44	36.43
Crematoria	ND	31	35	43	53	70	85	87	89	91

Table 7.8 Emission factors for cremation of corpses - 2022

Air pollutant	u.m.	Cremation
NO _x	kg/body	0.4739
CO	kg/body	0.0047
NMVOC	kg/body	0.0091
SO _x	kg/body	0.0093
PM10	g/body	2.1483
PM2.5	g/body	2.1483
Pb	mg/body	30.0300
Cd	mg/body	5.0300
Hg	mg/body	0.0070
As	mg/body	13.6100
Cr	mg/body	13.5600
Cu	mg/body	12.4300
Ni	mg/body	17.3300
PAH (benzo(a)pyrene)	µg/body	176.2000

Air pollutant	u.m.	Cremation
Dioxins	µg/body	0.0190

7.2.5 Small scale waste burning (5C2)

The open burning of agricultural waste is a key category for Cd and CO emissions. NMVOC, CO, NO_x, NH₃, SO₂, PM10, PM2.5, BC, As, Cr, Cu, Ni, Se, Zn, Pb, Cd, Hg, Dioxin and PAH emissions have been estimated, applying the tier 1 and tier 2 approach (for heavy metals, PAH emissions and BC). No estimations were performed for NH₃ emissions as well as other POPs. In the current submission EFs used for field burning, open burning of agricultural waste and fires have been made coherent and updated on the basis of the 2023 EMEP/EEA Guidebook and the 2006 IPCC Guidelines. This survey regarded CO, CH₄, NMVOC, NO_x, NH₃, N₂O, SO₂, PM, dioxins and PAH.

A country-specific methodology has been used. Parameters taken into consideration are the following:

1. Amount of removable residues (t), estimated with annual crop production (ISTAT, several years [a], [b]; ISTAT, 2017 [a], [b]) and removable residues/product ratio (IPCC, 1997; CESTAAT, 1988; Borgioli E., 1981).
2. Amount of dry residues in removable residue (t dry matter), calculated with amount of removable fixed residues and fraction of dry matter (IPCC, 1997; CESTAAT, 1988; Borgioli E., 1981).
3. Amount of removable dry residues oxidized (t dry matter), assessed with amount of dry residues in the removable residues, burnt fraction of removable residues (CESTAAT, 1988) and fraction of residues oxidized during burning (IPCC, 1997).
4. Amount of carbon from removable residues burning release in air (t C), calculated with the amount of removable dry residue oxidized and the fraction of carbon from the dry matter of residues (IPCC, 1997; CESTAAT, 1988).
5. C-CH₄ from removable residues burning (t C-CH₄), calculated with the amount of carbon from removable residues burning release in air and emission factor equal to 4.7g/kg dry matter burnt (IPCC, 2006).
6. C-CO from removable residues burning (t C-CO), calculated with the amount of carbon from removable residues burning release in air and emission factor equal to 107g/kg dry matter burnt (IPCC, 2006).
7. Amount of nitrogen from removable residues burning release in air (t N), calculated with the amount of removable dry residue oxidized and the fraction of nitrogen from the dry matter of residues. The fraction of nitrogen has been calculated considering raw protein content from residues (dry matter fraction) divided by 6.25.
8. N-NO_x from removable residues burning (t N-NO_x), calculated with the amount of nitrogen from removable residues burning release in air and emission factor equal to 3 g/kg dry matter burnt (IPCC, 2006).

NMVOC emissions have been considered equal to CH₄ emissions. As regards the other pollutants, heavy metals, Dioxin and PAH emission factors are from the EMEP/EEA Guidebook (EMEP/EEA, 2016) and emissions have been added as requested by the NECD review process (EEA, 2018) (Table 7.9).

Table 7.9 Emission factors for burning of agriculture residues

Air pollutant	u.m.	Removable residues				References
		Wheat	Barley	Maize	Rice	
Benzo(a)pyrene	g/t	0.393	0.771	7.162	0.072	

Air pollutant	u.m.	Removable residues				References
		Wheat	Barley	Maize	Rice	
Benzo(b)fluoranthene	g/t	1.097	2.398	3.495	0.120	EMEP/EEA, 2023
Benzo(k)fluoranthene	g/t	0.468	0.601	2.138	0.088	
Indeno(1,2,3-cd)pyrene	g/t	0.336	0.298	2.415	0.055	
PM10	g/t		3.3			EMEP/EEA, 2023
PM2.5	g/t		2.8			EMEP/EEA, 2023
Dioxins	g/t		10			EMEP/EEA, 2023
BC	g/t		1.2			EMEP/EEA, 2013

Removable residues from agriculture production are estimated for each crop type (cereal, green crop, permanent cultivation) taking into account the amount of crop produced, from national statistics (ISTAT, several years [a], [b]; ISTAT, 2017 [a], [b]), the ratio of removable residue in the crop, the dry matter content of removable residue, the ratio of removable residue burned, the fraction of residues oxidised in burning, the carbon and nitrogen content of the residues. Most of these wastes refer especially to the prunes of olives and wine, because of the typical national cultivation. Activity data (agricultural production) used for estimating burning of agriculture residues are reported in Table 7.10. Emissions due to stubble burning, which are emissions only from the agriculture residues burned on field, are reported in the agriculture sector, under 3.F. Under the waste sector the burning of removable agriculture residues that are collected and could be managed in different ways (disposed in landfills, used to produce compost or used to produce energy) is reported. Different percentages of the removable agriculture residue burnt for different residues are assumed, varying from 10% to 90%, according to national and international literature. Moreover, these removable wastes are assumed to be all burned in open air (e.g. on field), taking in consideration the highest available CO, NMVOC, PM and dioxins emission factors as reported in the table above. The amount of biomass from pruning used for domestic heating is reported in the energy sector in the 1A4b category as biomass fuel.

Table 7.10 Time series of crop productions (Gg)


Production	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Cereals										
Wheat	8,108.5	7,946.1	7,427.7	7,717.1	6,849.9	7,394.5	6,576.6	6,553.9	7,118.3	6,449.8
Rye	20.8	19.8	10.3	7.9	13.9	13.2	12.5	11.5	10.9	11.4
Barley	1,702.5	1,387.1	1,261.6	1,214.1	944.3	955.1	1,072.4	1,090.6	1,059.8	1,124.3
Oats	298.4	301.3	317.9	429.2	288.9	261.4	238.1	242.7	233.5	242.3
Rice	1,290.7	1,320.9	1,245.6	1,444.8	1,574.3	1,505.8	1,505.1	1,530.9	1,496.5	1,237.0
Maize	5,864	8,454	10,140	10,428	8,496	7,074	6,258.7	6,771.1	6,060.2	4,681.9
Sorghum	114.2	214.8	215.2	184.9	275.6	294.2	312.4	361.7	223.5	191.2
Woodycrops										
Grapes	8,438.0	8,447.7	8,869.5	8,553.6	7,839.7	7,915.0	7,862.9	8,222.4	8,149.4	8,438.0

Production	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Gg										
Olives	912.5	3,323.5	2,810.3	3,774.8	3,117.8	2,732.9	2,194.1	2,207.2	2,270.6	2,453.3
CitrusOrchards	2,868.8	2,607.7	3,100.2	3,518.1	3,820.6	3,151.5	2,895.9	2,940.0	3,098.0	3,094.4
Orchards	5,793.5	5,406.6	5,952.2	6,034.5	5,777.3	5,988.8	5,318.0	5,408.7	4,602.1	5,301.7
Carobs	29.2	44.4	38.1	31.7	25.3	31.5	35.9	36.9	37.6	35.6
Total	35,441	39,474	41,388	43,339	39,023	37,318	34,283	35,377	34,360	33,261

7.2.6 Wastewater treatments (5D)

Under source category 5D, NMVOC emissions are estimated both from domestic and industrial wastewater and NH₃ emissions are estimated from latrines. In Italy, domestic wastewaters follow the treatment systems and discharge pathways reported in the following figure. Commercial and some industrial wastewaters assimilated to municipal flows are co-discharged with domestics.

Figure 7.1 Domestic wastewater treatment system and discharge pathways



In urban areas, domestic wastewaters are collected and treated in plants mainly using a secondary treatment, with aerobic biological units: a wastewater treatment plant standard design consists of bar racks, grit chamber, primary sedimentation, aeration tanks (with return sludge), settling tank, chlorine contact chamber. The stabilization of sludge occurs in aerobic or anaerobic reactors; where anaerobic digestion is used, the reactors are covered and provided of gas recovery. The biogas collected from the anaerobic digestion of wastewaters is burned with heat/energy recovery and relevant emissions are

reported in the energy sector. On the contrary, in rural areas, wastewaters are treated in Imhoff tanks or in other on-site systems, such as latrines.

Industrial wastewaters are produced from most production industries like food and beverage, cloth and shoe production, paper and heavy production sites such as mining, oil and gas, as well as chemical manufacturing. Industrial wastewaters have very variable quality and volume depending on the type of industry producing and cannot be assimilated to domestic wastewaters, thus they are treated in specific plants, usually located within the industrial area.

By using EFs from the 2023 EEA/EMEP Guidebook both for domestic and industrial wastewater and the volumes of wastewater handled, NMVOC emissions resulted in the time series reported in the table below (Table 7.11). Volume of domestic wastewater is calculated starting from the inhabitants' equivalent treated in wastewater treatment plants multiplied by the average per capita water supply. For industrial wastewater, data have been collected for several industrial sectors (iron and steel, refineries, organic chemicals, food and beverage, paper and pulp, textiles and leather industry). The volume of wastewater production, for each industry selected, has been calculated multiplying the annual production ($t\ year^{-1}$) by the amount of wastewater consumption per unit of product ($m^3\ t^{-1}$), as reported in following box.

	Wastewater generation (m^3/t)	References
Coke	1.5	IPCC, 2000
Organic Chemicals	22.3	FEDERCHIMICA, several years
Paints	5.5	IPCC, 2000
Plastics and Resins	0.6	IPCC, 2000
Soap and Detergents	3	IPCC, 2000
Vegetables, Fruits and Juices	20	IPCC, 2000
Sugar Refining	4	ANPA-ONR, 2001
Vegetable Oils	3.1	IPCC, 2000
Dairy Products	3.87	ANPA-ONR, 2001
Wine and Vinegar	3.8	ANPA-ONR, 2001
Beer and Malt	420 (l/hl)	Assobirra, several years
Alcohol Refining	24	IPCC, 2000
Meat and Poultry	13	IPCC, 2000
Fish Processing	13	same value of Meat and Poultry
Paper	25	Assocarta, several years
Pulp	25	Assocarta, several years
Textiles (dyeing)	60	IPCC, 1995
Textiles (bleaching)	350	IPCC, 1995
Leather	0.10	UNIC, several years

In Table 7.11 NH_3 emissions are reported from domestic wastewater. An in-depth analysis of national circumstances has been made, collecting many statistical data on population and on urban wastewater treatment plants (BLUE BOOK, several years; COVIRI, several years; ISTAT, 1984; ISTAT, 1987; ISTAT, 1991; ISTAT, 1993; ISTAT [a], [b], 1998; ISTAT [d], [e], several years). Some data, such as the degree of collected or treated wastewater are available for specific year, so the entire time series has been reconstructed with interpolation of data. In 2022, the 99.6% of population is served by sewer systems, whereas 91% of population is served by wastewater treatment plants. In 1990, the percentage of population served by sewer system was 57%, whereas only 52% of population was served by wastewater treatment plants.

Rural population, for national circumstances, is intended as the population in "dispersed houses". Within this category (population living in "dispersed houses"), those covered by a local sewer systems, are treated only in Imhoff systems and latrines, not in WWTPs, thus activity data for calculation of NH_3 emissions are derived from the difference of rural population collected, minus rural population treated in Imhoff tanks: it is assumed that all the wastewater treated in other systems than the Imhoff is treated in latrines, consistently with the approach used to estimate GHG.

Table 7.11 Time series of NMVOC and NH₃ emissions (Gg)

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
NMVOC										
Domestic wastewater										
Equivalent inhabitants (*1000)	46,436	60,015	65,601	73,426	76,847	75,239	89,716	91,124	92,532	93,940
Domestic wastewaters production (10 ⁶ m ³)	4,237	5,476	5,986	6,700	7,012	6,866	8,187	8,315	8,444	8,572
Per capita water supply (lt./person*die)	250	250	250	250	250	250	250	250	250	250
EF (mg/m ³)	15	15	15	15	15	15	15	15	15	15
NMVOC Emissions (t)	63.6	82.1	89.8	100.5	105.2	103.0	122.8	124.7	126.7	128.6
Industrial wastewater										
Industrial wastewaters production (1000 m ³)	908,840	928,479	920,614	867,085	717,846	659,246	696,915	654,586	701,388	675,917
EF (mg/m ³)	15	15	15	15	15	15	15	15	15	15
NMVOC Emissions (t)	13.6	13.9	13.8	13.0	10.8	9.9	10.5	9.8	10.5	10.1
NH₃										
Domestic wastewater										
Rural population (*1000)	3,831	3,710	3,589	3,422	3,347	3,225	3,129	3,104	3,080	3,056
Population not served by collecting systems	1,647	1,120	502	582	330	19	16	15	14	13
Population served by collecting systems	2,184	2,590	3,087	2,840	3,016	3,206	3,113	3,090	3,066	3,043
treated in Imhoff tanks	421	647	845	468	635	967	1,156	1,174	1,192	1,210
treated in latrines	1,763	1,943	2,242	2,373	2,381	2,239	1,957	1,915	1,874	1,832
EF (kg/person/year)	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6	1.6
NH₃ Emissions (t)	2,821	3,109	3,587	3,796	3,810	3,583	3,131	3,065	2,998	2,932

7.2.7 Other waste (5E)

On the basis of the Final review report of the 2017 Comprehensive technical review of national emission inventories (EEA, 2017 [a]) emissions from category 5E – Car and Building Fires have been estimated. Buildings have been subdivided into 4 subcategories: detached house, undetached house, apartment buildings and industrial buildings and the distribution of population in the different typology of building has been derived from Eurostat. Data regarding the number of car and building fires have been derived from the Annually statistics of fire service in Italy (Annually statistics of fire service in Italy, several years) while EFs are coherent with the Guidebook EMEP/EEA 2016 deriving from Aasestad, 2007 for particulate matter (TSP=PM10=PM2.5) while BC EF has been derived from IIASA report (IIASA, 2004). No data about car and building fires are available before 2000 so 90's data have been reconstructed on the basis of the national population and the resulting time series are reported in Table 7.12. On the basis of the last review reports (EEA, 2019; EEA, 2020; EEA 2021) PCDD/F emissions have been estimated and revised. After the

technical correction during the 2019 NECD review, Italy investigated emission factors because those reported in the guidebook seems to be unreliable or not fitting to the Italian context. In particular, the emission factors reported in the EMEP/EEA Guidebook “used for particles in the inventory are given by scaling the emission factors used for combustion of fuelwood in the households” (Aasestad, 2007) but the Italian buildings are made up of the vast majority of reinforced concrete. Despite this, following the last review process, Italy decided to use these emission factors pending the availability of values deriving from new studies. Further, Cd, Hg and Pb emissions have been estimated for this source on the basis of 2019 EMEP/EEA Guidebook.

Table 7.12 PM2.5 and BC emissions from the category 5E

	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Detached house fires (n°)	10,580	10,592	10,614	10,861	9,213	9,577	13,222	13,554	14,174	13,151
Undetached house fires (n°)	8,186	8,195	8,212	8,404	6,928	11,484	12,728	12,078	12,685	12,378
Apartment building fires (n°)	18,843	18,865	18,904	19,344	18,891	23,278	28,913	29,020	32,695	33,980
Industrial building fires (n°)	4,931	4,936	4,947	5,062	4,560	4,872	3,599	3,192	3,278	4,128
Car fires (n°)	25,924	25,954	26,008	26,614	22,735	22,680	22,170	18,641	19,270	19,901
5E PM2.5 (Gg)	3.04	3.05	3.05	3.13	2.76	3.29	4.10	4.09	4.39	4.30
5E BC (Gg)	0.56	0.56	0.57	0.58	0.51	0.61	0.76	0.76	0.81	0.80
5E PCDD/F (g Iteq)	31.18	31.21	31.28	32.01	28.20	33.56	41.69	41.53	44.47	43.63
Cd (t)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Hg (t)	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03
Pb (t)	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03

7.3 TIME SERIES AND KEY CATEGORIES

The following Table 7.13 presents an outline of the weight of the different categories for each pollutant in the waste sector for the year 2022. Key categories are those shaded in blue.

Table 7.13 Key categories in the waste sector in 2022

	5A	5B1	5B2	5C1bi	5C1biii	5C1bi v	5C1bv	5C2	5D1	5D2	5E
	%										
SO _x				0.06	0.001	0.04	0.003	0.08			
NO _x				0.01	0.005	0.01	0.02	0.27			
NH ₃	1.69	0.04	0.96								
NMVOC	0.89	0.04		0.04	0.004	0.001	0.0003	0.02	0.02	0.001	
CO				0.001	0.0002	0.001	0.0001	3.20			
PM10	0.001			0.0001	0.0006	0.000	0.0003	0.66			2.09
PM2.5	0.0002			0.0001	0.0009	0.000	0.0004	0.89			2.98
BC				0.00003	0.0003	0.00004		4.31			4.38
Pb				0.003	0.0006	0.004	0.004	0.07			0.01
Cd				0.01	0.0014	0.38	0.04	6.97			0.31
Hg				0.004	0.03	0.40	0.038	0.91			0.45
PAH				0.0001	0.00001	0.00004	0.00008	1.53			
Dioxins				0.01	0.008	0.003	0.002	2.17			14.35
HCB				0.06	7.69	0.73	0.07				
PCB				0.02	0.93	0.08	0.02				

Note: key categories are shaded in blue

The following pie charts show, for the main pollutants, the contribution of each sub-category to the total emissions from the waste sector, both for 1990 and 2022 (Figure 7.2, Figure 7.3, Figure 7.4 and Figure 7.5).

Finally, in Table 7.14 Time series emissions in the waste sector by category and pollutant, emissions time series for each pollutant of the waste sector are reported. In the period 1990-2022, total emissions from incineration plants increase, but whereas emissions from plants with energy recovery show a strong growth, emissions from plants without energy recovery decreased because of the legal constraints which impose the energy production. For 2022, about 99% of the total amount of urban waste incinerated is treated in plants with energy recovery system reported in 1A4a.

On the basis the NEC review (EEA, 2022) Italy took in consideration the review of PM emissions from landfills and made a survey on possible activity data and, above all, emission factors. In the current submission estimates have been carried out.

Figure 7.2 Contribution of CO, NOx and NMVOC sub-category emissions to waste sector total emissions




Figure 7.3 Contribution of POPs Annex III sub-category emissions to waste sector total emissions




Figure 7.4 Contribution of priority heavy metals sub-category emissions to waste sector total emissions




Figure 7.5 Contribution of PM10 and PM2.5 sub-category emissions to waste sector total emissions




Table 7.14 Time series emissions in the waste sector by category and pollutant

WASTE SECTOR	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Solid waste disposal (5A)										
NMVOOC (Gg)	6.43	7.97	9.06	8.96	8.20	7.39	7.08	7.51	7.38	7.32
NH ₃ (Gg)	5.21	6.46	7.35	7.26	6.65	5.99	5.74	6.09	5.98	5.94
PM10 (Gg)	0.004	0.004	0.004	0.003	0.003	0.002	0.002	0.002	0.002	0.002
PM2.5 (Gg)	0.0006	0.0006	0.0006	0.0005	0.0004	0.0003	0.0003	0.0003	0.0003	0.0003
Biological treatment of waste (5B)										
NMVOOC (Gg)	0.01	0.03	0.14	0.28	0.36	0.37	0.36	0.35	0.35	0.33
NH ₃ (Gg)	0.07	0.20	0.70	0.95	0.41	2.66	3.25	3.44	3.58	3.51
Waste incineration (5C)										
CO (Gg)	61.69	67.73	65.89	72.48	68.53	68.57	65.10	66.56	65.97	60.64
NO _x (Gg)	2.70	2.69	2.27	2.44	2.20	2.11	2.01	2.02	2.10	1.96
NMVOOC (Gg)	1.99	1.93	1.01	1.42	0.98	0.47	0.43	0.40	0.51	0.52
SO _x (Gg)	0.55	0.49	0.30	0.34	0.23	0.17	0.15	0.12	0.17	0.16
NH ₃ (Gg)	0.0012	0.0008	0.0005	0.0002	0.0002	-	-	-	-	-
PM10 (Gg)	1.78	2.04	1.97	1.98	1.52	1.44	1.31	1.34	1.35	1.36
PM2.5 (Gg)	1.66	1.91	1.85	1.86	1.44	1.36	1.24	1.27	1.28	1.29
BC (Gg)	0.80	0.87	0.85	0.94	0.88	0.88	0.84	0.86	0.85	0.78
PAH (t)	1.27	1.23	1.16	1.21	0.98	1.04	1.00	1.01	1.01	0.96
Dioxins (g I-Teq))	103.15	79.52	28.60	8.04	7.52	7.50	7.12	7.27	7.23	6.65
HCB (kg)	12.86	13.97	9.87	8.27	0.49	0.98	0.92	0.85	1.07	1.07
PCB (kg)	5.36	4.62	2.08	1.55	0.57	1.03	0.98	0.91	1.14	1.14
As (t)	0.09	0.09	0.07	0.07	0.05	0.05	0.05	0.05	0.05	0.04
Cd (t)	0.59	0.57	0.46	0.50	0.32	0.35	0.32	0.31	0.33	0.30
Cr (t)	0.66	0.58	0.35	0.39	0.09	0.10	0.09	0.08	0.09	0.09
Cu (t)	0.99	0.86	0.55	0.49	0.11	0.16	0.13	0.10	0.15	0.14
Hg (t)	0.33	0.32	0.23	0.30	0.20	0.09	0.07	0.06	0.08	0.08
Ni (t)	6.79	4.37	2.84	1.05	0.04	0.05	0.04	0.04	0.05	0.04
Pb (t)	5.90	5.53	2.76	4.03	0.24	0.18	0.16	0.16	0.17	0.17
Se (t)	0.04	0.04	0.03	0.04	0.03	0.04	0.04	0.04	0.04	0.03
Zn (t)	5.28	6.32	4.73	5.95	3.60	3.41	3.08	3.18	3.19	3.31

WASTE SECTOR	1990	1995	2000	2005	2010	2015	2019	2020	2021	2022
Wastewater (5D)										
NMVOC (Gg)	0.08	0.10	0.10	0.11	0.12	0.11	0.13	0.13	0.14	0.14
NH ₃ (Gg)	2.82	3.11	3.59	3.80	3.81	3.58	3.13	3.06	3.00	2.93
Other waste (5E)										
PM2.5 (Gg)	3.04	3.05	3.05	3.13	2.76	3.29	4.10	4.09	4.39	4.30
BC (Gg)	0.56	0.56	0.57	0.58	0.51	0.61	0.76	0.76	0.81	0.80
PCDD/F (g I-Teq)	31.18	31.21	31.28	32.01	28.20	33.56	41.69	41.53	44.47	43.63
Cd (t)	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Hg (t)	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03
Pb (t)	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.02	0.03	0.03

7.4 RECALCULATIONS

In the following table the recalculations occurred in the 2024 submission with respect the last year submission are reported at category level.

WASTE SECTOR	1990	1995	2000	2005	2010	2015	2019	2020	2021
Solid waste disposal (5A)									
NMVOC	-	-	-	-	-	-	0.04%	0.05%	0.1%
NH ₃	-	-	-	-	-	-	0.04%	0.05%	0.1%
PM10	-3.0%	-27.2%	-31.8%	-26.2%	-27.9%	-5.1%	10.3%	9.0%	20.7%
PM2.5	-3.0%	-27.2%	-31.8%	-26.2%	-27.9%	-5.1%	10.3%	9.0%	20.7%
Biological treatment of waste (5B)									
NMVOC	-	-	-	-	-	-	-	-	-
NH ₃	900.0%	200.7%	616.0%	197.0%	-11.7%	-2.5%	6.8%	9.8%	10.0%
Waste incineration (5C)									
CO	54.7%	53.9%	54.1%	53.7%	54.4%	54.9%	55.6%	56.1%	56.2%
NO _x	0.6%	-4.4%	-3.8%	-5.2%	-5.9%	-4.7%	-4.3%	-4.9%	-1.2%
NMVOC	-56.0%	-58.2%	-67.8%	-60.9%	-68.7%	-83.1%	-83.8%	-85.1%	-80.4%
SO _x	-0.2%	-0.4%	-0.6%	-0.4%	-1.0%	-1.4%	-1.7%	-2.2%	12.3%
PM10	-	-	-	-	-	NA	NA	NA	NA
PM2.5	-15.1%	-11.7%	-11.5%	-18.8%	-33.5%	-37.2%	-40.2%	-39.9%	-37.7%
BC	-1.1%	-2.1%	-2.2%	-1.7%	-2.8%	-2.9%	-3.1%	-3.3%	-1.4%
PAH	-1.5%	-3.2%	-3.4%	-2.7%	-5.2%	-5.2%	-5.7%	-6.4%	-3.7%

WASTE SECTOR	1990	1995	2000	2005	2010	2015	2019	2020	2021
Dioxins	-0.1%	-0.2%	-0.6%	-1.6%	-2.8%	-2.9%	-3.1%	-3.3%	-1.2%
HCB	-	-	-	-	-	-	-	-	18.1%
PCB	-	-	-	-	-	-	-	-	18.4%
As	-0.1%	-0.1%	-0.2%	-0.1%	-0.3%	-0.3%	-0.02%	0.5%	1.4%
Cd	-1.2%	-2.7%	-3.3%	-2.6%	-6.1%	-5.8%	-6.8%	-7.9%	-4.2%
Cr	-0.1%	-0.2%	-0.4%	-0.3%	-2.0%	-2.0%	-2.2%	-2.5%	2.0%
Cu	-0.1%	-0.2%	-0.2%	-0.2%	-1.5%	-1.1%	-1.4%	-1.8%	4.3%
Hg	-0.4%	-0.8%	-1.1%	-0.7%	-1.6%	-3.7%	-4.8%	-6.7%	-2.3%
Ni	0.0%	0.0%	0.0%	-0.1%	-3.2%	-2.7%	-3.0%	-3.7%	-0.3%
Pb	0.0%	0.0%	-0.1%	0.0%	-1.1%	-1.5%	-1.7%	-1.9%	0.5%
Se	-0.4%	-0.9%	-1.1%	-0.8%	-1.3%	-1.4%	-1.2%	-1.1%	0.9%
Zn	-0.1%	-0.2%	-0.2%	-0.1%	-0.4%	-0.4%	-0.4%	-0.4%	0.5%
Wastewater (5D)									
NMVOG	-	-	-	-	-	-	-0.001%	0.004%	-0.02%
NH3	NA	NA	NA	NA	NA	NA	NA	NA	NA
Other waste (5E)									
PM2.5	-	-	-	-	-	-	-	-	22.4%
BC	-	-	-	-	-	-	-	-	22.4%
PCDD/F	-	-	-	-	-	-	-	-	22.2%
Cd	-	-	-	-	-	-	-	-	23.0%
Hg	-	-	-	-	-	-	-	-	22.9%
Pb	-	-	-	-	-	-	-	-	22.9%

In general recalculations occur in 2021 because of the update of activity data of industrial and clinical waste reported by ISPRA (ISPRA, several years) in December 2023.

As regards waste disposed of in landfills, minor recalculations occur for NMVOC and NH₃ because of a fixed error in the model for 2016. Minor recalculations start from 2017 because of the delay time. Recalculations occur also for PM because of the review process (EEA, 2023) and the subsequential estimates of PM emissions based on mineral waste data reported. PM emissions reduce because in the previous submission have been estimated on the basis of the total amount of waste in landfills while now only the inert part has been considered.

As regards NH₃ from 5B, recalculations occur because of the change in the coefficients of N contained in livestock manure sent to anaerobic digestion of cattle, swine, poultry. More information is reported in the NIR (Agriculture chapter).

As regards emissions from 5C2 open burning of agricultural residues emissions, the recalculation of all emissions is due to updating of emission factors for CO, NO_x, NMVOC, PM₁₀, PM_{2.5}, substituted with those in the 2023 EMEP/EEA Guidebook and in the 2006 IPCC guidelines (for CO, NO_x).

As regards waste incineration, on the basis of (EEA, 2023) PM2.5/PM10 emissions from 5c1bi (Industrial waste incineration) and NMVOC emissions from 5C1biii (Clinical waste incineration) implementing T1 EF from 2019 EMEP/EEA Guidebook.

As regards Domestic wastewater handling the review process has led to NH₃ estimates from dry toilets.

7.5 PLANNED IMPROVEMENTS

Recently, in the previous submissions, estimates of PM emissions have been included in the inventory for solid waste disposed on landfills. Starting with the last NEC review, Italy is studying the most appropriate way to consider the contribution of inert materials to PM emissions, not only to ensure that the best available data is used but also to integrate the estimation system within the landfill emissions model.

As regards composting, Italy is looking for further information to apply the penetration of technologies back to 1990 even if from the first analyzes it emerges that the Italian technological context was among the most advanced even in those years. At the moment Italy is looking for further information to satisfy the request of the review process (EEA, 2023) and be able to introduce the penetration rate of abatement technologies with particular focus on 90s.

Emissions from 5E are still under investigation, the results will be reported in the next submissions.

8 OTHER (6A)

NH₃ emissions from dog and cat droppings are reported in this category. Emissions have been estimated applying the tier 1 approach and NH₃ emission factors for cats and dogs from the EMEP/EEA Guidebook (EMEP/EEA, 2023).

The historical series of the number of cats has been estimated starting from the data of FEDIAF (European Pet Food Industry Federation) report 2021 and 2022 (FEDIAF, 2022; 2023). The number of cats per household has been calculated for the years 2021-2022 as the number of cats divided by the number of households (source ISTAT- <http://dati.istat.it/>). The number of cats for the period 1990-2020 has been estimated as the product of the number of households of a certain year (source ISTAT/EUROSTAT) and the average percentage 2021-2022 number of cats per household.


The historical series of the number of dogs has been estimated starting from the data of the Canine Registry managed by the Ministry of Health, collecting data of regional registries (<https://www.salute.gov.it/portale/caniGatti/menuContenutoCaniGatti.jsp?lingua=italiano&area=cani&menu=anagrafe>). The updated number of dogs referred to present (2023) has been considered as constant for the year 2022. The number of dogs per household for the year 2022 has been estimated as the number of dogs divided by the number of households. The number of dogs for the period 1990-2020 has been estimated as the product of number of households of a certain year (source ISTAT/EUROSTAT) and the number of dogs per household for the year 2022.

8.1 TIME SERIES AND KEY CATEGORIES

In 2022, key category level was identified for 6A NH₃ emissions and for trend analysis; on the other hand, in 1990 they are not.

In Figure 8.1 NH₃ emissions trend are reported.

Figure 8.1 NH₃ emissions from pets (dogs and cats)



8.2 QA/QC AND VERIFICATIONS

Other data sources were consulted to verify activity data on dogs bred by Italian families, such as FEDIAF, ENCI (*Ente Nazionale della Cinofilia Italiana* – National board of Italian dog breeding) and FCI (Federation Cynologique Internationale).

8.3 RECALCULATIONS

This emission category was processed and included in the 2024 submission for the first time. There is no recalculation.

8.4 PLANNED IMPROVEMENTS

No improvement is planned.

9 RECALCULATIONS AND IMPROVEMENTS

9.1 RECALCULATIONS

To meet the requirements of transparency, consistency, comparability, completeness and accuracy of the inventory, the entire time series is checked and revised every year during the annual compilation of the inventory. Measures to guarantee and improve these qualifications are undertaken and recalculations should be considered as a contribution to the overall improvement of the inventory.

Recalculations are elaborated on account of changes in the methodologies used to carry out emission estimates, changes due to different allocation of emissions as compared to previous submissions, changes due to error corrections and in consideration of new available information.

The complete NFR files from 1990 to 2022 have been submitted. The percentage difference between the time series reported in the 2023 submission and the series reported this year (2024 submission) are shown in Table 9.1 by pollutant. Further information is available in the relevant chapter.

Improvements in the calculation of emission estimates have led to a recalculation of the entire time series of the national inventory. Considering the total emissions, the emission levels for the year 2021 show a decrease for some pollutants and an increase for others; in particular, a significant decrease has been observed for NMVOC emissions, equal to -2.4%, but on the other side a significant increase of emissions has been observed in 2021, according to the review process, for NH₃ (11.7%) because of recalculations occurred in the agriculture sector.

For 1A1, recalculations occurred for CO, NMVOC, PM_{2.5}, BC, Cd, Hg, Pb, PAH and dioxins because of the review of heavy residual in refineries activity data in 2010 and in the period 2013-2020. Recalculations occurred in 2021 for PM and BC from coke production because of the correction of an error in PM emission factor.

For 1A2, recalculations occurred for NH₃ emissions because of the review process (EEA, 2023) and the relevant request to assess NH₃ emissions from 1A2e biomass. Recalculations also occurred for 1A2b non ferrous metal starting from 2014, because of the analysis of environmental authorization (IPPC permits) of the sole integrated plant of zinc/lead production in Italy. For 1A2f changes occurred for NO_x, SO_x and NH₃ because the review of the series between 2007 and 2021.

In 2024 submission the historical series has been revised mainly as a result of the upgrade of COPERT model version used (from version 5.6.1 in last submission to 5.7.3 in submission 2024 (EMISIA, 2024)): Emission factors of Euro 6 CNG passenger cars, Euro VI diesel buses, Euro VI diesel hybrid buses, non-exhaust emission factors have been updated. As regards the software, revisions relate: the removal of CO₂ correction tool, capability of alternative HDVs classification based on REG EU 2017/2400, improved labels of forms and headers. Furthermore various bugs have been corrected.

For 1.A.3.dii national navigation, natural gas emission factor has been corrected for year 2021. Minor corrections on the figures of the gasoline engines split between 2 stroke and 4 stroke from 2014 onwards. The corrections lead to any relevant impacts on the distribution of the ratio of 2 strokes 4 strokes in the time series.

Energy recovery from waste reported in the commercial heating (1A4ai) has been updated from 2013 because of the update of data single plant leading to minor recalculations. Thanks to the NEC review process an error has been identified in NMVOC EF from clinical waste and it has been updated. More, the ratio PM_{2.5}/PM₁₀ has been updated for industrial waste from 2006. Recalculations occurred for the whole time series because of the update of the energy balance, in particular the update of kerosene time series which has been added.

In the industrial processes sector recalculations occur for PM₁₀, PM_{2.5} along the whole time series because this submission includes PM emissions estimates from category 2.A.5.a quarrying and mining; revised PM emission estimates from category 2.A.5b construction and demolition of buildings and PM emission estimates from construction and demolition of roads. Recalculations occur for PM emissions

from 2A5b due to the correction of reporting mistakes along the timeseries form last year submission and for NMVOC emissions from lubricants use because of the update of the activity data for 2012-2021.

For the solvent sector recalculations occurred for the update of EFs in paint application in construction and buildings for the years 2019-2021. In chemical product processing and manufacturing, a recalculation occurred on account of a revision in activity data for polyester, from 2018, and polyurethane processing, from 2008, and a change in EFs for leather tanning from 2019 due to information available from the industrial association reports.

Recalculations were implemented for the agricultural emission inventory. Major recalculations of NH₃ emissions are due to: update of NH₃ emission factors from synthetic fertilizers, reported in the EMEP/EEA Guidebook 2023; change in the coefficients of N contained in livestock manure sent to anaerobic digestion of cattle, swine, poultry since 1991; correction of NH₃ EFs from storage for cattle in the years 2003-2005; correction of NH₃ EFs from storage for cattle and pigs since 2014; correction of N values of synthetics for the year 2016, N values of soil improvers for the years 2015 and 2016, N values of other organics for the year 2016. The estimate of NH₃ from the residues was added. More info in the relevant chapter. As regards field burning of agricultural residues emissions, the recalculation of all emissions is due to: updating of emission factors for CO, NO_x, NMVOC, PM₁₀, PM_{2.5}, substituted with those in the 2023 EMEP/EEA Guidebook and in the 2006 IPCC guidelines (for CO, NO_x); the estimate of sorghum residue since 1990 has been changed, now aligned with the estimate of emissions from agricultural soils; the percentage of rice straw burned since 2001 has been changed, and the production of the years 2019-2021 has been updated.

As regards the waste sector, in general recalculations occur in 2021 because of the update of activity data of industrial and clinical waste reported by ISPRA (ISPRA, several years) in December 2023. In particular, as regards waste disposed of in landfills, minor recalculations occur for NMVOC and NH₃ because of a fixed error in the model for 2016. Minor recalculations start from 2017 because of the delay time. As regards Domestic wastewater handling the review process has led to NH₃ estimates from dry toilets. As regards emissions from 5C2 open burning of agricultural residues emissions, the recalculation of all emissions is due to: updating of emission factors for CO, NO_x, NMVOC, PM₁₀, PM_{2.5}, substituted with those in the 2023 EMEP/EEA Guidebook and in the 2006 IPCC guidelines (for CO, NO_x).

Table 9.1 Recalculation between 2023 and 2024 submissions (%)

	SO _x	NO _x	NH ₃	NM VOC	CO	PM 10	PM 2.5	BC	Pb	Hg	Cd	DIO X	PAH	HCB	PCB
	%														
1990	0.01	0.02	12.91	-0.56	0.43	0.53	-0.86	-0.10	-0.04	-0.01	-0.15	0.02	-0.03	-	-
1991	0.01	0.02	13.38	-0.54	0.48	-19.03	-4.72	-0.11	-0.05	-0.01	-0.17	0.02	-0.04	-	-
1992	0.01	0.02	14.25	-0.53	0.44	0.52	-0.88	-0.13	-0.08	-0.02	-0.21	0.01	-0.05	-	-
1993	0.01	0.01	14.70	-0.49	0.47	1.22	-0.78	-0.13	-0.09	-0.02	-0.26	-0.01	-0.06	-	-
1994	0.01	0.01	14.05	-0.48	0.68	0.02	-0.99	-0.14	-0.09	-0.02	-0.28	-0.01	-0.06	-	-
1995	0.01	0.01	13.31	-0.50	0.73	-0.42	-1.01	-0.14	-0.10	-0.02	-0.27	-0.01	-0.06	-	-
1996	0.01	0.01	13.22	-0.51	1.01	0.01	-1.08	-0.14	-0.11	-0.02	-0.26	-0.01	-0.06	-	-
1997	0.01	0.00	13.99	-0.55	1.09	1.54	-0.86	-0.14	-0.12	-0.02	-0.23	-0.01	-0.04	-	-
1998	0.01	0.01	13.08	-0.59	1.34	-1.77	-1.35	-0.13	-0.13	-0.02	-0.23	-0.00	-0.04	-	-
1999	0.01	-0.00	13.07	-0.56	1.62	0.40	-1.02	-0.14	-0.14	-0.02	-0.27	-0.02	-0.06	-	-

	SO _x	NO _x	NH ₃	NM VOC	CO	PM 10	PM 2.5	BC	Pb	Hg	Cd	DIO X	PAH	HCB	PCB
	%														
2000	0.00	0.69	13.17	-0.52	1.83	1.41	-1.31	-1.29	-0.21	-0.02	-0.30	-0.02	-0.10	-0.00	-0.00
2001	0.01	-0.01	13.45	-0.64	2.01	0.04	-1.12	-0.16	-0.25	-0.02	-0.29	-0.03	-0.10	-	-
2002	0.01	-0.01	13.98	-0.70	2.56	-19.04	-5.61	-0.18	-0.64	-0.03	-0.36	-0.04	-0.14	-	-
2003	0.00	-0.02	13.88	-0.75	2.56	0.96	-1.15	-0.16	-0.59	-0.02	-0.31	-0.02	-0.08	-	-
2004	0.00	-0.02	14.36	-0.87	3.12	-4.85	-2.43	-0.18	-0.50	-0.03	-0.39	-0.05	-0.14	-	-
2005	0.00	0.07	14.21	-0.97	1.88	-3.50	-1.90	-0.04	-0.44	-0.03	-0.34	-0.03	-0.08	0.00	0.00
2006	0.04	0.11	14.31	-1.42	1.30	2.59	-1.00	0.10	-0.45	-0.04	-0.44	-0.03	-0.09	0.00	-0.00
2007	0.07	0.60	13.70	-1.33	1.43	-9.98	-2.75	0.57	-0.57	-0.04	-0.45	0.02	-0.04	0.00	0.00
2008	0.03	0.79	12.42	-1.32	0.84	-1.45	-1.31	0.74	-0.55	-0.05	-0.57	0.03	-0.03	0.00	0.00
2009	0.02	0.04	11.71	-1.25	0.90	-6.16	-2.17	0.35	-0.72	-0.08	-0.77	0.14	-0.01	0.00	0.00
2010	-1.10	1.03	11.94	-1.14	0.71	-11.92	-3.09	0.88	-0.72	-0.09	-0.99	0.04	-0.05	0.00	0.00
2011	0.03	1.16	12.24	-1.24	0.75	-16.12	-4.60	1.00	-0.68	-0.09	-1.07	0.03	-0.08	0.00	0.00
2012	0.02	1.23	13.43	-1.17	0.50	1.33	-1.47	0.90	-0.67	-0.06	-0.81	0.06	-0.01	0.00	0.00
2013	0.01	1.49	12.50	-1.14	0.53	-1.05	-1.73	1.02	-0.66	-0.10	-1.17	0.04	-0.04	0.00	-0.00
2014	-0.04	2.59	12.23	-1.04	0.99	0.32	-1.51	1.75	-0.65	-0.11	-1.34	0.08	-0.04	-0.00	-0.00
2015	-0.07	2.32	12.64	-0.89	1.02	-6.82	-2.21	2.74	-0.59	-0.11	-1.04	0.15	-0.01	0.00	0.00
2016	-0.05	2.02	14.57	-2.16	0.02	-4.50	-2.34	0.87	-0.63	-0.13	-1.29	0.02	-0.03	0.00	0.00
2017	0.03	1.08	11.57	-2.61	-0.59	-4.33	-2.22	0.54	-0.59	-0.10	-1.06	0.01	-0.02	0.00	0.00
2018	-0.05	0.75	11.60	-3.23	-1.25	-12.63	-3.98	0.25	-0.62	-0.12	-1.26	-0.04	-0.07	0.00	0.00
2019	-0.05	0.75	11.05	-3.61	-1.77	-3.90	-2.39	0.23	-0.79	-0.13	-1.36	-0.05	-0.09	0.00	0.00
2020	-0.06	-0.06	12.11	-3.28	-1.98	-9.10	-3.29	-0.70	0.11	-0.15	-1.51	-0.13	-0.15	-0.00	0.00
2021	0.72	-0.42	11.73	-2.39	-0.63	6.29	1.49	0.58	1.90	0.12	-0.50	3.48	2.84	2.11	0.70

9.2 PLANNED IMPROVEMENTS

Specific improvements are specified in the QA/QC plan (ISPRA, 2024[b]); they can be summarized as follows.

For the energy and industrial processes sectors, a major progress regards the harmonization of information collected in the framework of different obligations, Large Combustion Plant, E-PRTR and Emissions Trading, thus highlighting the main discrepancies in data and detecting potential errors, the use of data and country specific emission factors collected in national research involving road transport and biomass consumption in residential and for POPs emissions the use of the results of a national research in the potential update of emission factors and methodologies in the iron and steel sector.

For the agriculture and waste sectors, improvements will be related to the availability of new information, on emission factors, activity data as well as parameters necessary to carry out the estimates; specifically, a study on the best available technologies used in agriculture practices, the elaboration of data from the

last farm structure survey and the agricultural census, and availability of information from the exceeding biogas flared at wastewater treatment plants are under investigation.

The EMEP/EEA Guidebook 2019 chapters (EMEP/EEA, 2019) has continued to be considered but the 2023 version started to be used. The update of emission factors will be assessed and applied in the next year submission of the inventory with a focus to PAH estimates in order to improve the completeness, e.g. for PAH compounds, accuracy and reduce the uncertainty.

The comparison between local inventories and national inventory and the meetings and exchange of information with local environmental agencies will continue.

Further analyses will concern the collection of statistical data and information to estimate uncertainty in specific sectors.

10 PROJECTIONS

The projections chapter has not been updated with respect to the previous submission because, as already reported, the Italian National Energy Balance is currently (March 2024) under review. For this reason, the National Emissions Inventory will be revised as soon as the work on the energy balance is concluded. Only subsequently the projections will be revised as the national energy and climate plan is being prepared and this plan must be based on updated and coherent inventory and projections.

The pollutant emission projections presented in this chapter are consistent with the emission projections submitted at EU level according to the Directive (EU) 2016/2284 of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants (NECD).

At its 42nd session (Geneva, 12-16 December 2022), the Executive Body for the LRTAP Convention adopted the 2023 Guidelines for reporting emissions and projections data under the Convention. The Annex IV (template for projections) to the reporting Guidelines was revised and now corresponds to the structure of Annex I. The adopted revised Annex IV has to be applied in 2024 and subsequent years, even though Parties are encouraged to provisionally use it for reporting in 2023. Italy adopted the new format started from the current submission.

The review process carried out during the year 2021 required several improvements as stated in the review report (downloadable from the following link <https://circabc.europa.eu/ui/group/cd69a4b9-1a68-4d6c-9c48-77c0399f225d/library/b70f5178-6e09-443b-a21c-0d3f7e5f96f8/details>).

Major improvement required the consistency of the Italian projection with the Italian NECD inventory, the completeness of reporting NO_x, NMVOC and PM_{2.5} emission for the agricultural sector and the transparency of the projections.

Regarding the consistency, the emission projections are coherent with the actual historical emission submission (year x-2), while NO_x, NMVOC and PM_{2.5} emission from the agricultural sector have been now taken into account. More details will be provided in the following paragraphs.

10.1 THE NATIONAL FRAMEWORK

At national level, the Legislative Decree n. 155 of 2010 (D.Lgs. 2010), that implements the European Directive on air quality, 2008/50/EC (EC, 2008), and the Legislative Decree n. 81 of 2018 (D.Lgs. 2018), that implements the Directive (EU) 2016/2284, provide that ISPRA develops the energy scenario and the scenario of national production activities while ENEA, based on these scenarios, calculates the emission projections using the methodology developed for these purposes at European level.

In this framework, ENEA has elaborated the new national baseline emission scenario using the GAINS-Italy model.

GAINS-Italy is part of the MINNI model, an Integrated Modelling System that links atmospheric science with the economics of emission abatement measures and policy analysis and consists of several interdependent and interconnected components: the national AMS (Atmospheric Modeling System, Mircea et al., 2014; D'Elia et al., 2021) and the national GAINS-Italy (Piersanti et al., 2021; D'Elia et al., 2009). They interact in a feedback system through ATMs (Atmospheric Transfer Matrices) and RAIL (RAINS-Atmospheric Inventory link).

The GAINS-Italy model (Figure 10.1) explores cost-effective multi-pollutant emission control strategies (Ciucci et al., 2016) that meet environmental objectives on air quality impacts (on human health and ecosystems) and greenhouse gases. The current legislation (CLE) scenario represents the 'baseline' and reflects all policies legally in force, both those affecting activity levels (such as energy and agriculture policies), as well as pollution control policies for the period 1990-2050.

The GAINS-Italy model elaborates emission scenarios for air pollutants and greenhouse gases on 5-year time intervals, starting from 1990 to 2050, and evaluates cost-effective multi-pollutant emission control strategies to reach environmental objectives on air quality impacts. Moreover, GAINS-Italy performs fast-response calculations of regional background concentrations of PM_{2.5} and NO₂ in consequence of hypothesized emission reductions on the Italian territory. This last feature is enhanced by the Atmospheric


Transfer Matrices (ATMs), simplified (quasi-linear) relations between total regional emissions and concentrations, calibrated through a set of national Atmospheric Modelling System simulations, based on controlled pollutant emission reductions (Briganti et al., 2011).

The development of an emission scenario with the GAINS-Italy model requires the definition of anthropogenic activity levels, both energy and non-energy, and of a control strategy with a 5-year interval for the period 1990-2050 in the format required by the model. Starting from these information, GAINS-Italy produces alternative future emission and air quality scenarios and abatement costs at a 5-year interval starting from 1990 to 2050.

For the preparation of national emission scenarios, an acceptable harmonization, at a given base year, between the national emission inventory and the GAINS-Italy emissions (D'Elia and Peschi, 2013) has been carried out.

More details about the procedure to build an emission scenario could be found in D'Elia and Peschi, 2016.

Figure 10.1 The scheme of the MINNI national model.



10.2 INPUT SCENARIOS

10.2.1 The energy scenario

The Energy scenario used as input to the GAINS-Italy model has been produced by with the partial equilibrium model TIMES (The Integrated MARKAL-EFOM1 System / EFOM Energy Flow Optimization Model), a model generator for local, national or multi regional economies finalized to the analysis of whole energy systems (electricity generation and consumption, heat distribution, transports, industries,

civil, etc.). The model belongs to the family of MARKAL (Market Allocation, <http://www.iea-etsap.org/web/Markal.asp>) models, the so-called “3e models” (energy, economy, environment), and was developed by the International Energy Agency (IEA) under the program Energy Technology Systems Analysis Program (ETSAP). This model is recognized by the International Panel on Climate Change (IPCC).

The energy system thus simulated is composed by a number of different sectors and subsectors (e.g., electricity production, industrial activities, residential buildings, etc.), each one consisting of a set of technologies connected by input-output linear relationships. Inputs and outputs can be energy carriers, materials, emissions, or requests for services. TIMES is a bottom-up, demand-driven model in which each technology is identified by technical and economic parameters and the production of a good is conditioned to the effective demand by end-users.

The structure of energy scenarios is defined by variables and equations determined by input data constituting the regional database. The database contains qualitative and quantitative data describing the interaction between different components of the energy system.

TIMES identifies the optimal solution to provide energy services at the lowest cost, producing simultaneously investments in new technologies or using more intensively the available technologies in each region defined by the user. For example, an increase in electricity demand for residential use can be satisfied with a more intensive use of available power plants or through the installation of new power plants. Model choices are based on the analysis of technological characteristics of available alternatives, the cost of energy supply and environmental criteria and bounds

The model has been developed considering the detailed energy input needed by GAINS-Italy so that the two models are fully integrated and all the information needed by GAINS-Italy can be found in the TIMES output, that describes, for each sector, the amount of energy carriers, raw materials used, and goods or services produced. The scenarios are consistent with those submitted in 2021 to the EU Commission under Regulation (EU) 2018/1999.

The main assumptions of the scenario can be summarized as follow:

- GDP: economic growth from 2015 to 2020 with average annual rate of -1.32%. For the period 2020-2025 the average annual rate is +2.41%, for period 2025-2030 is +0.30%, and for period 2030-2050 is 0.54%;
- energy: increase of efficiency in finale uses and renewable sources toward EU 2030 climate targets;
- population: decrease with average annual rate of -0.24% in the period 2015-2050.

The scenario considers a slow recovery from the economic crisis that has hit all national activities, and it also considers the development of low carbon technologies and efficiency improvement. The main driving variables used for projections of demand for energy services in the end-use sectors, as well as for activity levels of the industrial processes, are:

- industry: gross value added (GVA) and, for some sub-sectors, physical productions;
- tertiary: GVA;
- residential: demographic trends (mainly population and number of households), increase in the number of appliances per household and growing demand for summer cooling;
- electricity generation: continuation of the ongoing growth of renewable sources;
- transport: dynamics of active population (along with assumptions about mobility per capita) and goods exchange, fleet renewal according to historic market trends.

General economic parameters used for the scenario are those provided by the draft EU reference scenario for Italy in December 2020. Table 10.1 shows actual and projected values for GDP and GVA.

As already mentioned, the GDP is assumed to be reduced from 2015 to 2020 with an average year rate of -1.32%, at it is expected to recover at +0.78% yearly up to 2040. The GDP value for 2020 considers the effect of lockdown due to SARS-Cov2 pandemic. As for value added from productive sectors, a higher recovery for services is projected as compared to industry after 2020. In future years, tertiary sector is expected to continue growing at higher rate than industry, further increasing its role in the Italian economy.

Table 10.2 shows the energy and carbon international prices according to the suggested projections by European Commission. The prices show increasing trends for all the commodities. The increase in carbon price is particularly steep during the periods 2015-2030 (average annual rate 9.4%) and 2030-2040 (5.9% yearly), while the foreseen growth for energy prices is much slower (average annual rate from 0.4% for natural gas to 4.1% for oil in the period 2015-2040).

Table 10.1 – Actual (up to 2015) and projected GDP, and GVA

billion € 2016 constant prices basis		2015	2020	2025	2030	2035	2040	2050
GDP		1,683	1,574	1,773	1,800	1,813	1,837	2,003
Average annual rate (%)		-0.50%	-1.32%	2.41%	0.30%	0.14%	0.27%	1.04%
GVA – industry		239	236	259	260	260	262	278
Average annual rate (%)		-0.16%	-0.26%	1.88%	0.08%	0.04%	0.10%	0.72%
GVA – construction		65	61	69	71	71	72	79
Average annual rate (%)		-6.08%	-1.36%	2.57%	0.36%	0.17%	0.32%	1.12%
GVA – tertiary		1,136	1,061	1,204	1,226	1,236	1,256	1,379
Average annual rate (%)		-0.17%	-1.36%	2.57%	0.36%	0.17%	0.32%	1.12%
GVA – agriculture		35	33	34	34	34	34	34
Average annual rate (%)		0.74%	-0.73%	0.14%	-0.03%	0.00%	0.00%	0.31%

		2015	2020	2025	2030	2035	2040	2050
Coal prices	€ 2016 / GJ	1.8	1.8	2.7	2.9	3.1	3.3	3.5
Oil prices	€ 2016 / GJ	6.0	6.6	11.8	14.0	15.1	16.2	16.6
Gas prices	€ 2016 / GJ	7.5	3.5	5.7	6.0	6.8	8.2	8.6
Carbon price	€ 2016 / t CO ₂	7.8	25	28	30	40	53	84

billion € 2016 constant prices basis		2015	2020	2025	2030	2035	2040	2050
GVA Iron and steel		5.10	5.58	5.95	5.94	5.93	5.92	6.00
Average annual rate (%)		6.18%	1.79%	1.31%	-0.06%	-0.02%	-0.04%	0.18%
GVA Non ferrous metals		2.65	2.72	2.93	2.94	2.93	2.93	2.95
Average annual rate (%)		7.47%	0.51%	1.54%	0.00%	-0.01%	-0.03%	0.08%
GVA Fertilisers/inorganic chemicals		1.88	1.91	1.93	1.89	1.88	1.85	1.71
Average annual rate (%)		-1.88%	0.27%	0.22%	-0.39%	-0.13%	-0.27%	-0.98%
GVA Petrochemicals		2.49	2.84	2.96	2.95	2.95	2.96	3.06
Average annual rate (%)		5.75%	2.63%	0.88%	-0.08%	0.00%	0.04%	0.34%
GVA Other chemicals/ cosmetics		6.65	6.65	7.12	7.18	7.22	7.29	7.99
Average annual rate (%)		-0.35%	-0.01%	1.38%	0.19%	0.09%	0.21%	1.08%
GVA Pharmaceuticals		8.99	9.89	10.01	10.18	10.27	10.46	11.51
Average annual rate (%)		2.01%	1.93%	0.25%	0.32%	0.18%	0.37%	1.13%
GVA Cement and derived products		2.26	2.11	2.39	2.43	2.44	2.47	2.73
Average annual rate (%)		-9.33%	-1.36%	2.48%	0.30%	0.12%	0.25%	1.22%
GVA Ceramics, bricks, etc.		2.74	3.03	3.26	3.30	3.30	3.32	3.44
Average annual rate (%)		-1.78%	1.98%	1.50%	0.24%	0.04%	0.10%	0.32%
GVA Glass production		2.21	2.32	2.44	2.47	2.48	2.51	2.70
Average annual rate (%)		-2.13%	0.95%	1.07%	0.23%	0.09%	0.18%	0.92%
GVA Other non metallic minerals		2.33	2.08	2.35	2.38	2.39	2.41	2.54

Average annual rate (%)	-4.09%	-2.18%	2.46%	0.20%	0.07%	0.16%	0.58%
GVA Paper and pulp production	5.18	5.10	5.24	5.26	5.28	5.31	5.68
Average annual rate (%)	-1.15%	-0.34%	0.56%	0.06%	0.07%	0.15%	0.78%
GVA Printing and publishing	4.24	4.44	4.68	4.71	4.72	4.75	5.09
Average annual rate (%)	-1.03%	0.92%	1.06%	0.13%	0.05%	0.13%	0.83%
GVA Food, drink and tobacco	27.28	28.01	29.73	30.10	30.27	30.61	33.15
Average annual rate (%)	0.38%	0.53%	1.20%	0.25%	0.11%	0.23%	0.93%
GVA Textiles	24.29	19.43	22.39	21.99	21.84	21.62	20.86
Average annual rate (%)	0.34%	-4.37%	2.88%	-0.36%	-0.13%	-0.20%	-0.36%
GVA Engineering	101.86	102.95	114.69	115.33	115.59	116.17	124.70
Average annual rate (%)	-0.44%	0.21%	2.18%	0.11%	0.05%	0.10%	0.87%
GVA Other industries	38.60	36.61	40.62	40.80	40.89	41.06	43.58
Average annual rate (%)	-0.99%	-1.05%	2.10%	0.09%	0.04%	0.08%	0.69%

The population grew significantly from 2005 to 2015 with annual average rate of 0.49% and started to decline from 2015 to 2020 with annual rate -0.19%. The declining trend is expected all along the time series until 2050. In forthcoming years, the difference with the previous submission is noticeable with a growing divergence (Table 10.4).

Table 10.4 – Population

	2015	2020	2025	2030	2035	2040	2050
Thousands	60,796	60,233	59,583	58,941	58,341	57,711	55,860

The next table shows the number of persons per household adopted for GHG projections in residential sector.

Table 10.5 – Inhabitants per household

	2015	2020	2025	2030	2035	2040	2050
Inhabitants per household	2.35	2.31	2.42	2.40	2.37	2.35	2.29

Table 10.6 shows data of transport demand for passengers, freights, domestic navigation, and air traffic. The expected activity scenario for transport shows a sharp decline in 2020 due to the mentioned lockdown. After 2020 projections show a steady growth up to 2050. The transport demand decreases up to 2020 compared to 2015 with annual rate of -4.3% for passengers while show a weak growth of 0.24% for goods. After 2020, up to 2050, the annual growth rate is around 1.2% for passengers and goods, with a much steeper increase until 2025 because of the end of the pandemic.

Table 10.6 – Transport demand for passengers and freights

		2015	2020	2025	2030	2035	2040	2050	
Passenger	Road	billion pass-km	820.2	670.7	875.2	877.1	885.4	898.1	919.6
	Rail	billion pass-km	59.5	31.7	60.3	64.8	66.5	68.1	72.4
	Domestic aviation	Number of Landing and Take-Off cycle (LTO)	380.6	121.0	383.2	434.1	471.5	512.2	567.8
	International aviation	Number of Landing and Take-Off cycle (LTO)	325.4	115.7	366.3	416.3	457.6	497.6	558.2
	Total	billion pass-km	876.8	702.4	935.5	941.9	951.9	966.1	992.0

	Road	billion ton-km	124.9	144.5	168.3	183.8	190.1	196.2	208.6
	Rail	billion ton-km	20.8	20.5	24.1	25.9	26.9	27.7	29.0
Freight	Domestic navigation (inland waterways and national maritime)	billion ton-km	51.2	44.6	59.8	61.1	61.8	62.6	65.2
	Total	billion ton-km	207.1	209.6	252.3	270.8	278.8	286.5	302.8

The gross inland consumption of energy (GIC), estimated according to the methodology adopted by Eurostat, is expected to be about 138.7 Mtoe in 2030 with an average yearly decrease rate of -0.8% since 2015. After the further fall in 2020 and the rebound effect in 2025 the projected gross inland consumption shows constant decrease to 124.6 Mtoe up to 2050.

GIC started to decrease since 2005, before the economic crisis, while in the period 1990-2005 it has constantly increased with an annual average equal to +1.7%. The share of natural gas increased constantly since 1990 counterbalancing the corresponding decrease of oil share. Since 2007 it is also evident the growing role of renewable energies (Figure 9.1).

Relevant changes are expected in the estimated total energy consumption between the previous projections and the last one due to updated data for base year and adoption of new measures. The most relevant changes occurred in 2020 following the measures adopted to slow down the diffusion of SARS-Cov2 pandemic followed by the rebound effect in the next years.

10.2.2 The scenario of non-energy activities

To develop an emission scenario, the GAINS-Italy model requires the definition also of non-energy activities level. The definition of such scenario is based on economic variables, like GDP (gross domestic product) or added value derived from the energy scenario, population data or specific sector statistics.

Livestock projection has been carried out with a statistical model where the number of animals has been linked to the projections of other variables, like meat consumption and production, or milk consumption and production (see equation 9.1)

$$(n^{\circ}heads_i) = \left(\frac{n^{\circ}heads_i}{MP_i} \right) \times \left(\frac{MP_i}{MC_i} \right) \times \left(\frac{MC_i}{MC_{tot}} \right) \times \left(\frac{MC_{tot}}{Pop} \right) \times (Pop) \quad (9.1)$$

where the head number of livestock *i* is linked to meat production (MP) and consumption (MC) of livestock *i*, total meat consumption (MC_{tot}) and population (Pop).

All the details about this methodology are provided in D'Elia and Peschi, 2013.

The updated livestock projections have been elaborated for the baseline scenario: population data (Pop) are the same of the energy scenario, while for meat production and consumption sectorial studies and statistics have been considered.

In the following figures, the results for the main livestock are reported where ISPRA stands for the activity data considered in the 2022 emission submission, that was the last submission available when the livestock projections were elaborated; WM are the projections elaborated for the baseline scenario.

Figure 10.2 Livestock scenario comparison for dairy cows and other cattle.





Figure 10.3 Livestock scenario comparison for pigs and sheep and goats.



All the activity data for the agricultural, solvent and industrial sector are reported in the Appendix.



10.2.3 The control strategy definition

In addition to energy, climate and agricultural policies assumed in the energy, non-energy and agricultural input scenarios, in the baseline emission projections a detailed inventory of national emission control legislation is considered (Amann et al., 2011).

In the WM scenario it is assumed that all the European and national regulations adopted before 2021 will be fully complied according to the foreseen time schedule. Examples of the legislations considered are the Directive on Industrial Emissions for large combustion plants, the Directives on Euro standards, Solvent Directive, the Code of Agricultural Good Practice.

An example of the WM control strategy for the road transport sector is reported in the following figures. Moreover, the emission factors for the different pollutant and technologies are reported in the Annex.

Figure 10.4 Control strategy for gasoline (on the left) and diesel (on the right) cars.



10.3 THE HARMONIZATION PROCESS

The first step for the preparation of a new national emission scenario is to align at a given base year the latest national emission inventory submission and the GAINS-Italy emissions, estimated with a top-down approach. Being a Party of the United Nations Economic Commission for Europe (UNECE) Convention on Long Range Transboundary Air Pollution (CLRTAP), Italy has to annually submit an emission inventory of air pollutants and provide a report on its data according to the Guidelines for reporting emissions and projections data (UNECE, 2015). On the other hand, to produce a reliable emission scenario, GAINS-Italy model produces its own emission estimates, for the years considered in the model, with its own classification system. Discrepancies between the inventory and the GAINS-Italy output exist and are due to different reasons, such as, for example, different coverage and aggregation of emission sources, different emission calculation methodologies. These discrepancies need so to be solved and the emission estimates to be aligned. This alignment step is called harmonization and is needed to validate the emission scenario to base emission time trends in GAINS-Italy on a reliable starting point. In the harmonization process, activity data, emission factors and technologies for each sector are compared. If discrepancies emerge (for example in fuel allocation across sectors or different assumptions on control measures in place in the year of comparison), the model parameters will be modified according to the inventory with the attempt to let GAINS-Italy reproduce emissions as closely as possible to the national emission inventory. Further details about the harmonization method are reported in D'Elia and Peschi, 2013. For all these reasons, a comparison between the last national emission inventory, and the GAINS-Italy emission estimates has been carried out considering three historical years, 2005, 2010 and 2015. Results of the harmonization process between the 2023 emission inventory submission (INVENTORY_sub2023) and GAINS-IT estimates are summarized in Table 10.7.

Table 10.7 Comparison of total emissions in the last submission of the national inventory report

Pollutant	Emissions 2005			Emissions 2010			Emissions 2015		
	Inventory_sub2023 (kt)	GAINS-IT (kt)	Δ (%)	Inventory_sub2023 (kt)	GAINS-IT (kt)	Δ%	Inventory_sub2023 (kt)	GAINS-IT (kt)	Δ (%)
SO ₂	411	421	2%	224	219	-2%	128	124	-3%
NO _x	1231	1236	0%	893	845	-5%	679	684	1%
PM _{2.5}	186	182	-2%	213	202	-5%	169	169	-0%
NM _{VOC}	1202	1247	4%	988	1018	3%	778	810	4%
NH ₃	421	422	0%	379	381	1%	357	378	6%

Discrepancies in reproducing the national emission inventory have been considered acceptable if differences remain within a few percentage points, i.e. in the interval between $\pm 5\%$. In the comparison, NO_x and NM_{VOC} emissions from the agricultural sector, even if calculated also by the model, have not been considered to be coherent with the compliance of the NEC ceilings.

In the following plots, details on sectoral emissions by NFR code (Nomenclature For Reporting; Table 10.8 reports the sectors considered) are illustrated for the year 2015.

Table 10.8 Definition of the NFR code used in the comparison between emission inventory and GAINS-IT

NFR code	Description
1A1	Energy industries (Combustion in power plants & Energy Production)
1A2	Manufacturing Industries and Construction (Combustion in industry including Mobile)

NFR code	Description
1A3b	Road and Off-road Transport
1A4	Other sectors (Commercial, institutional, residential, agriculture and fishing stationary and mobile combustion)
1A5	Other
1B	Fugitive emissions (Fugitive emissions from fuels)
2	Industrial Processes and Solvent use
3	Agriculture
5	Waste
6A	Other (included in National Total for Entire Territory)

For SO₂ emissions, the model shows a slight underestimate of the sectors 1A1 and 1A2, and for NO_x in the sector 1A3 and overestimate in the sector 1A4. PM_{2.5} shows a good agreement between the two estimates with the exception of the sector 6 – Other where the model estimates emissions from barbecue, fireworks not estimated in the inventory for their high uncertainties. The model shows a slight overestimate in NMVOC emissions from the sector 2 – Industrial process and solvent use and in NH₃ emissions from the sector 3 – Agriculture in the range of uncertainty considered acceptable.

Figure 10.6 NO_x national emission harmonization between the last emission inventory and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 10.7 NMVOC national emission harmonization between the last emission inventory (INVENTORY_sub2023) and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 10.8 SO_x national emission harmonization between the last emission inventory (INVENTORY_sub2023) and GAINS-IT estimates detailed by NFR sectors for the year 2015.




Figure 10.9 NH₃ national emission harmonization between the last emission inventory (INVENTORY_sub2023) and GAINS-IT estimates detailed by NFR sectors for the year 2015.





Figure 10.10 PM_{2.5} national emission harmonization between the last emission inventory (INVENTORY_sub2023) and GAINS-IT estimates detailed by NFR sectors for the year 2015.



10.4 THE EMISSION SCENARIO

The result of the activity input scenarios and of the harmonization process is an emission scenario.

In the following figures, a comparison of the 2023 emission inventory submissions, the WM and WAM scenarios are presented. Details by NFR sector are presented for the WM and WAM scenarios.

A huge decrease in SO₂ emissions is projected (Figure 10.11) driven by the energy and the maritime sector for the year 2030 while the industrial sector (1A2) in the year 2030 represents the main emitting sector (Figure 10.12).

Figure 10.11 Reported and projected (WM, WAM) SO_x emissions elaborated by the GAINS-Italy model.





Figure 10.12 SO_x emission scenario WM (on the left) and WAM (on the right) by



A huge decrease is estimated in NO_x emission scenarios (Figure 10.13) due to the diffusion of new diesel Euro 6 and electric vehicles. The road transport sector still represents the principle NO_x source (Figure 10.14).

Figure 10.13 Reported and projected (WM and WAM) NO_x emissions elaborated by the GAINS-Italy model.




Figure 10.14 NO_x emission scenario WM (on the left) and WAM (on the right) by sector.




Figure 10.15 Reported and projected (WM and WAM) PM2.5 emissions elaborated by the GAINS-Italy model.





Figure 10.16 PM2.5 emission scenario WM (on the left) and WAM (on the right) by sector.



The decrease of NMVOC emissions (Figure 10.17) is driven by the solvent that will remain the main emitting sector (Figure 10.18).

Figure 10.17 Reported and projected (WM and WAM) NMVOC emissions elaborated by the GAINS-Italy model.





Figure 10.18 NMVOC emission scenario WM (on the left) and WAM (on the right) by sector.



NH₃ is the pollutant with less variations (Figure 10.19) whose main contribution to total NH₃ emissions is due by the agricultural sector (Figure 10.20).

Figure 10.19 Reported and projected (WM and WAM) NH₃ emissions elaborated by the GAINS-Italy model.





Figure 10.20 NH₃ emission scenario WM (on the left) and WAM (on the right) by sector.



10.5 THE NEC EMISSION TARGETS

The NEC Directive (EU, 2016), implemented in the Italian legislation in the D.Lgs. 81/2018, defines for each Member States the emission reduction targets in the year 2020 and 2030 respect to the base year 2005 for the anthropogenic emissions of SO₂, NO_x, PM_{2.5}, NMVOC and NH₃.

In Table 10.9 the attainment of the national emission reductions in the year 2030 in the comparison with the new National Emission Ceilings Directive (NECD) targets is reported.

2030 EMISSION REDUCTIONS			
	NECD targets	WM	WAM
SO₂	-71%	-84%	-86%
NO_x	-65%	-73%	-79%
PM_{2.5}	-40%	-57%	-57%
NM_{VOC}	-46%	-53%	-53%
NH₃	-16%	-25%	-25%

According to the present emission projections, the WM scenario could attain the emission target for the year 2030. Additional measures in the WAM scenario have been considered especially in the road transport sector where higher electric vehicles spread is foreseen.

Detailed data on the drivers influencing the emission projections can be found in the Appendix 2.

11 REPORTING OF GRIDDED EMISSIONS AND LPS

Every four years, from 2017 with reference to 2015 emissions, ISPRA shall provide the disaggregation of the national inventory at provincial level as instituted by the Legislative Decree n. 81 of 30 May 2018. The emissions disaggregated at regional and provincial levels are compared to the results obtained by regional bottom-up inventories. Emissions disaggregated at local level are also used as input for air quality modelling. Methodologies and proxies are described in the relevant publication (ISPRA, 2009) (ISPRA, 2022) available on: <https://emissioni.sina.isprambiente.it/inventari-locali/> .

Simultaneously, ISPRA carries out the spatial disaggregation of national emissions on the 0.1x0.1° EMEP grid.

The grid definition for a country contains a dataset for each grid cell with information about (CEIP, 2019):

- the country or area (ISO2 or a three digits abbreviation for countries and other areas which you can see in the column Country code in the grid definition table above)
- the country-or area name
- the longitude position of the grid cell (centre of the cell)
- the latitude position of the grid cell (centre of the cell)
- and fraction of the grid cell (share of the cell area which belongs to the country/area, e.g. 1 for cells which are completely inside the country/area borders and e.g. 0.5 for cells where half of the area belongs to the country/area and the rest is outside the boundary)

According to the review process (EEA, 2020), the reporting of gridded data has been aligned with the last available reporting guidelines and the relevant GNFR codes in the next submission, as it is indicated in the Table 11.1. Proxy used for disaggregation have been reported too.

Table 11.1 Aggregation for Gridding and LPS of NFR sector

NFR Aggregation for Gridding and LPS (GNFR)	Proxy
A_PublicPower	point sources consumptions from ETS/LPS, see paragraph "LPS data"
B_Industry	combustion in industry prevalently point sources consumptions data from ETS/LPS, where not available production data. For industrial processes prevalently production data from E-PRTR or producers associations, in a few of case production capacity or data about employees at NUTS3 level, see paragraph "LPS data".
C_OtherStationaryComb	resident people and fuels sold at NUTS3 level
D_Fugitive	see paragraph "fugitive"
E_Solvents	prevalently employees at NUTS3 level. Data from the national institut of statistics (ISTAT)
F_RoadTransport	see paragraph "transport"
G_Shipping	see paragraph "transport"
H_Aviation	see paragraph "transport"
I_Offroad	see paragraph "transport"
J_Waste	amount of managed waste at NUTS3 level on the basis of the management system for SWDS, biological treatments and incineration. Source: ISPRA. For domestic wastewater resident people at Nuts3 level while data about employees at NUTS3 level in the case of industrial wastewater have been used. Source: ISPRA.
K_AgriLivestock	see paragraph "agriculture"
L_AgriOther	see paragraph "agriculture"
M_Other	
N_Natural	

NFR Aggregation for Gridding and LPS (GNFR)	Proxy
O_AviCruise	see paragraph "transport"
P_IntShipping	see paragraph "transport"
z_Memo	

The methodologies for spatial disaggregation are consistent with those reported in the EMEP/EEA 2019 Guidebook and described in relevant report (ISPRA, 2009). National emissions have been disaggregated at provincial level (NUTS3) because of the availability of proxy data, then the allocation to the 0.1x0.1° grid has been realized on the basis of the 2019 EMEP/EEA Guidebook (EMEP/EEA, 2019). In particular, point sources have been allocated directly to the grid within which they are contained by converting the x, y values to that of the coordinates used to geo-reference the grid. For area sources (NUTS3 polygons) the fraction of the area of the polygons has been used to distribute the emissions from the original polygon to the intersected grid cell. Finally, the spatially resolved (mapped) detailed NFR sectors have been aggregated in the GNFR code.

11.1 FUGITIVE

Proxy used for disaggregation of 1B1 and 1B2 are reported in the Table 11.2.

Table 11.2 Proxy for the disaggregation of fugitive emissions

Category	Proxy
1 B 1 a (SNAP 050102)	Amount of coal production
1 B 1 a (SNAP 050103)	Amount of solid fuels consumption from ETS plants
1 B 2 a iv	Amount of crude oil refined at plant level
1 B 2 a v (SNAP 050502)	Amount of gasoline sold at provincial level.
1 B 2 a v (SNAP 050503)	Amount of mineral oil and LPG in depots for industrial and services at regional level distributed according to the amount of gasoline sold at provincial level.
1 B 2 b (SNAP 050601)	Amount of natural gas consumption in the compressor stations (ETS data)
1 B 2 b (SNAP 050603)	Amount of natural gas provided to the distribution network at provincial level (other destination than industrial and thermoelectric users)
1 B 2 c	Amount of crude oil refined at plant level

11.2 TRANSPORT

Proxy used for disaggregation of the different transport mode are reported in the Table 11.3.

Table 11.3 Proxy for the disaggregation of transport emissions

Category	Proxy
1A2gvii	ISPRA study about industrial machinery data elaborated at NUTS 3 level.
1A3ai(i)	Emissions deriving from international LTO, in the detail of Italian airports, elaborated by Eurocontrol to the aim of the estimation of emissions from aviation for Member Countries Inventories, then aggregated at NUTS 3 level.

Category	Proxy
1A3aii(i)	Emissions deriving from domestic LTO, in the detail of Italian airports, elaborated by Eurocontrol to the aim of the estimation of emissions from aviation for Member Countries Inventories, then aggregated at NUTS 3 level.
1A3bi	Ministry of Transport data about passenger cars fleet at NUTS 3 level, according to Copert classification; traffic flows and length of motorway sections as regards highway share.
1A3bii	Ministry of Transport data about light duty vehicles fleet at NUTS 3 level, according to Copert classification; Value added as regards urban and rural share; traffic flows and length of motorway sections as regards highway share.
1A3biii	Ministry of Transport data about vehicles fleet at NUTS 3 level, according to Copert classification; Value added as regards urban and rural share for heavy duty vehicles; traffic flows and length of motorway sections as regards highway share.
1A3biv	Ministry of Transport data about mopeds and motorcycles fleet at NUTS 3 level, according to Copert classification; traffic flows and length of motorway sections for motorcycles categories as regards highway share.
1A3bv	Same proxies as gasoline vehicles categories. Evaporative emissions are added to those due to combustion, for each vehicle category, according to Copert classification, for each driving cycle, therefore disaggregated at NUTS 3 level according to the criterion used for other road transport activities. Therefore emissions related to this activity are included in the other NFR classes.
1A3bvi	Same proxies as corresponding vehicles categories, according to Copert classification. Non exhaust emissions are added to those due to combustion, for each vehicle category, according to Copert classification, for each driving cycle, therefore disaggregated at NUTS 3 level according to the criterion used for other road transport activities. Therefore emissions related to this activity are included in the other NFR classes.
1A3c	Ministry of Transport data about the length of non-electrified railway sections, relating to Italian State Railways Group and to the Regional and/or local railways network, elaborated at NUTS 3 level.
1A3dii	Data elaborated at NUTS 3 level covering: Eurostat data about vessels, by type, in harbours ; ships berths for: sailing boats, motorboats, watercrafts; simplified trajectories for cruise; inland waterways traffic is estimated by attributing a share to freight traffic (provinces of the Po basin) and a share to passenger traffic on the basis of Ministry of transport data about the number of boats for province.
1A4bii	Data at NUTS 3 level about mechanical means used, deriving from surveys on the structure and production of farms, performed by the Italian Institute of Statistics.
1A4cii	Data at NUTS 3 level about mechanical means used, deriving from surveys on the structure and production of farms, performed by the Italian Institute of Statistics.
1A4ciii	Data, from Economic Observatory on the production structures of maritime fishing in Italy, about fishing boats number and consumptions at NUTS 3 level.
1A5b	National Institute of Statistics data about resident population at NUTS 3 level have been used as proxy for military mobile activities.

11.3 AGRICULTURE

Proxy used for disaggregation of the agriculture sector are reported in the Table 11.4. The methods used for gridding are reported below Table 11.1 and are the same for all inventory sectors.

Table 11.4 Proxy for the disaggregation of agriculture sector

Category	Proxy
3Da1	Quantity of annual nitrogen fertilizers (tons) distributed in the Italian provinces (National Institute of Statistics - ISTAT); for CO ₂ from liming: annual amount of lime and dolomite distributed (ISTAT); for CO ₂ from urea application: annual amount of urea distributed (ISTAT)
3C	Cultivated area (hectares) for the production of rice (ISTAT)
3Da2, 3Da3, 3De	The activity includes estimates of emissions from spreading livestock manure, grazing, nitrogen-fixing process of leguminous crops (NH ₃ emissions only), use of other organic fertilizers (compost, other soil conditioners, animal and vegetable processing wastes, other), agricultural sludge spreading, soil input from crop residues (N ₂ O emissions only), and cultivation of organic soils (N ₂ O emissions only). NMVOC emissions arise only from the spreading of livestock manure and from grazing.

Category	Proxy
	<p>Provincial data used for disaggregation are: agricultural area used (hectares) and annual production harvested (tons), number of livestock, organic fertilizers used, tons of sludge spread in agriculture, area of leguminous crops most cultivated (alfalfa, soybean, clover, grain bean), organic soils.</p> <p>National emissions from this sector were disaggregated proportionally to: agricultural area used for legumes; livestock stock for emissions from grazing and manure spreading; agricultural production for emissions from nitrogen contributed by crop residues; amount of sludge spread on agricultural soils and agricultural area used for emissions from the use of sewage sludge; amount of organic manures and soil conditioners for emissions from the distribution of other organic fertilizers; provincial areas of organic soils for emissions generated by this type of soils.</p> <p>As for provincial NH₃ emissions from spreading, a provincial distribution was constructed of emissions from spreading from the provincial distribution of total ammonia emissions (i.e., from sheltering, storage, spreading, and grazing) by animal category and the percentage of emission from spreading (calculated considering the emission factors of the four stages of manure management).</p> <p>The provincial distribution of emissions total ammonia emissions was calculated from the provincial emission factors (totals of the four stages of management of manure) compiled by CRPA (in a 1997 study), to which a variation was applied based on the emission factors updated by animal category (but constant for all provinces), and the provincial animal numbers from ISTAT sources.</p> <p>For the disaggregation of emissions from grazing, the same procedure as for spreading was used as just described.</p> <p>As for provincial emissions of N₂O, NO_x and NMVOC from grazing and spreading, a provincial distribution of nitrogen excreted at grazing and shelter separately, based on the coefficients of nitrogen excreted at grazing and sheltering by animal category and the provincial distribution of livestock numbers from ISTAT sources. The provincial distribution of nitrogen at shelter was used to disaggregate emissions from spreading.</p>
3F	Annual production of cereal harvest (quintals) (ISTAT)
3A1	For cattle, provincial emissions were estimated by multiplying the number of heads available at provincial level (ISTAT) by the provincial emission factors from a 1994 study (Research Centre on Animal Production - CRPA). National emissions were then disaggregated on the basis of these provincial emissions
3A	Number of heads available at provincial level (ISTAT)
3B	<p>As regards CH₄, COVNM, PM₁₀, PM_{2.5}, NH₃, NO_x emissions, national emissions for cattle and swine are estimated according to IPCC and EMEP/EEA for manure management (housing and storage) and were distributed by province based on the number of livestock. The 2013 Farm Structure Survey (FSS) data were used to disaggregate 2015 national emissions, and the 2016 FSS was used for 2019 emissions. NMVOC emissions from storage were disaggregated based on the distribution provincial emissions of NMVOCs from storage, processed with provincial emission factors for cattle and swine, estimated by CRPA in a 1997 study, and provincial consistencies from ISTAT sources. Emissions of NH₃ from shelter and storage were disaggregated by constructing two distributions of provincial emissions from sheltering and storage separately. The procedure for calculating these distributions is the same as that used for spreading and described in SNAP category 1002 with respect to the emissions of NH₃.</p> <p>Considering emissions for buffalo, sheep, goats, equines, poultry, rabbits, fur animals national emissions are estimated according to IPCC and EMEP/EEA for manure management (housing and storage) and were distributed by province based on the number of animals. The 2013 FSS data were used to disaggregate 2015 national emissions, and the 2016 FSS was used for 2019 emissions. NMVOC emissions from storage were disaggregated based on the provincial distribution of methane emissions from storage, compiled with 2006 IPCC emission factors by climate band (Cool (< 15°C) and Temperate (15 to 25°C)) and provincial consistencies from ISTAT sources. NH₃ emissions from shelter and storage were disaggregated by constructing two distributions of provincial emissions from shelter and storage separately. The procedure for calculating these distributions is the same as that used for spreading and described in the SNAP category 1002 with respect to emissions of NH₃.</p> <p>As regards N₂O emissions, due to animal manure stored, were distributed at the provincial level, starting in 2000, based on the provincial distribution of nitrogen excreted at the shelter, estimated with the coefficients of excreted nitrogen at the shelter and the provincial distribution of the number of animals by animal category. For the years 1990 and 1995, national emissions were disaggregated with the distribution of provincial N₂O emissions from storage, estimated by CRPA.</p>
3Df	Data on the provincial sale of pesticides containing HCB (ISPRA elaborations on ISTAT data)

Category	Proxy
5C2	Annual production of cereal and woody crops harvest (quintals) (ISTAT)

11.4 LPS DATA

GNFR	n° plant	Height class/n°
A_PublicPower	207	2/194; 4/4; 5/9
B_Industry	330	1/328; 4/2
D_Fugitive	117	1/117
G_Shipping	77	1/77
H_Aviation	106	1/106
I_Offroad	22	1/22

LPS data for the year 2019 have been submitted in 2021. A brief description of data is reported in the following.

Table 11.5 Number of plants and relevant height class

GNFR	n° plant	Height class/n°
A_PublicPower	207	2/194; 4/4; 5/9
B_Industry	330	1/328; 4/2
D_Fugitive	117	1/117
G_Shipping	77	1/77
H_Aviation	106	1/106
I_Offroad	22	1/22

Following the review process, some errors on longitude and latitude of plants have been corrected, also thanks the implementation procedures of the IED (Industrial Emissions Directive) European Directive.

Italy has decided not to provide all and only PRTR data as LPS data as in Italy some PRTR categories do not respond to the characteristics of large point sources. This is especially true for farms that cannot be considered large point sources and therefore have been eliminated with respect to the previous submission. The current submission is instead consistent with gridded data and includes sources outside of PRTR which are LPS such as ports or extractive activities.

12 REFERENCES

12.1 INTRODUCTION

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2009. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 9/2009.

EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 12/2013.

EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

EMEP/EEA, 2023. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 6/2023.

EU, 2016. National Emissions Ceilings Directive 2016/2284/EU.

EEA, 2017 [a]. Final Review Report, 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy. 30 November 2017. Reference: No 07.0201/2016/741511/SER/ENV.C.3. Umweltbundesamt Wien.

EEA, 2018. Final Review Report, 2018 Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD'). Italy. 30 November 2018. Reference: No 070203/2017/765105/SER/ENV.C.3. Umweltbundesamt Wien.

EEA, 2019. Final Review Report, 2019 Third phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD'). Italy. 22 November 2019. Reference: Service request No 4 under Framework contract No ENV.C.3/FRA/2017/0012. International Institute for Applied Systems Analysis (IIASA) Laxenburg

EEA, 2020. Final Review Report, 2020 Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 20 November 2020. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2022. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2022 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 11 November 2022. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2023. Final Review Report, Review of National Air Pollutant Emission Inventory Data 2023 under Directive (EU) 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/815979/SER/ENV.C.3. Umweltbundesamt Wien.

IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html

ISPRA, 2014. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Procedures Manual. Report 112/2014. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/> .

ISPRA, 2018. National Greenhouse Gas Inventory System in Italy. March 2018, <https://emissioni.sina.isprambiente.it/inventario-nazionale/>

ISPRA, 2022. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. ISPRA, 369/2022. URL: <https://emissioni.sina.isprambiente.it/inventari-locali/>

ISPRA, 2024[a]. Italian Greenhouse Gas Inventory 1990-2022. National Inventory Report 2024. March 2024, <https://emissioni.sina.isprambiente.it/inventario-nazionale/> .

ISPRA, 2024[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2024. March 2024, <https://emissioni.sina.isprambiente.it/inventario-nazionale/> .

Romano D., Bernetti A., De Lauretis R., 2004. Different methodologies to quantify uncertainties of air emissions. Environment International vol 30 pp 1099-1107.

UNECE, 2008. Guidelines for Reporting Emission Data under the Convention on Long-range Transboundary Air Pollution, Emission Reporting Guidelines. ECE/EB.AIR/2008/4 23 September 2008.

UNECE, 2010. Report for the Stage 3 in-depth review of emission inventories submitted under the UNECE LRTAP Convention and EU National Emissions Ceilings Directive for Italy. CEIP/S3.RR/2010/ITALY (25/11/2010).

UNECE, 2013. Report for the Stage 3 in-depth review of emission inventories submitted under the UNECE LRTAP Convention and EU National Emissions Ceilings Directive for Italy. CEIP/S3.RR/2013/ITALY (20/08/2013)

UNECE, 2014. Guidelines for Reporting Emissions and Projections Data under the Convention on Long Range Transboundary Air Pollution, ECE/EB.AIR/125 13 March 2014.

12.2 ANALYSIS OF KEY TRENDS BY POLLUTANT

Decree of President of the Republic 24 May 1988, n. 203. Attuazione delle direttive CEE numeri 80/779, 82/884, 84/360 e 85/203 concernenti norme in materia di qualità dell'aria, relativamente a specifici agenti inquinanti, e di inquinamento prodotto dagli impianti industriali, ai sensi dell'art. 15 della L. 16 aprile 1987, n. 183 (2) (2/a). Gazzetta Ufficiale del 16 giugno 1988, n. 140, S.O.

EC, 1975. Council Directive 75/716/EEC of 24 November 1975 on the approximation of the laws of the Member States relating to the sulphur content of certain liquid fuels.

EC, 1988. Council Directive 88/609/EEC of 24 November 1988 on the limitation of emissions of certain pollutants into the air from large combustion plants.

EC, 1991. Council Directive 91/441/EEC of 26 June 1991 amending Directive 70/220/EEC on the approximation of the laws of the Member States relating to measures to be taken against air pollution by emissions from motor vehicles.

EC, 1994. Directive 94/12/EC of the European Parliament and the Council of 23 March 1994 relating to measures to be taken against air pollution by emissions from motor vehicles and amending Directive 70/220/EEC.

EC, 1996. Council directive 96/61/EC of 24 September 1996 concerning integrated pollution prevention and control. Official Journal L 257 , 10/10/1996 P. 0026 – 0040.

EC, 1997 [a]. Directive 97/68/EC of the European Parliament and of the Council of 16 December 1997 on the approximation of the laws of the Member States relating to measures against the emission of gaseous

and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EC, 1997 [b]. Directive 97/24/EC of the European Parliament and of the Council of 17 June 1997 on certain components and characteristics of two or three-wheel motor vehicles.

EC, 1998. Directive 98/69/EC of the European Parliament and of the Council of 13 October 1998 relating to measures to be taken against air pollution by emissions from motor vehicles and amending Council Directive 70/220/EEC.

EC, 1999. Council Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations.

EC, 2001. National Emission Ceilings Directive: Directive 2001/81/EC.

EC, 2004. Council Directive 2004/26/EC of 21 April 2004 amending Directive 97/68/EC on the approximation of the laws of the Member States relating to measures against the emission of gaseous and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EU, 2016. National Emissions Ceilings Directive 2016/2284/EU.

ISTAT, 2014. I consumi energetici delle famiglie, anno 2013. <http://www.istat.it/it/archivio/142173>

Ministerial Decree 8 May 1989. Limitazione delle emissioni in atmosfera di taluni inquinanti originati dai grandi impianti di combustione. Gazzetta Ufficiale Italiana n. 124 del 30/05/1989.

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

Ministerial Decree 16 May 1996, n. 392. Regolamento recante norme tecniche relative alla eliminazione degli olii usati. G. U. n. 173 del 25/07/1996.

Ministerial Decree 21 January 2000, n. 107. Regolamento recante norme tecniche per l'adeguamento degli impianti di deposito di benzina ai fini del controllo delle emissioni dei vapori. G. U. 02/05/2000 n. 100.

MASE, several years [a]. Bilancio Energetico Nazionale (BEN). Ministero delle Attività Produttive, Direzione Generale delle Fonti di Energia ed industrie di base. <http://dgerm.sviluppoeconomico.gov.it/dgerm/ben.asp> .

12.3 ENERGY (NFR SECTOR 1)

ACI, several years. Annuario statistico. Automobile Club d'Italia, Roma. <http://www.aci.it/index.php?id=54>.

AEEG, several years. Qualità del servizio gas. Autorità per l'energia elettrica e il gas. http://www.autorita.energia.it/it/dati/elenco_dati.htm .

AISCAT, several years. Aiscat in cifre. Data and reports available on website at: http://www.aiscat.it/pubbl_cifre.htm?ck=1&sub=3&idl=4&nome=pubblicazioni&nome_sub=aiscat%20in%20cifre .

ALTROCONSUMO, 2018. Characterization of residential biomass fired appliances emissions with "real-world" combustion cycles. Technical Report (rev.2). Laboratorio Energia e Ambiente Piacenza. Società Consortile partecipata dal Politecnico di Milano. Piacenza December, 3rd 2018.

AMA-Comune di Roma, 1996. Nuovo impianto per l'incenerimento dei rifiuti ospedalieri. Rapporto AMA.

ANCMA, several years. Data available on website at: <http://www.ancma.it/statistiche/-/statistics> .

ANPA, 2001. Redazione di inventari nazionali delle emissioni in atmosfera nei settori del trasporto aereo e marittimo e delle emissioni biogeniche. Rapporto finale. Gennaio 2001

ARPA, 2007 – Stima dei consumi di legna da ardere ed uso domestico in Italia. Ricerca commissionata da APAT e ARPA Lombardia, Rapporto finale, marzo 2007.

ASSOTERMICA, several years. Studio Statistico, <http://www.anima.it/contenuti/10670/studi-di-mercato> .

Berdowski J.J.M., Baas J., Bloos J.P.J., Visschedijk A.J.H., Zandveld P.Y.J., 1997. The European Emission Inventory of Heavy Metals and Persistent Organic Pollutants for 1990. TNO Institute of Environmental Sciences, Energy Research and Process Innovation, UBA-FB report 104 02 672/03.

CESI, 2005. Caratterizzazione delle emissioni di caldaie residenziali, CESI, 2005.

CNR STEMS, Innovhub SSI, 2020. Database of fuel consumption and emission factors derived from national measurements campaign. Internal documentation.

CONFETRA, several years. Il trasporto merci su strada in Italia. Data and reports available on website at: <http://www.confetra.it/it/centrostudi/statistiche.htm> .

Decree of President of the Republic 24 May 1988, n. 203. Attuazione delle direttive CEE numeri 80/779, 82/884, 84/360 e 85/203 concernenti norme in materia di qualità dell'aria, relativamente a specifici agenti inquinanti, e di inquinamento prodotto dagli impianti industriali, ai sensi dell'art. 15 della L. 16 aprile 1987, n. 183 (2) (2/a). Gazzetta Ufficiale del 16 giugno 1988, n. 140, S.O.

De Stefanis P., 1999. Personal communication.

De Stefanis P., 2012. Personal communication (mail 16 November 2012).

EC, 2004. Council Directive 2004/26/EC of 21 April 2004 amending Directive 97/68/EC on the approximation of the laws of the Member States relating to measures against the emission of gaseous and particulate pollutants from internal combustion engines to be installed in non-road mobile machinery.

EDISON, several years. Rendiconto ambientale e della sicurezza.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report, 2020 Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 20 November 2020. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, 2022. Final Review Report, 2022 Review of National Air Pollutant Emission Inventory Data 2022 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 11 November 2022. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.

EEA, several years, Monitoring CO₂ emissions from passenger cars and vans, EEA technical reports.

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 12/2013.

EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

EMEP/EEA, 2023. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 6/2023.

EMISIA SA, 2024. COPERT 5 v 5.7.3, Computer programme to calculate emissions from road transport, January, 2024. <http://www.emisia.com/copert/> .

ENAC/MIT, several years. Annuario Statistico. Ente Nazionale per l'Aviazione Civile, Ministero delle Infrastrutture e dei Trasporti.

ENEA, several years. Rapporto Energia Ambiente. Ente per le Nuove tecnologie, l'Energia e l'Ambiente, Roma.

ENEA, 2000. Il riciclaggio delle batterie al piombo-acido esauste.

ENEA-AIB-MATT, 2002. "Valutazione delle emissioni di inquinanti organici persistenti da parte dell'industria metallurgica secondaria".

ENEA-federAmbiente, 2012. Rapporto sul recupero energetico da rifiuti urbani in Italia. 3° ed.

ENI, several years [a]. Health Safety Environment report. ENI.

ENI, several years [b]. Personal communications.

EUROCONTROL, several years. EUROCONTROL Fuel and Emissions Inventory. Personal Communication to EU Member States. Last communication November 2017.

FEDERACCAI, several years. La siderurgia in cifre. Federazione Imprese Siderurgiche Italiane.

FEDERAMBIENTE, 1998. Impianti di smaltimento: analisi sui termocombustori RSU – prima edizione. Indagine a cura di Motawi A.

Gerardi V., Perrella G., 2001 - I consumi energetici di biomasse nel settore residenziale in Italia nel 1999. ENEA, Ente Nazionale per l'Energia e l'Ambiente, 2001, ENEA RT/ERG/2001/07.

Giordano R., 2007. Trasporto merci: criticità attuali e potenziali sviluppi nel contesto europeo. National road transporters central committee.

INNOVHUB, 2016. Studio comparativo sulle emissioni di apparecchi a gas, GPL, gasolio e pellet. INNOVHUB, Stazioni sperimentali per l'industria.

INNOVHUB, 2018. Fuel consumption, regulated and unregulated exhaust emission tests on five Euro 6 b/c bifuel LPG passenger cars.

INNOVHUB, 2021. Final report on emissions from biomass combustion for heating.

INNOVHUB, several years. Report on the physico-chemical characterization of fossil fuels used in Italy.

IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009.

ISPRA, 2022. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. ISPRA, 369/2022. URL: <https://emissioni.sina.isprambiente.it/inventari-locali/> .

ISPRA, 2024[a]. Italian Greenhouse Gas Inventory 1990-2022. National Inventory Report 2024. March 2024, <http://emissioni.sina.isprambiente.it/inventario-nazionale/> .

ISPRA, 2024[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2024. March 2024, <http://emissioni.sina.isprambiente.it/inventario-nazionale/> .

ISPRA, 2023 [c]. Aggiornamento delle modalità di calcolo delle emissioni navali con particolare riferimento all'ambito portuale a livello nazionale e locale, Rapporti 382/2023

ISPRA, 2023 [d]. Update of estimates of atmospheric emissions from integrated plant for non ferrous metals production. Technical note n.1/2023.

ISPRA, several years. Fuel Quality Monitoring Annual Report.

ISTAT, 2009. Personal communication.

ISTAT, 2014. I consumi energetici delle famiglie, 2013. Nota metodologica. Istituto Nazionale di Statistica www.istat.it .

ISTAT, several years [a]. Annuario Statistico Italiano. Istituto Nazionale di Statistica

ISTAT, several years [b]. Trasporto merci su strada. Istituto Nazionale di Statistica. http://www3.istat.it/dati/dataset/20110729_00/ .

JRC, 2013. JRC reference report. Best Available Techniques (BAT) Reference Document for Iron and Steel Production. Industrial Emissions Directive 2010/75/EU (Integrated Pollution Prevention and Control). Report EUR 25521 EN.

Katsis P., Mellios G., Ntziachristos L., 2012. Description of new elements in COPERT 4 v 10.0, December 2012.

Kouridis C., Gkatzoflias D., Kioutsioukis I., Ntziachristos L., Pastorello C., Dilara P., 2010. Uncertainty Estimates and Guidance for Road Transport Emission Calculations, European Commission, Joint Research Centre, Institute for Environment and Sustainability, 2010, <http://www.emisia.com/docs/COPERT%20uncertainty.pdf> .

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

MIT, several years. Conto Nazionale delle Infrastrutture e dei Trasporti (CNT). Ministero delle Infrastrutture e dei Trasporti. http://www.mit.gov.it/mit/site.php?o=vc&lm=2&id_cat=148 .

MASE, several years [a]. Bilancio Energetico Nazionale (BEN). Ministero dell'Ambiente e della Sicurezza Energetica. <https://dgsaie.mise.gov.it/ben.php> .

MASE, several years [b]. Bollettino Petrolifero Trimestrale (BPT). Ministero dello sviluppo economico. <https://dgsaie.mise.gov.it/bollettino-petrolifero> .

Riva A., 1997. Methodology for methane emission inventory from SNAM transmission system. Snam Spa Italy.

Romano D., Gaudio D., De Lauretis R., 1999. Aircraft Emissions: a comparison of methodologies based on different data availability. Environmental Monitoring and Assessment. Vol. 56 pp. 51-74.

SCENARI/ISPRA, 2013. Indagine sull'uso e la disponibilità delle biomasse in Italia. 2013.

Sempos I., 2018. Note on fossil carbon content in biofuels. IPCC Working Group I, 10 October 2018.

SNAM, several years. Bilancio di sostenibilità.

SSC, 2012. Final report on emissions from biomass combustion for heating. Stazione Sperimentale dei Combustibili. Prot. ENEA/2009/34883/APU-UGA March 2012.

TECHNE, 2009. Stima delle emissioni in atmosfera nel settore del trasporto aereo e marittimo. Final report. TECHNE Consulting, March 2009.

TERNA, several years. Dati statistici sugli impianti e la produzione di energia elettrica in Italia, Gestore Rete Trasmissione Nazionale. www.terna.it .

Trozzi C., Vaccaro R., De Lauretis R., Romano D., 2002 [a]. Air pollutant emissions estimate from global air traffic in airport and in cruise: methodology and case study. Presented at Transport and Air Pollution 2002.

Trozzi C., Vaccaro R., De Lauretis R., 2002 [b]. Air pollutant emissions estimate from global ship traffic in port and in cruise: methodology and case study. Presented at Transport and Air Pollution 2002.

UCINA, several years. La nautica in cifre. Analisi annuale del mercato. Unione Nazionale Cantieri Industrie Nautiche ed Affini.

UNEM, several years. Previsioni di domanda energetica e petrolifera in Italia. Unione Energia per la Mobilità.

12.4 IPPU - INDUSTRIAL PROCESSES (NFR SECTOR 2)

APAT, 2003. Il ciclo industriale dell'acciaio da forno elettrico. Agenzia per la Protezione dell'Ambiente e per i servizi tecnici, Rapporti 38/2003.

AITEC, 2017. Relazione annuale 2017. Associazione italiana tecnico economica del cemento. www.aitecweb.com.

CORINAIR, 1994. Default emission factors handbook. EUR 12586/2, Belgium.

EEA, 2017 [a]. Final Review Report. 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy 30 November 2017.

EEA, 2018. Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report 2018, 30 November 2018.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

EMEP/CORINAIR, 2006. Atmospheric Emission Inventory Guidebook. Technical report No 11/2006.

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2013. Air Pollutant Emission Inventory Guidebook. Technical report n. 12/2013.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

ENEA-AIB-MATT, 2002. "Valutazione delle emissioni di inquinanti organici persistenti da parte dell'industria metallurgica secondaria".

EUROCHLOR, 1998. Mercury process for making chlorine.

EUROCHLOR, 2001. Reduction of Mercury Emissions from the West European Chlor-Alkali Industry, June 2001 available on the web <http://www.chem.unep.ch/mercury/2001-ngo-sub/eurochlor/sub1ngoatt8.pdf>

FEDERACCIAI, several years. La siderurgia in cifre. Federazione Imprese Siderurgiche Italiane.

ILVA, 1997. Personal communication.

IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook.

IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.

IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.

IPPC, 2001. Best Available Techniques Reference Document on the Production of Iron and Steel. Integrated Pollution Prevention and Control. European Commission. December 2001.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. <http://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale>.

ISPRA, 2022. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. ISPRA, 369/2022. URL: <https://emissioni.sina.isprambiente.it/inventari-locali/>.

ISPRA, 2023. Update of estimates of atmospheric emissions from integrated plant for non ferrous metals production. Technical note n.1/2023.

ISTAT, several years. Annuario Statistico Italiano. Istituto Nazionale di Statistica.

Legambiente, 2007. Lo stato dell'arte sulle riconversioni degli impianti cloro soda in Italia, Ottobre 2007 available on the web: http://risorse.legambiente.it/docs/Lo_stato_dell_arte_sulle_riconversioni_degli_impianti_cloro-soda_in_Italia_ott_2007.0000001526.pdf

MASE, several years [b]. Bollettino Petrolifero Trimestrale (BPT). Ministero dello sviluppo economico. <https://dgsaie.mise.gov.it/bollettino-petrolifero> .

MATTM, 2011. Autorizzazione integrata ambientale per l'esercizio dello stabilimento siderurgico della società ILVA s.p.a. ubicato nel comune di Taranto. IPPC permit.

MSE, several years. Statistiche produzione cementi. <http://www.sviluppoeconomico.gov.it/index.php/it/per-i-media/statistiche/2009708-statistiche-produzione-cementi> .

TNO, 1992. Emission factors manual PARCOM – ATMOS. Emission factors for air pollutants 1992.

12.5 IPPU - SOLVENT AND OTHER PRODUCT USE (NFR SECTOR 2)

ACI, several years. Annuario statistico. Automobile Club d'Italia, Roma. <http://www.aci.it/index.php?id=54>

AIA, several years [a]. Personal Communication. Associazione Italiana Aerosol.

AIA, several years [b]. Relazioni annuali sulla produzione italiana aerosol. Associazione Italiana Aerosol.

Assocasa, several years. Personal Communication.

AVISA, several years. Personal Communication.

CEPE, 1999. European guidelines for car repairing.

EC, 1999. Council Directive 1999/13/EC of 11 March 1999 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain activities and installations.

EC, 2002. Screening study to identify reduction in VOC emissions due to the restrictions in the VOC content of products. Final Report of the European Commission, February 2002.

EC, 2004. Directive 2004/42/ce of the european parliament and of the council of 21 April 2004 on the limitation of emissions of volatile organic compounds due to the use of organic solvents in certain paints and varnishes and vehicle refinishing products and amending Directive 1999/13/EC.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.

EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 21/2016.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.

ENEA, 1997. Piano nazionale di tutela della qualità dell'aria. Technical report produced for the Italian Ministry of Environment in the framework of a national project on air quality issue.

ENEA/USLRMA, 1995. Lavanderie a secco.

FAO, several years. Food balance. <http://faostat.fao.org> .

Federchimica, several years. Personal Communication.

FIAT, several years. Rendiconto Ambientale. Gruppo Fiat.

GIADA, 2006. Progetto Giada and Personal Communication. ARPA Veneto – Provincia di Vicenza.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html

ISTAT, several years [a]. Annuario Statistico Italiano.

ISTAT, several years [b]. Bollettino mensile di statistica.

ISTAT, several years [c]. Statistica annuale della produzione industriale. <http://www.istat.it/it/archivio/73150>

MICA, 1999. L'industria Italiana delle vernici. Ministero dell'Industria del Commercio e dell'Artigianato. Dicembre 1999.

Offredi P., several years. Professione Verniciatore del Legno. Personal communication.

Regione Campania, 2005. Inventario regionale delle emissioni di inquinanti dell'aria della Regione Campania, marzo 2005.

Regione Toscana, 2001. Inventario regionale delle sorgenti di emissione in aria ambiente, febbraio 2001.

Techne, 1998. Personal communication.

Techne, 2004. Progetto MeditAiraneo. Rassegna dei fattori di emissione nazionali ed internazionali relativamente al settore solventi. Rapporto Finale, novembre 2004.

Techne, 2008. Fattori di emissione per l'utilizzo di solventi. Rapporto Finale, marzo 2008.

UNIPRO, several years. Rapporto Annuale - Consumi cosmetici in Italia.

Vetrella G., 1994. Strategie ottimali per la riduzione delle emissioni di composti organici volatili. Thesis in Statistics.

12.6 AGRICULTURE (NFR SECTOR 3)

APAT, 2005. Methodologies used in Italy for the estimation of air emission in the agriculture sector. Technical report 64/2005. Rome - Italy. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/methodologies-used-in-italy-for-the-estimation-of>.

Bassanino M., Sacco D., Zavattaro L., Grignania C., 2011. Nutrient balance as a sustainability indicator of different agro-environments in Italy. *Ecol Indic* 11(2): 715-723.

Bechini L., Castoldi N., 2009. On-farm monitoring of economic and environmental performances of cropping systems: Results of a 2-year study at the field scale in northern Italy. *Ecol Indic* 9: 1096-1113.

Berdowski J.J.M., Baas J, Bloos J.P.J., Visschedijk A.J.H., Zandveld P.Y.J., 1997. The European Atmospheric Emission Inventory for Heavy Metals and Persistent Organic Pollutants. Umweltforschungsplan des Bundesministers für Umwelt, Naturschutz und Reaktorsicherheit Luftreinhaltung. Forschungsbericht 104 02 672/03. TNO, Apeldoorn, The Netherlands.

Bittman S. et al, 2014. Bittman, S., Dedina, M., Howard C.M., Oenema, O., Sutton, M.A., (eds), 2014. Options for Ammonia Mitigation: Guidance from the UNECE Task Force on Reactive Nitrogen, Centre for Ecology and Hydrology, Edinburgh, UK www.clrtap-tfrn.org.

Bonazzi G., Crovetto M., Della Casa G., Schiavon S., Sirri F., 2005. Evaluation of Nitrogen and Phosphorus in Livestock manure: Southern Europe (Italy), presentato al Workshop: Nutrients in livestock manure, Bruxelles, 14 February 2005.

Cóndor R.D., Di Cristofaro E., De Lauretis R., 2008. Agricoltura: inventario nazionale delle emissioni e disaggregazione provinciale. Istituto superiore per la protezione e la ricerca ambientale, ISPRA Rapporto tecnico 85/2008. Roma, Italia. <https://www.isprambiente.gov.it/it/archivio/notizie-e-novita-normative/notizie-ispra/anno-2009/rapporto-agricoltura>.

Cóndor R.D., 2011. Agricoltura: emissioni nazionali in atmosfera dal 1990 al 2009. Istituto superiore per la protezione e la ricerca ambientale (ISPRA). Rapporto ISPRA 140/2011. Roma, Italia. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/agricoltura-emissioni-nazionali-in-atmosfera-dal>.

Cóndor R.D., Valli L., 2011. Emissioni nazionali di ammoniaca e scenari emissivi derivanti dalla fase di spandimento agronomico e all'uso dei fertilizzanti azotati in Italia. e-book edited by the Centro Ricerche Produzioni Animali (CRPA). http://www.crpa.it/media/documents/crpa_www/Pubblicazi/E-book/Ammoniaca2011/EmissioniAmmoniaca.pdf.

CREA, 2023. Provision of data on green manure areas by crop for the years 2019-2021, extracted from the FADN database by Dr Antonio Giampaolo in January 2024.

CREA, 2021. Accordo di collaborazione (ex art. 15 L. 241/90) tra ISPRA e CREA - Rapporto tecnico. Valutazione delle emissioni connesse all'uso dei fertilizzanti azotati nell'ambito della direttiva NEC. Novembre 2021.

CRPA, 1997. Piani Regionali di Risanamento e tutela della qualità dell'aria. Quadro delle azioni degli enti locali per il settore zootecnico delle aree padane. Relazione di dettaglio sulla metodologia adottata per la quantificazione delle emissioni di ammoniaca. Febbraio 1997.

CRPA, 2000. Aggiornamento dell'inventario delle emissioni in atmosfera di ammoniaca, metano e protossido di azoto dal comparto agricolo. Centro Ricerche Produzioni Animali. Gennaio 2000.

CRPA, 2006[a]. Progetto MeditAIRaneo: settore Agricoltura. Relazione finale. Technical report on the framework of the MeditAIRaneo project for the Agriculture sector, Reggio Emilia -- Italy.

CRPA, 2006[b]. Predisposizione di scenari di emissione finalizzati alla progettazione di interventi per la riduzione delle emissioni nazionali di ammoniaca ed alla valutazione di misure e di progetti per la tutela della qualità dell'aria a livello regionale. Final report. Reggio Emilia -- Italy.

CRPA, 2010[a]. Personal communication - experts Laura Valli and Maria Teresa Pacchioli from the Research Centre on Animal Production (expert consultation on N excretion and national production systems). Reggio Emilia, Italy.

CRPA, 2010[b]. Valutazione dell'entità delle emissioni ammoniacali derivanti dall'applicazione al suolo dei fertilizzanti, delle loro possibilità di riduzione e individuazione degli elementi per un monitoraggio statistico delle tecniche di applicazione utilizzate. Report. Reggio Emilia – Italy.

CRPA, 2018. Studio per la valutazione degli effetti sulle emissioni delle trasformazioni in corso nel settore degli allevamenti. Report. Reggio Emilia – Italy.

EC, 1986. Council Directive 86/278/EC. Council Directive 86/278/EC of 12 June 1986 on the protection of the environment, and in particular of the soil, when sewage sludge is used in agriculture. Official Journal of the European Communities 4 July 1986.

EC, 2003. Integrated Pollution Prevention and Control (IPPC). Reference Document on Best Available Techniques for Intensive Rearing of Poultry and Pigs. European Commission, July 2003.

EEA, 2016. Analysis of key trends and drivers in greenhouse gas emissions in the EU between 1990 and 2014. Technical paper, June 2016. <http://www.eea.europa.eu/publications/analysis-of-key-trends-ghg/>.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report, 22 November 2019.

EEA, 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive). Final Review Report 2020, 20 November 2020.

-
- EMEP/CORINAIR, 2006. Atmospheric Emission Inventory Guidebook. Technical report No 11/2006.
- EMEP/EEA, 2023. Air Pollutant Emission Inventory Guidebook 2023. Technical guidance to prepare national emission inventories. EEA report 06/2023.
- ENEA, 2003. Emissione di ammoniaca e di composti ad effetto serra dagli allevamenti di galline ovaiole. Fattori di emissione e tecniche di riduzione. RT/2003/65/PROT. Italy.
- EPA, 1995. AP-42 Compilation of Air Emission Factors, 5th edition January 1995. <https://www.epa.gov/air-emissions-factors-and-quantification/ap-42-compilation-air-emission-factors>.
- Giardini L., 1983. Agronomia Generale, Patron, Bologna.
- Gislon G., Colombini S., Borreani G., Crovetto G. M., Sandrucci A., Galassi G., Tabacco E., Rapetti L., 2020. Milk production, methane emissions, nitrogen, and energy balance of cows fed diets based on different forage systems. *J. Dairy Sci.* 103:8048–8061 <https://doi.org/10.3168/jds.2019-18134>.
- GU, Gazzetta Ufficiale della Repubblica Italiana, 2006. Criteri e norme tecniche generali per la disciplina regionale dell'utilizzazione agronomica degli effluenti di allevamento e di acque reflue di cui all'articolo 38 del decreto legislativo 11 maggio 1999 N. 152. G.U. n. 109 del 12/05/06 - Suppl. Ordinario n.120. Ministero delle Politiche Agricole e Forestali. Italy. <http://www.gazzettaufficiale.it/>.
- GU, Gazzetta Ufficiale della Repubblica Italiana, 2016. Criteri e norme tecniche generali per la disciplina regionale dell'utilizzazione agronomica degli effluenti di allevamento e delle acque reflue, nonché per la produzione e l'utilizzazione agronomica del digestato. G.U. n. 90 del 18/04/16 - Suppl. Ordinario n. 9. Ministero delle Politiche Agricole Alimentari e Forestali. Italy. <http://www.gazzettaufficiale.it/>.
- IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook.
- IPCC, 2006. 2006 IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the National Greenhouse Gas Inventories Programme, Eggleston H.S., Buendia L., Miwa K., Ngara T. and Tanabe K. (eds). Published: IGES, Japan.
- ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale>.
- ISPRA, 2021. Database della disaggregazione a livello provinciale dell'Inventario nazionale delle emissioni:1990-1995-2000-2005-2010-2015-2019. Istituto Superiore per la Protezione e la Ricerca Ambientale, ISPRA. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- ISPRA, 2022. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni, Rapporto 369/2022.<https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale-dell2019inventario-nazionale-delle-emissioni>.
- ISPRA, several years [a]. Italian Greenhouse Gas Inventory - National Inventory Report. <http://emissioni.sina.isprambiente.it/inventario-nazionale/>.
- ISPRA, several years [b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory, <http://emissioni.sina.isprambiente.it/inventario-nazionale/>.
- ISPRA, several years [c]. Annuario dei dati ambientali. <http://annuario.isprambiente.it/>.
- ISPRA, several years [d]. Serie storiche delle emissioni nazionali di inquinanti atmosferici, Rete del Sistema Informativo Nazionale Ambientale - SINANET. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>.
- Jenkins B.M. (1996a). 'Atmospheric Pollutant Emission Factors from Open Burning of Agricultural and Forest Biomass by Wind Tunnel Simulations'. Final report (3 Vols.). CARB Project A932-126, California Air Resources Board, Sacramento, California.
- MASE, several years [b]. Personal communication: E-mail request for sewage sludge applied to agricultural soils in Italy. Ministero dell'Ambiente e della Tutela del Territorio e del Mare, Roma -Italia.

Nicholson F.A., Chambers B.J., Walker A.W., 2004. Ammonia emissions from broiler litter and laying hen manure management system, *Biosystems Engineering* (2004) 89 (2), 175-185.

Perelli, M., 2007. Prezzi dei prodotti agricoli e fertilizzazione. *Fertilizzanti Maggio 2007*. Anno IX N3. 10-13pp.

Regione Emilia-Romagna, Servizio sviluppo sistema agroalimentare, (2001) L. R. 28/98 – p.s.a. 2000 - n. Prog. 3 tab. D - Tecniche di riduzione delle emissioni in atmosfera originate dagli allevamenti zootecnici, a cura di CRPA, Settembre 2001.

Regione Emilia Romagna, 2004 L. R. 28/98 – P.S.A. 2001 - N. PROG. 3 TAB. B3 - Bilancio dell'azoto nelle specie di interesse zootecnico, Relazione finale, a cura di C.R.P.A., September 2004, Reggio Emilia, Italy.

Xiccato G., Schiavon S., Gallo L., Bailoni L., Bittante G., 2005. Nitrogen excretion in dairy cow, beef and veal cattle, pig, and rabbit farms in Northern Italy. *Ital. J.Anim.Sci.* Vol. 4 (Suppl.), 103-111.

12.7 WASTE (NFR SECTOR 5)

Aasestad, 2007. The Norwegian Emission Inventory 2007. Documentation of methodologies for estimating emissions of greenhouse gases and long-range transboundary air pollutants.

AMA-Comune di Roma, 1996. Nuovo impianto per l'incenerimento dei rifiuti ospedalieri. Rapporto AMA.

ANPA, 1998. Il sistema ANPA di contabilità dei rifiuti, prime elaborazioni dei dati. Agenzia Nazionale per la Protezione dell'Ambiente.

APAT-ONR, several years. Rapporto Rifiuti. Agenzia per la Protezione dell'Ambiente e per i servizi Tecnici.

AUSITRA-Assoambiente, 1995. Impianti di trattamento dei rifiuti solidi urbani e assimilabili. Indagine a cura di Merzagora W., Ferrari S.P.

Borgioli E., 1981. Nutrizione e alimentazione degli animali domestici. Ed Agricole, p. 464.

CESTAAT, 1988. Impieghi dei sottoprodotti agricoli ed agroindustriali, Vol. 1. Centro Studi sull'Agricoltura, l'Ambiente e il Territorio, edizione fuori commercio, p. 311.

CREA, 2017. Fornitura dati meteo-climatici georeferenziati nell'ambito della collaborazione CREA-AA/ISPRA. CREA - Consiglio per la Ricerca in Agricoltura e l'Analisi dell'Economia Agraria Centro di ricerca Agricoltura e Ambiente (CREA-AA), delivery data mail 19/10/2017.

CRPA, 2018. Studio per la valutazione degli effetti sulle emissioni delle trasformazioni in corso nel settore degli allevamenti. Report. Reggio Emilia – Italy

De Stefanis P., 1999. Personal communication.

De Stefanis P., 2012. Personal communication (mail 16 November 2012)

EC, 1999. Council Directive 1999/31/EC. Council Directive 99/31/EC of 26 April 1999 on the landfill of waste. *Official Journal of the European Communities* 16 July 1999.

EEA, 2017 [a]. Final Review Report. 2017 Comprehensive Technical Review of National Emission Inventories pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Italy 30 November 2017.

EEA, 2017 [b]. Final Review Report. 2017 annual review of national greenhouse gas inventory data pursuant to Article 19(2) of Regulation (EU) No 525/2013. Italy 30 June 2017.

EEA, 2018. Second phase of review of national air pollution emission inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284). Final Review Report 2018, 30 November 2018.

EEA, 2019. Third phase of review of national air pollution inventory data pursuant to the Directive on the Reduction of National Emissions of Certain Atmospheric Pollutants (Directive (EU) 2016/2284 or 'NECD') Italy. Draft Review Report 2019, 18 October 2019.

-
- EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.
- EEA, 2021. Final Review Report, 2021 Review of National Air Pollutant Emission Inventory Data 2021 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 29 October 2021. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.
- EEA, 2022. Final Review Report, 2022 Review of National Air Pollutant Emission Inventory Data 2022 under Directive 2016/2284 (National Emission reduction Commitments Directive). Italy. 11 November 2022. Reference: Service request No. 070201/2019/8159797/SER/ENV.C.3 Umweltbundesamt Wien.
- EEA, 2023. Final Review Report, Review of National Air Pollutant Emission Inventory Data 2023 under Directive (EU) 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/815979/SER/ENV.C.3. Umweltbundesamt Wien.
- EMEP/CORINAIR, 2007. Atmospheric Emission Inventory Guidebook. Technical report No 16/2007.
- EMEP/EEA, 2016. Air Pollutant Emission Inventory Guidebook. Technical report n. 21/2016.
- EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.
- ENEA-federAmbiente, 2012. Rapporto sul recupero energetico da rifiuti urbani in Italia. 3° ed.
- Favoino E., Cortellini L., 2001. Composting and biological treatment in southern European countries: an overview. Conference Proceedings Soil and Biowaste in Southern Europe. Rome 18-19 January, 2001.
- Favoino E., Girò F., 2001. An assessment of effective, optimised schemes for source separation of organic waste in Mediterranean districts. Conference Proceedings Soil and Biowaste in Southern Europe. Rome 18-19 January, 2001.
- FEDERAMBIENTE, 1998. Impianti di smaltimento: analisi sui termocombustori RSU – prima edizione. Indagine a cura di Motawi A.
- Finn L., Spencer R., 1997. Managing biofilters for consistent odor and VOC treatment. Biocycle, January 1997 Vol. 38 Iss.1.
- Gaudio et al., 1993. Emissioni in atmosfera dalle discariche di rifiuti in Italia. RS, Rifiuti Solidi vol. VII n. 5, Sept.-Oct. 1993.
- IIASA, 2004. Interim Report IR-04-079. Primary Emissions of Submicron and Carbonaceous Particles in Europe and the Potential for their Control.
- IPCC, 1997. Revised 1996 IPCC Guidelines for National Greenhouse Gas Emission Inventories. Three volumes: Reference Manual, Reporting Manual, Reporting Guidelines and Workbook. IPCC/OECD/IEA. IPCC WG1 Technical Support Unit, Hadley Centre, Meteorological Centre, Meteorological Office, Bracknell, UK.
- IPCC, 2000. Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories. IPCC National Greenhouse Gas Inventories Programme, Technical Support Unit, Hayama, Kanagawa, Japan.
- ISPRA, 2017. Update of CH4 emission factor from composting. Technical note n.1/2017.
- ISPRA, 2024[a]. Italian Greenhouse Gas Inventory 1990-2022. National Inventory Report 2024. March 2024, <https://emissioni.sina.isprambiente.it/inventario-nazionale/> .
- ISPRA, several years [a]. Rapporto Rifiuti. Istituto Superiore per la Protezione e la Ricerca Ambientale. <http://www.isprambiente.gov.it/it/pubblicazioni/rapporti>
- ISTAT, several years [a]. Statistiche dell'agricoltura, zootecnia e mezzi di produzione - Annuari (1990-1993), Istituto Nazionale di Statistica.
- ISTAT, several years [b]. Statistiche dell'agricoltura - Annuari (1994-2000), Istituto Nazionale di Statistica.

ISTAT, 2015. Censimento delle acque per uso civile. Istituto nazionale di statistica, also available at website <http://www.istat.it>.

ISTAT, 2017[a]. Dati congiunturali sulle coltivazioni 2014. Istituto Nazionale di Statistica. <http://agri.istat.it/jsp/Introduzione.jsp> .

ISTAT, 2017[b]. Personal communication with R. Moro: E-mail request for last updated information on agricultural surface and production, year 2015. Istituto Nazionale di Statistica.

MASE, several years [a]. RSA - Rapporto sullo stato dell'ambiente 1989, 1992, 1997, 2001. Ministero dell'Ambiente e della Tutela del Territorio e del Mare.

Ministerial Decree 19 November 1997, n. 503. Regolamento recante norme per l'attuazione delle Direttive 89/369/CEE e 89/429/CEE concernenti la prevenzione dell'inquinamento atmosferico provocato dagli impianti di incenerimento dei rifiuti urbani e la disciplina delle emissioni e delle condizioni di combustione degli impianti di incenerimento di rifiuti urbani, di rifiuti speciali non pericolosi, nonché di taluni rifiuti sanitari. G.U. 29 gennaio 1998, n. 23.

Ministerial Decree 12 July 1990. Linee Guida per il contenimento delle emissioni inquinanti degli impianti industriali e la fissazione dei valori minimi di emissione. G.U. 30 luglio 1990, n. 176.

Pastorelli et al., 2001. Sviluppo di fattori di emissione da inceneritori di rifiuti urbani lombardi e loro applicazione all'inventario nazionale delle diossine. Ingegneria Ambientale, ANNO XXX N.1 January 2001.

SEFIT, several years. Personal Communication with Daniele Fogli: E-mail request for activity data regarding cremation of corpses in Italy.

SEFIT, 2015. Emissioni inquinanti in atmosfera per i crematori italiani. Indagine conoscitiva ed elaborazione dati. Novembre 2015.

Tchobanoglous G. et al., 1993. Tchobanoglous G., Theisen H., Vigil A. Integrated Waste Management, McGraw-Hill, 1993.

US EPA, 1990. Air Emissions Species Manual, vol. I: Volatile Organic Compound Species Profiles, Second Edition. EPA-450/2-90-001a (United States Environmental Protection Agency – Office of Air Quality Planning and Standards, Research Triangle Park, NC 27711), January 1990.

VV.FF., several years. Annually statistics of fire service in Italy, several years.

12.8 OTHER (NFR SECTOR 6)

EMEP/EEA, 2023. Air Pollutant Emission Inventory Guidebook 2023. Technical guidance to prepare national emission inventories. EEA report 06/2023.

FEDIAF, 2022. Facts & Figures 2021. FEDIAF European Pet Food - Annual Report 2022. Statistics | FEDIAF (europeanpetfood.org).

FEDIAF, 2023. Facts & Figures 2022. FEDIAF European Pet Food - Annual Report 2023. Statistics | FEDIAF (europeanpetfood.org).

12.9 RECALCULATIONS AND IMPROVEMENTS

EEA, 2023. Final Review Report, Review of National Air Pollutant Emission Inventory Data 2023 under Directive (EU) 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/815979/SER/ENV.C.3. Umweltbundesamt Wien.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. Technical report n. 13/2019.

EMEP/EEA, 2023. Air Pollutant Emission Inventory Guidebook. Technical report n. 6/2023.

EMISIA SA, 2024. COPERT 5 v 5.7.3, Computer programme to calculate emissions from road transport, January, 2024. <http://www.emisia.com/copert/> .

ISPRA, 2024[b]. Quality Assurance/Quality Control plan for the Italian Emission Inventory. Year 2022. March 2024, <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>

ISPRA, several years. Rapporto Rifiuti. Istituto Superiore per la Protezione e la Ricerca Ambientale. <http://www.isprambiente.gov.it/it/pubblicazioni/rapporti>

12.10 PROJECTIONS

Amann, M., Bertok, J., Borken-Kleefeld, J., Cofala, J., Heyes, C., Höglund-Isaksson, L., Klimont, Z., Nguyen, B., Posch, M., Rafaj, P., Sandler, R., Schöpp, W., Wagner, F., Winiwarter, W., 2011. Cost-effective control of air quality and greenhouse gases in Europe: modelling and policy applications. *Environmental Modeling Software*, 26, 1489–1501.

Briganti, G., Calori, G., Cappelletti, A., Ciancarella, L., D'Isidoro, M., Finardi, S., Vitali, L., 2011. Determination of multi-year atmospheric transfer Matrices for GAINS-Italy model. In: *High Performance Computing on CRESCO Infrastructure: Research Activities and Results*, vol. 2010. pp. 45–51 Report ENEA Cresco, ISBN 978-88-8286-268-8.

D'Elia, I., Peschi, E., 2013. Lo scenario emissivo nazionale nella negoziazione internazionale. ENEA Technical Report, RT/2013/10/ENEA (in Italian). <http://openarchive.enea.it/handle/10840/4505> .

D'Elia, I., Peschi, E., 2016. How National integrated air quality models can be used in defining environmental policies: the revision of the NEC directive. ENEA Technical Report, RT/2016/30/ENEA. <http://openarchive.enea.it/handle/10840/8153> .

D.Lgs., 2010. Decreto Legislativo 13 agosto 2010, n. 155. Attuazione della direttiva 2008/50/CE relativa alla qualità dell'aria ambiente e per un'aria più pulita in Europa (GU n.216 del 15-9-2010 - Suppl. Ordinario n. 217).

D.Lgs., 2018. Decreto Legislativo 30 maggio 2018, n. 81. Attuazione della direttiva (UE) 2016/2284 del Parlamento europeo e del Consiglio, del 14 dicembre 2016, concernente la riduzione delle emissioni nazionali di determinati inquinanti atmosferici, che modifica la direttiva 2003/35/CE e abroga la direttiva 2001/81/CE (GU n. 151 del 2-7-2018).

EC, 2008. Directive 2008/50/EC of the European parliament and of the council of 21 may 2008 on ambient air quality and a cleaner air for Europe. *Official Journal of the European Union L*. 152 of 11.06.2008.

EU, 2016. Directive (EU) 2016/2284 of the European Parliament and of the Council of 14 December 2016 on the reduction of national emissions of certain atmospheric pollutants, amending Directive 2003/35/EC and repealing Directive 2001/81/EC. *Official Journal of the European Union L*. 344/1 of 17.12.2016.

IIR, 2021. Italian Emission Inventory 1990-2019. Informative Inventory Report 2021. ISPRA Technical Report 342/2021. ISBN 978-88-448-1047-4

Loulou, R., Goldstein, G., Noble, K., 2004. Documentation for the MARKAL Family of Models. ETSAP.

Loulou, R., Remne, U., Kanudia, A., Lehtila, A., Goldstein, G., 2005. Documentation for the TIMES Model - PART I 1–78. <https://iea-etsap.org/index.php/documentation>

Mircea, M., Ciancarella, L., Briganti, G., Calori, G., Cappelletti, A., Cionni, I., Costa, M., Cremona, G., D'Isidoro, M., Finardi, S., Pace, G., Piersanti, A., Righini, G., Silibello, C., Vitali, L., Zanini, G., 2014. Assessment of the AMS-MINNI system capabilities to predict air quality over Italy for the calendar year 2005. *Atmospheric Environment*, 84, 178–188, ISSN 1352-2310, <http://dx.doi.org/10.1016/j.atmosenv.2013.11.006>

UNECE, 2015. Guidelines for reporting emissions and projections data under the Convention on Long-range Transboundary Air Pollution. ECE/EB.AIR/128/2015. <https://www.unece.org/fileadmin/DAM/env/documents/2015/AIR/EB/English.pdf>

12.11 REPORTING OF GRIDDED EMISSIONS AND LPS

CEIP, 2019. Updated documentation of the EMEP gridding system v.2019 for the spatial disaggregation of emission data with a resolution of 0.1°x0.1°(long-lat). Technical report CEIP 06/2019.

EEA, 2020. Final Review Report 2020. Review of National Air Pollutant Emission Inventory Data 2020 under Directive 2016/2284 (National Emission reduction Commitments Directive) Service Contract No. 070201/2019/8159797/SER/ENV.C.3. Italy, 20 November 2020.

EMEP/EEA, 2019. Air Pollutant Emission Inventory Guidebook. EEA. Technical report No 13/2019.

ISPRA, 2009. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni. Anni 1990-1995-2000-2005. ISPRA, 92/2009. URL: http://www.apat.gov.it/site/it-IT/APAT/Pubblicazioni/Rapporti/Documento/rapporti_92_2009.html.


ISPRA, 2021. Database della disaggregazione a livello provinciale dell'Inventario nazionale delle emissioni:1990-1995-2000-2005-2010-2015-2019. Istituto Superiore per la Protezione e la Ricerca Ambientale, ISPRA. <http://emissioni.sina.isprambiente.it/serie-storiche-emissioni/>

ISPRA, 2022. La disaggregazione a livello provinciale dell'inventario nazionale delle emissioni, Rapporto 369/2022. <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/la-disaggregazione-a-livello-provinciale-dell2019inventario-nazionale-delle-emissioni>.

APPENDIX 1 SUMMARY INFORMATION ON CONDENSABLE IN PM

In order to improve atmospheric modelling and support the design of efficient and relevant policy for reducing the levels of air pollutants, emission inventory data need to be complete, accurate and comparable. With this aim, Italy immediately accepted the EMEP proposal on the necessity of accounting for condensable in PM emissions and generally applies these emission factors to all the categories. Of course, for certain categories is not possible to define if the emission factors includes condensable or not, as reported also in the 2019 Guidebook EMEP/EEA, consequently it is hard to fill the following table at category level but it is possible to provide more information. In particular, Italy uses emission factors with condensable for PM emissions from road transport thanks to the Copert model and in domestic and residential heating thanks several studies carried on in the last years about heating appliances, burning wood or other fuels. In particular, as concerns emissions from small combustion, a paper discussed during the 2019 meeting of TFEIP reported some independent verification on these estimates and Italy resulted in a good comparability with estimates of TNO (see Figure A1 1).

Figure A1 1 Comparison between TNO and countries estimates.



APPENDIX 2 UNCERTAINTY ASSESSMENT

In the following table detailed results, concerning NOx emissions, are reported for 2021 and for the trend (data submitted in 2023). In table A2.2 results for SOx 2022 emissions have been reported.

Table A2.1 Results of the uncertainty analysis (Approach 1) for NOx. 2021 and trend assessment.

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Energy	1A1a - Public electricity and heat production	NOx	408.63	25.06	10.0	10.0	14.1	0.337	-93.9	0.6
Energy	1A1b - Petroleum refining	NOx	37.26	7.97	10.0	10.0	14.1	0.034	-78.6	0.0
Energy	1A1c - Manufacture of solid fuels and other energy industries	NOx	8.59	2.17	10.0	10.0	14.1	0.003	-74.8	0.0
Energy	1A2a - Stationary combustion in manufacturing industries and construction: Iron and steel	NOx	21.51	6.30	10.0	10.0	14.1	0.021	-70.7	0.0
Energy	1A2b - Stationary combustion in manufacturing industries and construction: Non-ferrous metals	NOx	2.85	0.82	10.0	10.0	14.1	0.000	-71.4	0.0
Energy	1A2c - Stationary combustion in manufacturing industries and construction: Chemicals	NOx	55.87	2.95	10.0	10.0	14.1	0.005	-94.7	0.1
Energy	1A2d - Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	NOx	2.84	4.42	10.0	10.0	14.1	0.010	55.4	0.0
Energy	1A2e - Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	NOx	11.13	1.77	10.0	10.0	14.1	0.002	-84.1	0.0
Energy	1A2f - Stationary combustion in manufacturing industries and construction: Non-metallic minerals	NOx	120.64	31.70	10.0	10.0	14.1	0.539	-73.7	0.0
Energy	1A2gvii - Mobile Combustion in manufacturing industries and construction: (please specify in the IIR)	NOx	30.76	4.54	20.0	20.0	28.3	0.044	-85.2	0.1
Energy	1A2gviii - Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	NOx	35.79	1.66	10.0	10.0	14.1	0.001	-95.4	0.1
Energy	1A3ai(i) - International aviation LTO (civil)	NOx	1.60	2.33	3.0	10.0	10.4	0.002	45.3	0.0

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Energy	1A3aii(i) - Domestic aviation LTO (civil)	NOx	1.36	1.59	3.0	10.0	10.4	0.001	17.0	0.0
Energy	1A3bi - Road transport: Passenger cars	NOx	590.92	128.06	3.0	10.0	10.4	4.793	-78.3	0.2
Energy	1A3bii - Road transport: Light duty vehicles	NOx	60.60	43.76	3.0	10.0	10.4	0.560	-27.8	0.1
Energy	1A3biii - Road transport: Heavy duty vehicles and buses	NOx	340.33	79.65	3.0	10.0	10.4	1.854	-76.6	0.1
Energy	1A3biv - Road transport: Mopeds & motorcycles	NOx	4.29	2.83	3.0	10.0	10.4	0.002	-34.1	0.0
Energy	1A3c - Railways	NOx	10.27	1.48	3.0	20.0	20.2	0.002	-85.5	0.0
Energy	1A3dii - National navigation (shipping)	NOx	95.55	78.89	2.0	10.0	10.2	1.735	-17.4	0.2
Energy	1A3ei - Pipeline transport	NOx	2.89	0.57	3.0	10.0	10.4	0.000	-80.4	0.0
Energy	1A4ai - Commercial/institutional: Stationary	NOx	11.26	30.85	10.0	10.0	14.1	0.510	173.9	0.2
Energy	1A4bi - Residential: Stationary	NOx	51.53	41.27	10.0	10.0	14.1	0.914	-19.9	0.2
Energy	1A4bii - Residential: Household and gardening (mobile)	NOx	0.02	0.00	10.0	10.0	14.1	0.000	-93.9	0.0
Energy	1A4ci - Agriculture/Forestry/Fishing: Stationary	NOx	0.97	12.94	10.0	20.0	22.4	0.224	1238.9	0.1
Energy	1A4cii - Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	NOx	102.39	23.70	10.0	20.0	22.4	0.753	-76.8	0.1
Energy	1A4ciii - Agriculture/Forestry/Fishing: National fishing	NOx	8.37	6.20	10.0	20.0	22.4	0.052	-25.9	0.0
Energy	1A5b - Other, Mobile (including military, land based and recreational boats)	NOx	11.16	1.70	3.0	50.0	50.1	0.019	-84.8	0.0
Energy	1B2aiv - Fugitive emissions oil: Refining / storage	NOx	5.00	5.33	3.0	20.0	20.2	0.031	6.7	0.0
Energy	1B2c - Venting and flaring (oil, gas, combined oil and gas)	NOx	0.27	0.19	3.0	20.0	20.2	0.000	-29.1	0.0
IPPU	2B1 - Ammonia production	NOx	1.46	0.22	3.0	10.0	10.4	0.000	-84.8	0.0
IPPU	2B2 - Nitric acid production	NOx	3.46	0.27	3.0	10.0	10.4	0.000	-92.3	0.0
IPPU	2B3 - Adipic acid production	NOx	0.02	0.02	3.0	10.0	10.4	0.000	42.8	0.0
IPPU	2B6 - Titanium dioxide production	NOx	0.05	0.04	3.0	10.0	10.4	0.000	-27.9	0.0
IPPU	2B7 - Soda ash production	NOx	0.09	0.14	3.0	10.0	10.4	0.000	58.5	0.0
IPPU	2B10a - Chemical industry: Other (please specify in the IIR)	NOx	16.97	1.78	3.0	10.0	10.4	0.001	-89.5	0.0

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
IPPU	2C1 - Iron and steel production	NOx	2.25	2.59	3.0	10.0	10.4	0.002	15.2	0.0
IPPU	2C2 - Ferroalloys production	NOx	0.01	0.00	3.0	10.0	10.4	0.000	-100.0	0.0
IPPU	2C3 - Aluminium production	NOx	0.50	0.00	3.0	10.0	10.4	0.000	-100.0	0.0
IPPU	2G - Other product use (please specify in the IIR)	NOx	0.17	0.13	5.0	30.0	30.4	0.000	-25.2	0.0
IPPU	2H1 - Pulp and paper industry	NOx	0.08	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
AFOLU	3B1a - Manure management - Dairy cattle	NOx	0.75	0.36	5.0	20.0	20.6	0.000	-51.8	0.0
AFOLU	3B1b - Manure management - Non-dairy cattle	NOx	0.58	0.52	5.0	20.0	20.6	0.000	-11.3	0.0
AFOLU	3B2 - Manure management - Sheep	NOx	0.17	0.13	5.0	20.0	20.6	0.000	-23.0	0.0
AFOLU	3B3 - Manure management - Swine	NOx	0.02	0.02	5.0	20.0	20.6	0.000	2.7	0.0
AFOLU	3B4a - Manure management - Buffalo	NOx	0.03	0.06	5.0	20.0	20.6	0.000	118.9	0.0
AFOLU	3B4d - Manure management - Goats	NOx	0.02	0.02	5.0	20.0	20.6	0.000	-15.7	0.0
AFOLU	3B4e - Manure management - Horses	NOx	0.03	0.04	5.0	20.0	20.6	0.000	27.7	0.0
AFOLU	3B4f - Manure management - Mules and asses	NOx	0.01	0.01	5.0	20.0	20.6	0.000	-13.6	0.0
AFOLU	3B4gi - Manure management - Laying hens	NOx	0.00	0.14	5.0	20.0	20.6	0.000	5711.1	0.0
AFOLU	3B4gii - Manure management - Broilers	NOx	0.15	0.16	5.0	20.0	20.6	0.000	8.7	0.0
AFOLU	3B4giii - Manure management - Turkeys	NOx	0.09	0.06	5.0	20.0	20.6	0.000	-32.7	0.0
AFOLU	3B4giv - Manure management - Other poultry	NOx	0.02	0.03	5.0	20.0	20.6	0.000	63.2	0.0
AFOLU	3B4h - Manure management - Other animals (please specify in IIR)	NOx	0.10	0.07	5.0	20.0	20.6	0.000	-31.9	0.0
AFOLU	3Da1 - Inorganic N-fertilizers (includes also urea application)	NOx	30.38	21.56	20.0	20.0	28.3	0.997	-29.0	0.2
AFOLU	3Da2a - Animal manure applied to soils	NOx	20.93	18.64	20.0	50.0	53.9	2.701	-11.0	0.3
AFOLU	3Da2b - Sewage sludge applied to soils	NOx	0.20	0.30	20.0	50.0	53.9	0.001	49.7	0.0
AFOLU	3Da2c - Other organic fertilisers applied to soils	NOx	0.66	4.40	20.0	50.0	53.9	0.151	566.5	0.1
AFOLU	3Da3 - Urine and dung deposited by grazing animals	NOx	7.07	5.70	20.0	50.0	53.9	0.253	-19.3	0.1
Waste	3F - Field burning of agricultural residues	NOx	0.45	0.47	30.0	50.0	58.3	0.002	3.9	0.0
Waste	5C1a - Municipal waste incineration	NOx	0.46	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
Waste	5C1bi - Industrial waste incineration	NOx	0.43	0.05	10.0	20.0	22.4	0.000	-88.4	0.0
Waste	5C1bii - Hazardous waste incineration	NOx	0.00	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
Waste	5C1biii - Clinical waste incineration	NOx	0.07	0.03	10.0	20.0	22.4	0.000	-61.3	0.0

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Waste	5C1bv - Cremation	NOx	0.06	0.06	10.0	20.0	22.4	0.000	-5.6	0.0
Waste	5C2 - Open burning of waste	NOx	0.01	0.13	10.0	20.0	22.4	0.000	2342.9	0.0
TOTAL			2124.05	610.69			4.1		-71.2	0.9

Table A2.2 Results of the uncertainty analysis (Approach 1) for SO_x. 2022 and trend assessment.

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Energy	1A1a - Public electricity and heat production	SO _x	769.28	6.93	10.0	10.0	14.1	1.235	-99.1	0.2
Energy	1A1b - Petroleum refining	SO _x	192.21	5.44	10.0	10.0	14.1	0.761	-97.2	0.0
Energy	1A1c - Manufacture of solid fuels and other energy industries	SO _x	39.28	1.29	10.0	10.0	14.1	0.043	-96.7	0.0
Energy	1A2a - Stationary combustion in manufacturing industries and construction: Iron and steel	SO _x	20.79	3.17	10.0	10.0	14.1	0.259	-84.7	0.0
Energy	1A2b - Stationary combustion in manufacturing industries and construction: Non-ferrous metals	SO _x	15.07	1.25	10.0	10.0	14.1	0.041	-91.7	0.0

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Energy	1A2c - Stationary combustion in manufacturing industries and construction: Chemicals	SOx	129.09	0.30	10.0	10.0	14.1	0.002	-99.8	0.0
Energy	1A2d - Stationary combustion in manufacturing industries and construction: Pulp, Paper and Print	SOx	4.37	0.04	10.0	10.0	14.1	0.000	-99.0	0.0
Energy	1A2e - Stationary combustion in manufacturing industries and construction: Food processing, beverages and tobacco	SOx	25.47	0.03	10.0	10.0	14.1	0.000	-99.9	0.0
Energy	1A2f - Stationary combustion in manufacturing industries and construction: Non-metallic minerals	SOx	63.55	23.26	10.0	10.0	14.1	13.930	-63.4	0.2
Energy	1A2gvii - Mobile Combustion in manufacturing industries and construction: (please specify in the IIR)	SOx	3.50	0.01	20.0	20.0	28.3	0.000	-99.8	0.0
Energy	1A2gviii - Stationary combustion in manufacturing industries and construction: Other (please specify in the IIR)	SOx	65.75	0.41	10.0	10.0	14.1	0.004	-99.4	0.0
Energy	1A3ai(i) - International aviation LTO (civil)	SOx	0.13	0.30	3.0	5.0	5.8	0.000	131.5	0.0
Energy	1A3aii(i) - Domestic aviation LTO (civil)	SOx	0.11	0.18	3.0	5.0	5.8	0.000	58.2	0.0
Energy	1A3bi - Road transport: Passenger cars	SOx	61.39	0.24	3.0	5.0	5.8	0.000	-99.6	0.0
Energy	1A3bii - Road transport: Light duty vehicles	SOx	15.97	0.05	3.0	5.0	5.8	0.000	-99.7	0.0
Energy	1A3biii - Road transport: Heavy duty vehicles and buses	SOx	49.64	0.11	3.0	5.0	5.8	0.000	-99.8	0.0
Energy	1A3biv - Road transport: Mopeds & motorcycles	SOx	2.29	0.01	3.0	5.0	5.8	0.000	-99.7	0.0
Energy	1A3c - Railways	SOx	1.18	0.00	3.0	5.0	5.8	0.000	-100.0	0.0
Energy	1A3dii - National navigation (shipping)	SOx	77.94	10.91	2.0	5.0	5.4	0.444	-86.0	0.0
Energy	1A3ei - Pipeline transport	SOx	0.01	0.01	3.0	5.0	5.8	0.000	-18.5	0.0
Energy	1A4ai - Commercial/institutional: Stationary	SOx	2.47	4.11	10.0	5.0	11.2	0.272	66.0	0.0
Energy	1A4bi - Residential: Stationary	SOx	72.87	5.08	10.0	5.0	11.2	0.414	-93.0	0.0
Energy	1A4bii - Residential: Household and gardening (mobile)	SOx	0.03	0.00	10.0	5.0	11.2	0.000	-100.0	0.0
Energy	1A4ci - Agriculture/Forestry/Fishing: Stationary	SOx	6.92	0.03	10.0	10.0	14.1	0.000	-99.6	0.0

Inventory sector	Category code and name	Gas	Base year emissions	Year t emissions	Activity data uncertainty	Emission factor /estimation parameter uncertainty	Combined uncertainty	Contribution to variance by category in year t	Inventory trend in national emissions for year t increase with respect to base year	Uncertainty introduced into the trend in total national emissions with respect to base year
					%	%	%	(fraction)	(% of base year)	%
Energy	1A4cii - Agriculture/Forestry/Fishing: Off-road vehicles and other machinery	SOx	13.00	0.03	10.0	10.0	14.1	0.000	-99.8	0.0
Energy	1A4ciii - Agriculture/Forestry/Fishing: National fishing	SOx	1.18	0.00	10.0	10.0	14.1	0.000	-99.8	0.0
Energy	1A5b - Other, Mobile (including military, land based and recreational boats)	SOx	1.19	0.16	3.0	50.0	50.1	0.008	-86.7	0.0
Energy	1B2aiv - Fugitive emissions oil: Refining / storage	SOx	67.40	14.14	3.0	20.0	20.2	10.531	-79.0	0.1
Energy	1B2c - Venting and flaring (oil, gas, combined oil and gas)	SOx	12.32	3.72	3.0	20.0	20.2	0.728	-69.8	0.0
IPPU	2B1 - Ammonia production	SOx	0.06	0.00	3.0	10.0	10.4	0.000	-93.3	0.0
IPPU	2B6 - Titanium dioxide production	SOx	2.60	0.14	3.0	10.0	10.4	0.000	-94.7	0.0
IPPU	2B7 - Soda ash production	SOx	0.12	0.15	3.0	10.0	10.4	0.000	22.0	0.0
IPPU	2B10a - Chemical industry: Other (please specify in the IIR)	SOx	58.36	5.20	3.0	10.0	10.4	0.380	-91.1	0.0
IPPU	2C1 - Iron and steel production	SOx	2.89	1.23	3.0	10.0	10.4	0.021	-57.3	0.0
IPPU	2C2 - Ferroalloys production	SOx	0.01	0.00	3.0	10.0	10.4	0.000	-100.0	0.0
IPPU	2C3 - Aluminium production	SOx	3.50	0.00	3.0	10.0	10.4	0.000	-100.0	0.0
IPPU	2G - Other product use (please specify in the IIR)	SOx	0.01	0.02	5.0	30.0	30.4	0.000	148.0	0.0
IPPU	2H1 - Pulp and paper industry	SOx	1.04	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
AFOLU	3F - Field burning of agricultural residues	SOx	0.08	0.05	30.0	50.0	58.3	0.001	-39.6	0.0
Waste	5C1a - Municipal waste incineration	SOx	0.16	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
Waste	5C1bi - Industrial waste incineration	SOx	0.28	0.05	10.0	20.0	22.4	0.000	-81.4	0.0
Waste	5C1bii - Hazardous waste incineration	SOx	0.00	0.00	10.0	20.0	22.4	0.000	-100.0	0.0
Waste	5C1biii - Clinical waste incineration	SOx	0.00	0.00	10.0	20.0	22.4	0.000	-53.7	0.0
Waste	5C1biv - Sewage sludge incineration	SOx	0.04	0.04	10.0	20.0	22.4	0.000	-6.1	0.0
Waste	5C1bv - Cremation	SOx	0.00	0.00	10.0	20.0	22.4	0.000	269.5	0.0
Waste	5C2 - Open burning of waste	SOx	0.07	0.07	30.0	50.0	58.3	0.002	-1.5	0.0
TOTAL			1,783.60	88.15			5.4			0.3

APPENDIX 3 PROJECTED ACTIVITY DATA

Table A9.1 – WM Scenario for the agricultural sector.

Activity	Sector	Unit	2025	2030
DL	AGR_COWS	M animals	0.99339444	0.9902746
DS	AGR_COWS	M animals	0.77747334	0.77503162
OL	AGR_BEEF	M animals	1.80844404	1.80276447
OS	AGR_BEEF	M animals	1.89918562	1.89322106
PL	AGR_PIG	M animals	8.76716875	8.97492333
PS	AGR_PIG	M animals	0	0
LH	AGR_POULT	M animals	39.5044537	40.1684939
OP	AGR_POULT	M animals	139.387329	139.557617
SH	AGR_OTANI	M animals	7.87069575	7.69874828
HO	AGR_OTANI	M animals	0.43764918	0.43764918
FU	AGR_OTANI	M animals	0.175	0.175
BS	AGR_OTANI	M animals	0.42008917	0.45859968
CM	AGR_OTANI	M animals	12.3088445	11.5933201
NOF	PR_FERT	Mt	0.931876	0.931876
NOF	FERTPRO	kt N	273	273
NOF	FCON_UREA	kt N	288.366534	286.187442
NOF	FCON_OTHN	kt N	197.2433	195.752797
NOF	IO_NH3_EMISS	kt NH3	4.373	4.373
NOF	WT_NH3_EMISS	kt NH3	7.152	6.778
NOF	OTH_NH3_EMISS	kt NH3	15.4029785	15.4029785
FIRE_MASS	GRASSLAND	Mt biomass	0	0
FIRE_MASS	FOREST	Mt biomass	0	0
NOF	WASTE_AGR	Mt	1.26016363	1.34469675
NOF	AGR_ARABLE	M ha	6.78360116	7.02995197
RICE_AREA	AGR_ARABLE	M ha	0.25448362	0.27382456
AREA	RICE_FLOOD	M ha	0.25448362	0.27382456
AREA	RICE_INTER	M ha	0	0
AREA	RICE_UPLAND	M ha	0	0
AREA	GRASSLAND	M ha	3.962246	3.855024
AREA	HISTOSOLS	M ha	0.02324734	0.02324734
N_INPUT	CROP_RESID	kt N	153.295175	154.432265

Table A9.2 – WM Scenario for the solvent sector.

Activity	Sector	Unit	2025	2030
VEH	AUTO_P	kveh	0.00	0.00
VEH	AUTO_P_NEW	kveh	1149.89	1199.79
SC	COIL	Mm2	14.50	14.50
PNT	DECO_P	kt	580.50	578.52
SLV	DEGR	kt	15.43	15.43
SLV	DEGR_NEW	kt	0.00	0.00
POP	DOM_OS	mIn	59.58	58.94
TEX	DRY	kt	0.00	0.00
TEX	DRY_NEW	kt	343.00	343.00
EMI	EXD_GAS	kt	17.06	15.00
EMI	EXD_GAS_NEW	kt	0.00	0.00
EMI	EXD_LQ	kt	14.46	14.46
EMI	EXD_LQ_NEW	kt	0.00	0.00
SD	FATOIL	kt	5514.46	5514.46
POP	FOOD	mIn	59.58	58.94
ADH	GLUE_INH	kt	43.96	43.96
ADH	GLUE_INT	kt	175.84	175.84
EMI	IND_OTH	kt	18.11	18.15
PNT	IND_P_CNT	kt	13.44	13.47
PNT	IND_P_OT	kt	74.31	74.48
PNT	IND_P_PL	kt	11.83	11.86
EMI	INORG	kt	0.20	0.20
CTG	LEATHER	kt	20.68	20.68
EMI	ORG_STORE	kt	0.00	0.00
EMI	OTH_ORG_PR	kt	2.26	2.26
SLV	PHARMA	kt	131.56	132.83
PG	PIS	kt	1243.67	1239.89
EPS	PLSTYR_PR	kt	196.30	200.13
INK	PRT_OFFS	kt	0.00	0.00
INK	PRT_OFFS_NEW	kt	16.40	16.40
INK	PRT_PACK	kt	0.00	0.00
INK	PRT_PACK_NEW	kt	20.40	20.40
INK	PRT_PUB	kt	0.00	0.00
INK	PRT_PUB_NEW	kt	14.25	14.25
INK	PRT_SCR	kt	0.00	0.00
INK	PRT_SCR_NEW	kt	28.54	28.54
PVC	PVC_PR	kt	0.00	0.00
SHO	SHOE	Mp	177.89	176.10
EP	STCRACK_PR	kt	1593.14	1624.19
RUB	SYNTH_RUB	kt	478.56	487.89
TYR	TYRES	kt	200.72	204.63
PNT	VEHR_P	kt	26.93	33.86
PNT	VEHR_P_NEW	kt	0.00	0.00
POP	VEHTR	mIn	59.58	58.94
EMI	WASTE_VOC	kt	10.17	10.17
ENW	WIRE	kt	112.38	112.38
TIM	WOOD	Mm3	0.00	0.00
TIM	WOOD_CR	Mm3	0.00	0.00

SC	WOOD_P	Mm2	470.48	471.56
EMI	IND_OS	kt	0.26	0.26
EMI	OTHER_VOC	kt	0.00	0.00

Table A9.3 – WM Scenario for the industrial sector.

Activity	Sector	Unit	2025	2030
NOF	CONSTRUCT	M m2	32.24	32.83
NOF	MINE_BC	Mt	0.00	0.00
NOF	MINE_HC	Mt	0.00	0.00
NOF	MINE_OTH	Mt	0.03	0.03
NOF	OTHER_CO2	Mt CO2	4.76	4.76
NOF	OTHER_NOX	kt NOx	0.00	0.00
NOF	OTHER_PM	kt TSP	1.49	1.49
NOF	OTHER_SO2	kt SO2	2.00	2.00
NOF	OTHER_CH4	kt CH4	6.63	6.63
NOF	OTHER_N2O	kt N2O	2.98	2.98
NOF	PR_CEM	Mt	17.91	17.92
NOF	PR_LIME	Mt	2.61	2.64
NOF	PR_ALPRIM	Mt	0.00	0.00
NOF	PR_ALSEC	Mt	0.77	0.76
NOF	PR_BAOX	Mt	4.75	6.00
NOF	PR_BRIQ	Mt	0.00	0.00
NOF	PR_BRICK	Mt	5.08	5.15
NOF	PR_CAST	Mt	0.62	0.78
NOF	PR_CAST_F	Mt	0.62	0.78
NOF	PR_CBLACK	Mt	0.18	0.18
NOF	PR_COKE	Mt	2.07	2.06
NOF	PR_EARC	Mt	16.83	14.44
NOF	PR_GLASS	Mt	5.80	5.96
NOF	PR_HEARTH	Mt	0.00	0.00
NOF	PR_NIAC	Mt	0.15	0.15
NOF	PR_OTHER	Mt	0.54	0.54
NOF	PR_OT_NFME	Mt	0.00	0.00
NOF	PR_PELL	Mt	0.00	0.00
NOF	PR_PIGI	Mt	4.75	6.00
NOF	PR_PIGI_F	Mt	4.75	6.00
NOF	PR_PULP	Mt	0.00	0.00
NOF	PR_REF	Mt	82.36	69.22
NOF	PR_SINT	Mt	6.32	6.32
NOF	PR_SINT_F	Mt	6.32	6.32
NOF	PR_SUAC	Mt	0.92	0.97
CRU	PROD	PJ oil produced	228.00	225.00
GAS	PROD	PJ gas produced	254.17	250.00
CB_GAS	PROD	PJ gas produced	0.00	0.00
S_GAS	PROD	PJ gas produced	0.00	0.00
T_GAS	PROD	PJ gas produced	0.00	0.00
NOF	STH_AGR	Mt	31.43	31.43
NOF	STH_COAL	Mt	4.23	4.33
NOF	STH_FEORE	Mt	47.50	49.40
NOF	STH_NPK	Mt	0.29	0.28
NOF	STH_OTH_IN	Mt	83.01	85.08
GAS	TRANS	PJ gas transported	2432.95	2417.84

NOF	WASTE_FLR	PJ gas flared	5.01	4.21
NOF	MSW_URB	Mt waste	29.40	29.31
NOF	MSW_RUR	Mt waste	0.00	0.00
NOF	INW_TOT	Mt waste	28.96	29.99
NOF	SWD_HIST	Mt waste	0.00	0.00
NOF	MSW_COMP	Mt waste	0.00	0.00
NOF	INW_COMP	Mt waste	0.00	0.00
NOF	IND_PAP_COD	kt COD	17.11	17.84
NOF	IND_FOOD_COD	kt COD	707.34	761.07
NOF	IND_OTH_COD	kt COD	678.12	709.43
NOF	PR_ADIP	Mt	0.00	0.00
POP	ANY	M people	59.58	58.94

Table A9.4 – Emission factors for NOx emissions.

Sector	Fuel activity	Measure	Unit	FACTOR_ABTD [kt/unit]
APPLIC_L	MANURE_N	INHIB	[kt N]	0.012
APPLIC_L	MANURE_N	NOC	[kt N]	0.022
APPLIC_L	MANURE_N	PRECFARM	[kt N]	0.007
APPLIC_L	MANURE_N	VRT	[kt N]	0.018
APPLIC_L	RICE_N	NOC	[kt N]	0.011
APPLIC_L	SOIL_N	INHIB	[kt N]	0.011
APPLIC_L	SOIL_N	NOC	[kt N]	0.02
APPLIC_L	SOIL_N	PRECFARM	[kt N]	0.007
APPLIC_L	SOIL_N	VRT	[kt N]	0.016
APPLIC_M	MANURE_N	INHIB	[kt N]	0.012
APPLIC_M	MANURE_N	NOC	[kt N]	0.022
APPLIC_M	MANURE_N	PRECFARM	[kt N]	0.007
APPLIC_M	MANURE_N	VRT	[kt N]	0.018
APPLIC_M	RICE_N	NOC	[kt N]	0.011
APPLIC_M	SOIL_N	INHIB	[kt N]	0.011
APPLIC_M	SOIL_N	NOC	[kt N]	0.02
APPLIC_M	SOIL_N	PRECFARM	[kt N]	0.007
APPLIC_M	SOIL_N	VRT	[kt N]	0.016
APPLIC_S	MANURE_N	INHIB	[kt N]	0.012
APPLIC_S	MANURE_N	NOC	[kt N]	0.022
APPLIC_S	MANURE_N	PRECFARM	[kt N]	0.007
APPLIC_S	MANURE_N	VRT	[kt N]	0.018
APPLIC_S	RICE_N	NOC	[kt N]	0.011
APPLIC_S	SOIL_N	INHIB	[kt N]	0.011
APPLIC_S	SOIL_N	NOC	[kt N]	0.02
APPLIC_S	SOIL_N	PRECFARM	[kt N]	0.007
APPLIC_S	SOIL_N	VRT	[kt N]	0.016
CON_COMB	GAS	IOGCM	[PJ]	0.06
CON_COMB	GAS	IOGCSC	[PJ]	0.024
CON_COMB	GAS	IOGCSN	[PJ]	0.036

CON_COMB	GAS	NOC	[PJ]	0.12
CON_COMB	HC1	ISFCM	[PJ]	0.13
CON_COMB	HC1	ISFCSC	[PJ]	0.052
CON_COMB	HC1	ISFCSN	[PJ]	0.078
CON_COMB	HC1	NOC	[PJ]	0.26
CON_COMB	HF	IOGCM	[PJ]	0.11
CON_COMB	HF	IOGCSC	[PJ]	0.044
CON_COMB	HF	IOGCSN	[PJ]	0.066
CON_COMB	HF	NOC	[PJ]	0.22
CON_COMB	LPG	IOGCM	[PJ]	0.035
CON_COMB	LPG	NOC	[PJ]	0.07
CON_LOSS	HC1	NOC	[PJ]	0
DOM	GAS	DGCCOM	[PJ]	0.043
DOM	GAS	DGCCR	[PJ]	0.028
DOM	GAS	NOC	[PJ]	0.055
DOM	LPG	DMDCCO	[PJ]	0.053
DOM	LPG	DMDCCR	[PJ]	0.042
DOM	LPG	NOC	[PJ]	0.06
DOM	MD	DMDCCO	[PJ]	0.044
DOM	MD	DMDCCR	[PJ]	0.035
DOM	MD	NOC	[PJ]	0.05
DOM	OS1	NOC	[PJ]	0.107
GRAZE_L	EX_CTTL	INHIB	[kt N]	0.029
GRAZE_L	EX_CTTL	NOC	[kt N]	0.038
GRAZE_L	EX_SH	NOC	[kt N]	0.022
GRAZE_M	EX_CTTL	INHIB	[kt N]	0.029
GRAZE_M	EX_CTTL	NOC	[kt N]	0.038
GRAZE_M	EX_SH	NOC	[kt N]	0.022
GRAZE_S	EX_CTTL	INHIB	[kt N]	0.029
GRAZE_S	EX_CTTL	NOC	[kt N]	0.038
GRAZE_S	EX_SH	NOC	[kt N]	0.022
IN_BO_CHEM	GAS	IOGCM	[PJ]	0.035
IN_BO_CHEM	GAS	IOGCSC	[PJ]	0.014
IN_BO_CHEM	GAS	IOGCSN	[PJ]	0.021
IN_BO_CHEM	GAS	NOC	[PJ]	0.07
IN_BO_CHEM	HF	IOGCM	[PJ]	0.1
IN_BO_CHEM	HF	IOGCSC	[PJ]	0.04
IN_BO_CHEM	HF	ISFCSN	[PJ]	0.06
IN_BO_CHEM	HF	NOC	[PJ]	0.2
IN_BO_CHEM	LPG	IOGCM	[PJ]	0.035
IN_BO_CHEM	LPG	NOC	[PJ]	0.07
IN_BO_CON	GAS	IOGCM	[PJ]	0.035
IN_BO_CON	GAS	IOGCSC	[PJ]	0.014
IN_BO_CON	GAS	IOGCSN	[PJ]	0.021
IN_BO_CON	GAS	NOC	[PJ]	0.07

IN_BO_CON	HF	IOGCM	[PJ]	0.1
IN_BO_CON	HF	IOGCSC	[PJ]	0.04
IN_BO_CON	HF	ISFCSN	[PJ]	0.06
IN_BO_CON	HF	NOC	[PJ]	0.2
IN_BO_CON	LPG	IOGCM	[PJ]	0.035
IN_BO_CON	LPG	NOC	[PJ]	0.07
IN_BO_OTH	GAS	IOGCM	[PJ]	0.035
IN_BO_OTH	GAS	IOGCSC	[PJ]	0.014
IN_BO_OTH	GAS	IOGCSN	[PJ]	0.021
IN_BO_OTH	GAS	NOC	[PJ]	0.07
IN_BO_OTH	HF	IOGCM	[PJ]	0.1
IN_BO_OTH	HF	IOGCSC	[PJ]	0.04
IN_BO_OTH	HF	ISFCSN	[PJ]	0.06
IN_BO_OTH	HF	NOC	[PJ]	0.2
IN_BO_OTH	OS1	ISFCM	[PJ]	0.065
IN_BO_OTH	OS1	NOC	[PJ]	0.13
IN_BO_OTH_L	HC1	ISFCM	[PJ]	0.13
IN_BO_OTH_L	HC1	ISFCSC	[PJ]	0.052
IN_BO_OTH_L	HC1	ISFCSN	[PJ]	0.078
IN_BO_OTH_L	HC1	NOC	[PJ]	0.26
IN_BO_PAP	GAS	IOGCM	[PJ]	0.035
IN_BO_PAP	GAS	IOGCSC	[PJ]	0.014
IN_BO_PAP	GAS	IOGCSN	[PJ]	0.021
IN_BO_PAP	GAS	NOC	[PJ]	0.07
IN_BO_PAP	HF	IOGCM	[PJ]	0.1
IN_BO_PAP	HF	IOGCSC	[PJ]	0.04
IN_BO_PAP	HF	ISFCSN	[PJ]	0.06
IN_BO_PAP	HF	NOC	[PJ]	0.2
IN_OC	DC	NOC	[PJ]	0.02
IN_OC	GAS	IOGCM	[PJ]	0.06
IN_OC	GAS	IOGCSC	[PJ]	0.024
IN_OC	GAS	IOGCSN	[PJ]	0.036
IN_OC	GAS	NOC	[PJ]	0.12
IN_OC	HC1	ISFCM	[PJ]	0.052
IN_OC	HC1	ISFCSC	[PJ]	0.021
IN_OC	HC1	ISFCSN	[PJ]	0.031
IN_OC	HC1	NOC	[PJ]	0.104
IN_OC	HF	IOGCM	[PJ]	0.099
IN_OC	HF	IOGCSC	[PJ]	0.04
IN_OC	HF	ISFCSN	[PJ]	0.059
IN_OC	HF	NOC	[PJ]	0.198
IN_OC	MD	IOGCM	[PJ]	0.04
IN_OC	MD	NOC	[PJ]	0.08
MSW_URB_FOOD	NOF	NOC	[Mt waste]	0
MSW_URB_FOOD	NOF	SWD_FLA	[Mt waste]	0.03

MSW_URB_FOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_FOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_FOOD	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_FOOD	NOF	UNC_BURN	[Mt waste]	5.73
MSW_URB_GLA	NOF	NOC	[Mt waste]	0
MSW_URB_MET	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	SWD_FL A	[Mt waste]	0.03
MSW_URB_OTH	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_OTH	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_OTH	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_OTH	NOF	UNC_BURN	[Mt waste]	5.73
MSW_URB_PAP	NOF	NOC	[Mt waste]	0
MSW_URB_PAP	NOF	SWD_FL A	[Mt waste]	0.03
MSW_URB_PAP	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_PAP	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_PAP	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_PAP	NOF	UNC_BURN	[Mt waste]	5.73
MSW_URB_PLA	NOF	NOC	[Mt waste]	0
MSW_URB_PLA	NOF	SWD_FL A	[Mt waste]	0.03
MSW_URB_PLA	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_PLA	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_PLA	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_PLA	NOF	UNC_BURN	[Mt waste]	5.73
MSW_URB_TEX	NOF	NOC	[Mt waste]	0
MSW_URB_TEX	NOF	SWD_FL A	[Mt waste]	0.03
MSW_URB_TEX	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_TEX	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_TEX	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_TEX	NOF	UNC_BURN	[Mt waste]	5.73
MSW_URB_WOOD	NOF	NOC	[Mt waste]	0
MSW_URB_WOOD	NOF	SWD_FL A	[Mt waste]	0.03
MSW_URB_WOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.146
MSW_URB_WOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.573
MSW_URB_WOOD	NOF	TREAT_INC	[Mt waste]	0.03
MSW_URB_WOOD	NOF	UNC_BURN	[Mt waste]	5.73
PP_ENG	GAS	LFEUI	[PJ]	0.29
PP_ENG	GAS	LFEUII	[PJ]	0.13
PP_ENG	GAS	LFEUIII	[PJ]	0.08
PP_ENG	GAS	LFEUIV	[PJ]	0.04
PP_ENG	GAS	LFEUV	[PJ]	0.03
PP_ENG	GAS	LFEUVI	[PJ]	0.02
PP_ENG	GAS	NOC	[PJ]	1
PP_ENG	MD	NOC	[PJ]	1.3
PP_ENG	MD	TIWEUI	[PJ]	0.858

PP_ENG	MD	TIWEUII	[PJ]	0.78
PP_ENG	MD	TIWEUIII	[PJ]	0.585
PP_ENG	MD	TIWEUIV	[PJ]	0.39
PP_ENG	MD	TIWEUV	[PJ]	0.247
PP_ENG	MD	TIWEUVI	[PJ]	0.059
PP_EX_OTH	GAS	NOC	[PJ]	0.2
PP_EX_OTH	GAS	POGCM	[PJ]	0.07
PP_EX_OTH	GAS	POGCSC	[PJ]	0.04
PP_EX_OTH	HF	NOC	[PJ]	0.22
PP_EX_OTH	HF	POGCM	[PJ]	0.077
PP_EX_OTH	HF	POGCSC	[PJ]	0.044
PP_EX_OTH	MD	NOC	[PJ]	0.07
PP_EX_OTH	MD	POGCM	[PJ]	0.025
PP_EX_OTH	OS2	NOC	[PJ]	0.13
PP_EX_OTH	OS2	PHCCM	[PJ]	0.065
PP_EX_OTH	OS2	PSFCSN	[PJ]	0.039
PP_MOD	GAS	NOC	[PJ]	0.03
PP_MOD	GAS	POGSCR	[PJ]	0.006
PP_NEW	GAS	NOC	[PJ]	0.06
PP_NEW	GAS	POGSCR	[PJ]	0.012
PP_NEW	GSL	NOC	[PJ]	0.07
PP_NEW	HF	NOC	[PJ]	0.1
PP_NEW	HF	POGSCR	[PJ]	0.02
PP_NEW	OS1	NOC	[PJ]	0.065
PP_NEW	OS1	PSFSNC	[PJ]	0.039
PP_NEW	OS2	NOC	[PJ]	0.065
PP_NEW	OS2	PHCSCR	[PJ]	0.013
PP_NEW	OS2	PSFSNC	[PJ]	0.039
PROD_AGAS	CRU	NOC	[PJ]	0.004
PROD_AGAS	CRU	REC_USE	[PJ]	0
PROD_AGAS	CRU	REC_USE2	[PJ]	0
PR_ALSEC	NOF	NOC	[Mt]	0.007
PR_ALSEC	NOF	PRNOX1	[Mt]	0.004
PR_ALSEC	NOF	PRNOX2	[Mt]	0.003
PR_ALSEC	NOF	PRNOX3	[Mt]	0.001
PR_BRICK	NOF	DDK	[Mt]	0
PR_BRICK	NOF	FCBTK	[Mt]	0
PR_BRICK	NOF	HK	[Mt]	0.067
PR_BRICK	NOF	IDK_BTK	[Mt]	0.492
PR_BRICK	NOF	MCBTK	[Mt]	0
PR_BRICK	NOF	MK	[Mt]	0
PR_BRICK	NOF	NOC	[Mt]	0.984
PR_BRICK	NOF	TK_COAL	[Mt]	0.018
PR_BRICK	NOF	TK_EOF	[Mt]	0.197
PR_BRICK	NOF	TK_OTHER	[Mt]	0.237

PR_BRICK	NOF	VS BK	[Mt]	0.394
PR_BRICK	NOF	ZIG_ZAG	[Mt]	0
PR_CEM	NOF	NOC	[Mt]	2.35
PR_CEM	NOF	PRNOX1	[Mt]	1.41
PR_CEM	NOF	PRNOX2	[Mt]	0.94
PR_CEM	NOF	PRNOX3	[Mt]	0.235
PR_COKE	NOF	NOC	[Mt]	1.1
PR_COKE	NOF	PRNOX1	[Mt]	0.66
PR_COKE	NOF	PRNOX2	[Mt]	0.44
PR_COKE	NOF	PRNOX3	[Mt]	0.11
PR_EARC	NOF	NOC	[Mt]	0.008
PR_EARC	NOF	PRNOX1	[Mt]	0.005
PR_EARC	NOF	PRNOX2	[Mt]	0.003
PR_EARC	NOF	PRNOX3	[Mt]	0.001
PR_GLASS	NOF	NOC	[Mt]	8.12
PR_GLASS	NOF	PRNOX1	[Mt]	4.872
PR_GLASS	NOF	PRNOX2	[Mt]	3.248
PR_GLASS	NOF	PRNOX3	[Mt]	0.812
PR_LIME	NOF	NOC	[Mt]	2.35
PR_LIME	NOF	PRNOX1	[Mt]	1.41
PR_LIME	NOF	PRNOX2	[Mt]	0.94
PR_LIME	NOF	PRNOX3	[Mt]	0.235
PR_NIAC	NOF	NOC	[Mt]	3
PR_NIAC	NOF	PRNOX1	[Mt]	1.8
PR_NIAC	NOF	PRNOX2	[Mt]	1.2
PR_NIAC	NOF	PRNOX3	[Mt]	0.3
PR_PIGI	NOF	NOC	[Mt]	0
PR_PIGI	NOF	PRNOX1	[Mt]	0
PR_PIGI	NOF	PRNOX2	[Mt]	0
PR_PIGI	NOF	PRNOX3	[Mt]	0
PR_REF	NOF	NOC	[Mt]	0.2
PR_REF	NOF	PRNOX1	[Mt]	0.12
PR_REF	NOF	PRNOX2	[Mt]	0.08
PR_REF	NOF	PRNOX3	[Mt]	0.02
PR_SINT	NOF	NOC	[Mt]	1.33
PR_SINT	NOF	PRNOX1	[Mt]	0.798
PR_SINT	NOF	PRNOX2	[Mt]	0.532
PR_SINT	NOF	PRNOX3	[Mt]	0.133
PR_SUAC	NOF	NOC	[Mt]	0
PR_SUAC	NOF	PRNOX1	[Mt]	0
PR_SUAC	NOF	PRNOX2	[Mt]	0
PR_SUAC	NOF	PRNOX3	[Mt]	0
TRA_OTL_L	GAS	NOC	[PJ]	0.17
TRA_OTL_L	GAS	STLMCM	[PJ]	0.085
TRA_OTL_L	GAS	STLSCR	[PJ]	0.017

TRA_OT_S_L	HF	NOC	[PJ]	1.6
TRA_OT_S_L	HF	STLHCM	[PJ]	0.8
TRA_OT_S_L	HF	STLSCR	[PJ]	0.16
TRA_OT_S_L	MD	NOC	[PJ]	1.346
TRA_OT_S_L	MD	STLMCM	[PJ]	0.673
TRA_OT_S_L	MD	STLSCR	[PJ]	0.135
TRA_OT_S_M	GAS	NOC	[PJ]	0.17
TRA_OT_S_M	GAS	STMCM	[PJ]	0.085
TRA_OT_S_M	GAS	STMSCR	[PJ]	0.017
TRA_OT_S_M	MD	NOC	[PJ]	1.346
TRA_OT_S_M	MD	STMCM	[PJ]	0.673
TRA_OT_S_M	MD	STMSCR	[PJ]	0.135
TRA_OT_AGR	GSL	LFEUI	[PJ]	0.249
TRA_OT_AGR	GSL	LFEUII	[PJ]	0.112
TRA_OT_AGR	GSL	LFEUIII	[PJ]	0.069
TRA_OT_AGR	GSL	LFEUIV	[PJ]	0.034
TRA_OT_AGR	GSL	LFEUV	[PJ]	0.026
TRA_OT_AGR	GSL	LFEUVI	[PJ]	0.1
TRA_OT_AGR	GSL	NOC	[PJ]	0.3
TRA_OT_AGR	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_AGR	MD	CAGEUI	[PJ]	0.693
TRA_OT_AGR	MD	CAGEUII	[PJ]	0.525
TRA_OT_AGR	MD	CAGEUIII	[PJ]	0.315
TRA_OT_AGR	MD	CAGEUIII_PF	[PJ]	0.315
TRA_OT_AGR	MD	CAGEUIV	[PJ]	0.273
TRA_OT_AGR	MD	CAGEUV	[PJ]	0.063
TRA_OT_AGR	MD	CAGEUVI	[PJ]	0.032
TRA_OT_AGR	MD	NOC	[PJ]	1.05
TRA_OT_AGR	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_AIR	GSL	NOC	[PJ]	0.022
TRA_OT_CNS	MD	CAGEUI	[PJ]	0.858
TRA_OT_CNS	MD	CAGEUII	[PJ]	0.65
TRA_OT_CNS	MD	CAGEUIII	[PJ]	0.39
TRA_OT_CNS	MD	CAGEUIII_PF	[PJ]	0.39
TRA_OT_CNS	MD	CAGEUIV	[PJ]	0.338
TRA_OT_CNS	MD	CAGEUV	[PJ]	0.078
TRA_OT_CNS	MD	CAGEUVI	[PJ]	0.039
TRA_OT_CNS	MD	NOC	[PJ]	1.3
TRA_OT_CNS	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_CNS	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_INW	GSL	LFEUI	[PJ]	0.066
TRA_OT_INW	GSL	LFEUII	[PJ]	0.03
TRA_OT_INW	GSL	LFEUIII	[PJ]	0.018

TRA_OT_INW	GSL	LFEUIV	[PJ]	0.009
TRA_OT_INW	GSL	LFEUV	[PJ]	0.007
TRA_OT_INW	GSL	LFEUVI	[PJ]	0.005
TRA_OT_INW	GSL	NOC	[PJ]	0.227
TRA_OT_INW	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	GAS	LFEUI	[PJ]	0.143
TRA_OT_LB	GAS	LFEUII	[PJ]	0.061
TRA_OT_LB	GAS	LFEUIII	[PJ]	0.035
TRA_OT_LB	GAS	LFEUIV	[PJ]	0.019
TRA_OT_LB	GAS	LFEUV	[PJ]	0.017
TRA_OT_LB	GAS	LFEUVI	[PJ]	0.018
TRA_OT_LB	GAS	NOC	[PJ]	0.86
TRA_OT_LB	GSL	LFEUI	[PJ]	0.084
TRA_OT_LB	GSL	LFEUII	[PJ]	0.038
TRA_OT_LB	GSL	LFEUIII	[PJ]	0.023
TRA_OT_LB	GSL	LFEUIV	[PJ]	0.012
TRA_OT_LB	GSL	LFEUV	[PJ]	0.009
TRA_OT_LB	GSL	LFEUVI	[PJ]	0.006
TRA_OT_LB	GSL	NOC	[PJ]	0.29
TRA_OT_LB	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LB	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	MD	HDEUI	[PJ]	0.764
TRA_OT_LB	MD	HDEUII	[PJ]	0.811
TRA_OT_LB	MD	HDEUIII	[PJ]	0.702
TRA_OT_LB	MD	HDEUIV	[PJ]	0.404
TRA_OT_LB	MD	HDEUV	[PJ]	0.246
TRA_OT_LB	MD	HDEUVI	[PJ]	0.05
TRA_OT_LB	MD	NOC	[PJ]	1.1
TRA_OT_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_RAI	H2	NOC	[PJ]	0.007
TRA_OT_RAI	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_RAI	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDB	GAS	HDSEI	[PJ]	0.129
TRA_RD_HDB	GAS	HDSEII	[PJ]	0.069
TRA_RD_HDB	GAS	HDSEIII	[PJ]	0.029
TRA_RD_HDB	GAS	NOC	[PJ]	0.562
TRA_RD_HDB	MD	HDEUI	[PJ]	0.773
TRA_RD_HDB	MD	HDEUII	[PJ]	0.866
TRA_RD_HDB	MD	HDEUIII	[PJ]	0.683
TRA_RD_HDB	MD	HDEUIV	[PJ]	0.483
TRA_RD_HDB	MD	HDEUV	[PJ]	0.439

TRA_RD_HDB	MD	HDEUVI	[PJ]	0.05
TRA_RD_HDB	MD	HDEUVII	[PJ]	0.036
TRA_RD_HDB	MD	NOC	[PJ]	0.939
TRA_RD_HDB	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDB	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDT	GAS	HDSEI	[PJ]	0.129
TRA_RD_HDT	GAS	HDSEII	[PJ]	0.069
TRA_RD_HDT	GAS	HDSEIII	[PJ]	0.029
TRA_RD_HDT	GAS	NOC	[PJ]	0.562
TRA_RD_HDT	MD	HDEUI	[PJ]	0.759
TRA_RD_HDT	MD	HDEUII	[PJ]	0.829
TRA_RD_HDT	MD	HDEUIII	[PJ]	0.637
TRA_RD_HDT	MD	HDEUIV	[PJ]	0.432
TRA_RD_HDT	MD	HDEUV	[PJ]	0.298
TRA_RD_HDT	MD	HDEUVI	[PJ]	0.043
TRA_RD_HDT	MD	HDEUVII	[PJ]	0.019
TRA_RD_HDT	MD	NOC	[PJ]	0.987
TRA_RD_HDT	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDT	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD2	GSL	MMO2I	[PJ]	0.14
TRA_RD_LD2	GSL	MMO2II	[PJ]	0.273
TRA_RD_LD2	GSL	MMO2III	[PJ]	0.088
TRA_RD_LD2	GSL	MMO2IV	[PJ]	0.261
TRA_RD_LD2	GSL	MMO2V	[PJ]	0.208
TRA_RD_LD2	GSL	MMO2VI	[PJ]	0.208
TRA_RD_LD2	GSL	NOC	[PJ]	0.034
TRA_RD_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	GAS	LFEUI	[PJ]	0.129
TRA_RD_LD4C	GAS	LFEUII	[PJ]	0.069
TRA_RD_LD4C	GAS	LFEUIII	[PJ]	0.029
TRA_RD_LD4C	GAS	LFEUIV	[PJ]	0.02
TRA_RD_LD4C	GAS	LFEUV	[PJ]	0.018
TRA_RD_LD4C	GAS	LFEUVI	[PJ]	0.018
TRA_RD_LD4C	GAS	NOC	[PJ]	0.562
TRA_RD_LD4C	GSL	LFEUI	[PJ]	0.133
TRA_RD_LD4C	GSL	LFEUII	[PJ]	0.063
TRA_RD_LD4C	GSL	LFEUIII	[PJ]	0.025
TRA_RD_LD4C	GSL	LFEUIV	[PJ]	0.016
TRA_RD_LD4C	GSL	LFEUV	[PJ]	0.01
TRA_RD_LD4C	GSL	LFEUVI	[PJ]	0.012
TRA_RD_LD4C	GSL	NOC	[PJ]	0.72
TRA_RD_LD4C	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	LPG	LFEUI	[PJ]	0.161

TRA_RD_LD4C	LPG	LFEUII	[PJ]	0.072
TRA_RD_LD4C	LPG	LFEUIII	[PJ]	0.037
TRA_RD_LD4C	LPG	LFEUIV	[PJ]	0.02
TRA_RD_LD4C	LPG	LFEUV	[PJ]	0.021
TRA_RD_LD4C	LPG	LFEUVI	[PJ]	0.032
TRA_RD_LD4C	LPG	NOC	[PJ]	0.921
TRA_RD_LD4C	MD	MDEUI	[PJ]	0.282
TRA_RD_LD4C	MD	MDEUII	[PJ]	0.264
TRA_RD_LD4C	MD	MDEUIII	[PJ]	0.311
TRA_RD_LD4C	MD	MDEUIV	[PJ]	0.228
TRA_RD_LD4C	MD	MDEUV	[PJ]	0.233
TRA_RD_LD4C	MD	MDEUVI	[PJ]	0.043
TRA_RD_LD4C	MD	MDEUVII	[PJ]	0.021
TRA_RD_LD4C	MD	MDEUVip	[PJ]	0.39
TRA_RD_LD4C	MD	MDEUVit	[PJ]	0.135
TRA_RD_LD4C	MD	NOC	[PJ]	0.277
TRA_RD_LD4C	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4T	GAS	LFEUI	[PJ]	0.092
TRA_RD_LD4T	GAS	LFEUII	[PJ]	0.038
TRA_RD_LD4T	GAS	LFEUIII	[PJ]	0.022
TRA_RD_LD4T	GAS	LFEUIV	[PJ]	0.014
TRA_RD_LD4T	GAS	LFEUV	[PJ]	0.012
TRA_RD_LD4T	GAS	LFEUVI	[PJ]	0.013
TRA_RD_LD4T	GAS	NOC	[PJ]	0.617
TRA_RD_LD4T	MD	MDEUI	[PJ]	0.363
TRA_RD_LD4T	MD	MDEUII	[PJ]	0.359
TRA_RD_LD4T	MD	MDEUIII	[PJ]	0.297
TRA_RD_LD4T	MD	MDEUIV	[PJ]	0.24
TRA_RD_LD4T	MD	MDEUV	[PJ]	0.419
TRA_RD_LD4T	MD	MDEUVI	[PJ]	0.348
TRA_RD_LD4T	MD	MDEUVII	[PJ]	0.013
TRA_RD_LD4T	MD	MDEUVip	[PJ]	0.124
TRA_RD_LD4T	MD	MDEUVit	[PJ]	0.082
TRA_RD_LD4T	MD	NOC	[PJ]	0.421
TRA_RD_LD4T	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4T	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_M4	GSL	MOT4I	[PJ]	0.232
TRA_RD_M4	GSL	MOT4II	[PJ]	0.164
TRA_RD_M4	GSL	MOT4III	[PJ]	0.04
TRA_RD_M4	GSL	MOT4IV	[PJ]	0.018
TRA_RD_M4	GSL	MOT4V	[PJ]	0.018
TRA_RD_M4	GSL	MOT4VI	[PJ]	0.018
TRA_RD_M4	GSL	NOC	[PJ]	0.166
TRA_RD_M4	GSL_NV_HE	NOC	[thousand of vehicles]	0

TRA_RD_M4	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
WASTE_AGR	NOF	BAN	[Mt]	1
WASTE_AGR	NOF	NOC	[Mt]	3
WASTE_FLR	NOF	FLR_GP	[PJ]	0.002
WASTE_FLR	NOF	NOC	[PJ]	0.03

Table A9.5 – Emission factors for SOx emissions.

Sector	Fuel activity	Measure	Unit	FACTOR_ABTD [kt/unit]
CON_COMB	GAS	NOC	[PJ]	0
CON_COMB	HC1	IWFGD	[PJ]	0.018
CON_COMB	HC1	LINJ	[PJ]	0.14
CON_COMB	HC1	LSCO	[PJ]	0.35
CON_COMB	HC1	NOC	[PJ]	0.35
CON_COMB	HF	IWFGD	[PJ]	0.077
CON_COMB	HF	LSHF	[PJ]	0.25
CON_COMB	HF	NOC	[PJ]	1.55
CON_COMB	HF	RFGD	[PJ]	0.031
CON_LOSS	HC1	NOC	[PJ]	0
DOM	GAS	NOC	[PJ]	0
DOM	MD	LSMD1	[PJ]	0.094
DOM	MD	LSMD2	[PJ]	0.021
DOM	MD	LSMD3	[PJ]	0
DOM	MD	NOC	[PJ]	0.376
DOM	OS1	NOC	[PJ]	0.013
IN_BO_CHEM	GAS	NOC	[PJ]	0
IN_BO_CHEM	HF	IWFGD	[PJ]	0.024
IN_BO_CHEM	HF	LSHF	[PJ]	0.25
IN_BO_CHEM	HF	NOC	[PJ]	0.48
IN_BO_CON	GAS	NOC	[PJ]	0
IN_BO_CON	HF	IWFGD	[PJ]	0.024
IN_BO_CON	HF	LSHF	[PJ]	0.25
IN_BO_CON	HF	NOC	[PJ]	0.48
IN_BO_OTH	GAS	NOC	[PJ]	0
IN_BO_OTH	HF	IWFGD	[PJ]	0.024
IN_BO_OTH	HF	LSHF	[PJ]	0.25
IN_BO_OTH	HF	NOC	[PJ]	0.48
IN_BO_OTH	OS1	NOC	[PJ]	0.037
IN_BO_OTH_L	HC1	IWFGD	[PJ]	0.018
IN_BO_OTH_L	HC1	LINJ	[PJ]	0.14
IN_BO_OTH_L	HC1	LSCO	[PJ]	0.35
IN_BO_OTH_L	HC1	NOC	[PJ]	0.35
IN_BO_PAP	GAS	NOC	[PJ]	0
IN_BO_PAP	HF	IWFGD	[PJ]	0.024

IN_BO_PAP	HF	LSHF	[PJ]	0.25
IN_BO_PAP	HF	NOC	[PJ]	0.48
IN_OC	DC	NOC	[PJ]	0.027
IN_OC	GAS	NOC	[PJ]	0
IN_OC	HC1	IWFGD	[PJ]	0.003
IN_OC	HC1	LINJ	[PJ]	0.022
IN_OC	HC1	LSCO	[PJ]	0.055
IN_OC	HC1	NOC	[PJ]	0.055
IN_OC	HF	IWFGD	[PJ]	0.022
IN_OC	HF	LSHF	[PJ]	0.225
IN_OC	HF	NOC	[PJ]	0.432
IN_OC	MD	LSMD1	[PJ]	0.094
IN_OC	MD	LSMD2	[PJ]	0.021
IN_OC	MD	NOC	[PJ]	0.376
MSW_URB_FOOD	NOF	NOC	[Mt waste]	0
MSW_URB_FOOD	NOF	SWD_FL A	[Mt waste]	0.002
MSW_URB_FOOD	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_FOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_FOOD	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_FOOD	NOF	UNC_BURN	[Mt waste]	0.225
MSW_URB_GLA	NOF	NOC	[Mt waste]	0
MSW_URB_MET	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	SWD_FL A	[Mt waste]	0.002
MSW_URB_OTH	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_OTH	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_OTH	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_OTH	NOF	UNC_BURN	[Mt waste]	0.225
MSW_URB_PAP	NOF	NOC	[Mt waste]	0
MSW_URB_PAP	NOF	SWD_FL A	[Mt waste]	0.002
MSW_URB_PAP	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_PAP	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_PAP	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_PAP	NOF	UNC_BURN	[Mt waste]	0.225
MSW_URB_PLA	NOF	NOC	[Mt waste]	0
MSW_URB_PLA	NOF	SWD_FL A	[Mt waste]	0.002
MSW_URB_PLA	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_PLA	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_PLA	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_PLA	NOF	UNC_BURN	[Mt waste]	0.225
MSW_URB_TEX	NOF	NOC	[Mt waste]	0
MSW_URB_TEX	NOF	SWD_FL A	[Mt waste]	0.002
MSW_URB_TEX	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_TEX	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_TEX	NOF	TREAT_INC	[Mt waste]	0.002

MSW_URB_TEX	NOF	UNC_BURN	[Mt waste]	0.225
MSW_URB_WOOD	NOF	NOC	[Mt waste]	0
MSW_URB_WOOD	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_WOOD	NOF	SWD_UNM_HIGH	[Mt waste]	0.045
MSW_URB_WOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.022
MSW_URB_WOOD	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_WOOD	NOF	UNC_BURN	[Mt waste]	0.225
NONEN	GSL	NOC	[PJ]	0
OTHER_SO2	NOF	NOC	[kt SO2]	1
PP_ENG	GAS	NOC	[PJ]	0.002
PP_ENG	MD	LSMD1	[PJ]	0.094
PP_ENG	MD	LSMD2	[PJ]	0.021
PP_ENG	MD	LSMD3	[PJ]	0
PP_ENG	MD	NOC	[PJ]	0.376
PP_EX_OTH	GAS	NOC	[PJ]	0.002
PP_EX_OTH	HF	LSHF	[PJ]	0.25
PP_EX_OTH	HF	NOC	[PJ]	2
PP_EX_OTH	HF	PRWFGD	[PJ]	0.2
PP_EX_OTH	HF	PWFGD	[PJ]	0.1
PP_EX_OTH	MD	LSMD1	[PJ]	0.094
PP_EX_OTH	MD	LSMD2	[PJ]	0.021
PP_EX_OTH	MD	NOC	[PJ]	0.376
PP_EX_OTH	OS2	LINJ	[PJ]	0.05
PP_EX_OTH	OS2	NOC	[PJ]	0.125
PP_EX_OTH	OS2	PRWFGD	[PJ]	0.013
PP_EX_OTH	OS2	PWFGD	[PJ]	0.006
PP_MOD	GAS	NOC	[PJ]	0.002
PP_NEW	GAS	NOC	[PJ]	0.002
PP_NEW	GSL	NOC	[PJ]	0.004
PP_NEW	HF	LSHF	[PJ]	0.25
PP_NEW	HF	NOC	[PJ]	2
PP_NEW	HF	PWFGD	[PJ]	0.1
PP_NEW	HF	RFGD	[PJ]	0.04
PP_NEW	OS1	NOC	[PJ]	0.037
PP_NEW	OS2	LINJ	[PJ]	0.05
PP_NEW	OS2	NOC	[PJ]	0.125
PP_NEW	OS2	PWFGD	[PJ]	0.006
PROD_AGAS	CRU	NOC	[PJ]	0.106
PROD_AGAS	CRU	REC_USE	[PJ]	0
PROD_AGAS	CRU	REC_USE2	[PJ]	0
PR_ALSEC	NOF	NOC	[Mt]	0.013
PR_ALSEC	NOF	SO2PR1	[Mt]	0.006
PR_ALSEC	NOF	SO2PR2	[Mt]	0.004
PR_ALSEC	NOF	SO2PR3	[Mt]	0.001
PR_BRICK	NOF	DDK	[Mt]	0.4

PR_BRICK	NOF	FCBTK	[Mt]	0
PR_BRICK	NOF	HK	[Mt]	0.72
PR_BRICK	NOF	IDK_BTK	[Mt]	1.729
PR_BRICK	NOF	MCBTK	[Mt]	0.4
PR_BRICK	NOF	MK	[Mt]	0.1
PR_BRICK	NOF	NOC	[Mt]	3.458
PR_BRICK	NOF	TK_COAL	[Mt]	0
PR_BRICK	NOF	TK_EOF	[Mt]	0.692
PR_BRICK	NOF	TK_OTHER	[Mt]	0.05
PR_BRICK	NOF	VSBK	[Mt]	1.037
PR_BRICK	NOF	ZIG_ZAG	[Mt]	0.32
PR_CEM	NOF	NOC	[Mt]	0.95
PR_CEM	NOF	SO2PR1	[Mt]	0.475
PR_CEM	NOF	SO2PR2	[Mt]	0.285
PR_CEM	NOF	SO2PR3	[Mt]	0.095
PR_COKE	NOF	NOC	[Mt]	1.3
PR_COKE	NOF	SO2PR1	[Mt]	0.65
PR_COKE	NOF	SO2PR2	[Mt]	0.195
PR_COKE	NOF	SO2PR3	[Mt]	0.039
PR_EARC	NOF	NOC	[Mt]	0.001
PR_EARC	NOF	SO2PR1	[Mt]	0
PR_EARC	NOF	SO2PR2	[Mt]	0
PR_EARC	NOF	SO2PR3	[Mt]	0
PR_GLASS	NOF	NOC	[Mt]	1.74
PR_GLASS	NOF	SO2PR1	[Mt]	0.87
PR_GLASS	NOF	SO2PR2	[Mt]	0.522
PR_GLASS	NOF	SO2PR3	[Mt]	0.174
PR_LIME	NOF	NOC	[Mt]	0.95
PR_LIME	NOF	SO2PR1	[Mt]	0.475
PR_LIME	NOF	SO2PR2	[Mt]	0.285
PR_LIME	NOF	SO2PR3	[Mt]	0.095
PR_NIAC	NOF	NOC	[Mt]	0
PR_NIAC	NOF	SO2PR1	[Mt]	0
PR_NIAC	NOF	SO2PR2	[Mt]	0
PR_NIAC	NOF	SO2PR3	[Mt]	0
PR_PIGI	NOF	NOC	[Mt]	0
PR_PIGI	NOF	SO2PR1	[Mt]	0
PR_PIGI	NOF	SO2PR2	[Mt]	0
PR_PIGI	NOF	SO2PR3	[Mt]	0
PR_REF	NOF	NOC	[Mt]	0.54
PR_REF	NOF	SO2PR1	[Mt]	0.27
PR_REF	NOF	SO2PR2	[Mt]	0.162
PR_REF	NOF	SO2PR3	[Mt]	0.054
PR_SINT	NOF	NOC	[Mt]	1.3
PR_SINT	NOF	SO2PR1	[Mt]	0.65

PR_SINT	NOF	SO2PR2	[Mt]	0.39
PR_SINT	NOF	SO2PR3	[Mt]	0.26
PR_SUAC	NOF	NOC	[Mt]	10
PR_SUAC	NOF	SO2PR1	[Mt]	5
PR_SUAC	NOF	SO2PR2	[Mt]	3
PR_SUAC	NOF	SO2PR3	[Mt]	1
TRA_OT_S_L	GAS	NOC	[PJ]	0
TRA_OT_S_L	HF	LSHF	[PJ]	0.241
TRA_OT_S_L	HF	NOC	[PJ]	1.542
TRA_OT_S_L	MD	LSMD1	[PJ]	0.094
TRA_OT_S_L	MD	LSMD2	[PJ]	0.021
TRA_OT_S_L	MD	NOC	[PJ]	0.141
TRA_OT_S_M	GAS	NOC	[PJ]	0
TRA_OT_S_M	MD	LSMD1	[PJ]	0.094
TRA_OT_S_M	MD	LSMD2	[PJ]	0.021
TRA_OT_S_M	MD	NOC	[PJ]	0.141
TRA_OT_AGR	GSL	LSGSL	[PJ]	0
TRA_OT_AGR	GSL	NOC	[PJ]	0.004
TRA_OT_AGR	MD	LSMD1	[PJ]	0.094
TRA_OT_AGR	MD	LSMD2	[PJ]	0.021
TRA_OT_AGR	MD	LSMD3	[PJ]	0
TRA_OT_AGR	MD	NOC	[PJ]	0.376
TRA_OT_AIR	GSL	NOC	[PJ]	0.003
TRA_OT_CNS	MD	LSMD1	[PJ]	0.094
TRA_OT_CNS	MD	LSMD2	[PJ]	0.021
TRA_OT_CNS	MD	LSMD3	[PJ]	0
TRA_OT_CNS	MD	NOC	[PJ]	0.376
TRA_OT_INW	GSL	LSGSL	[PJ]	0
TRA_OT_INW	GSL	NOC	[PJ]	0.004
TRA_OT_LB	GAS	NOC	[PJ]	0
TRA_OT_LB	GSL	LSGSL	[PJ]	0
TRA_OT_LB	GSL	NOC	[PJ]	0.004
TRA_OT_LB	MD	LSMD1	[PJ]	0.094
TRA_OT_LB	MD	LSMD2	[PJ]	0.021
TRA_OT_LB	MD	LSMD3	[PJ]	0
TRA_OT_LB	MD	NOC	[PJ]	0.376
TRA_RD_HDB	GAS	NOC	[PJ]	0
TRA_RD_HDB	MD	LSMD1	[PJ]	0.094
TRA_RD_HDB	MD	LSMD2	[PJ]	0.021
TRA_RD_HDB	MD	LSMD3	[PJ]	0
TRA_RD_HDB	MD	NOC	[PJ]	0.376
TRA_RD_HDT	GAS	NOC	[PJ]	0
TRA_RD_HDT	MD	LSMD1	[PJ]	0.094
TRA_RD_HDT	MD	LSMD2	[PJ]	0.021
TRA_RD_HDT	MD	LSMD3	[PJ]	0

TRA_RD_HDT	MD	NOC	[PJ]	0.376
TRA_RD_LD2	GSL	LSGSL	[PJ]	0
TRA_RD_LD2	GSL	NOC	[PJ]	0.004
TRA_RD_LD4C	GAS	NOC	[PJ]	0
TRA_RD_LD4C	GSL	LSGSL	[PJ]	0
TRA_RD_LD4C	GSL	NOC	[PJ]	0.004
TRA_RD_LD4C	MD	LSMD1	[PJ]	0.094
TRA_RD_LD4C	MD	LSMD2	[PJ]	0.021
TRA_RD_LD4C	MD	LSMD3	[PJ]	0
TRA_RD_LD4C	MD	NOC	[PJ]	0.376
TRA_RD_LD4T	GAS	NOC	[PJ]	0
TRA_RD_LD4T	MD	LSMD1	[PJ]	0.094
TRA_RD_LD4T	MD	LSMD2	[PJ]	0.021
TRA_RD_LD4T	MD	LSMD3	[PJ]	0
TRA_RD_LD4T	MD	NOC	[PJ]	0.376
TRA_RD_M4	GSL	LSGSL	[PJ]	0
TRA_RD_M4	GSL	NOC	[PJ]	0.004
WASTE_AGR	NOF	BAN	[Mt]	0
WASTE_AGR	NOF	NOC	[Mt]	0.6
WASTE_FLR	NOF	FLR_GP	[PJ]	0.599
WASTE_FLR	NOF	NOC	[PJ]	0.749

Table A9.6 – Emission factors for PM2.5 emissions.

Sector	Fuel activity	Measure	Unit	FACTOR_ABTD [kt/unit]
AGR_ARABLE	NOF	ALTER	[M ha]	0.178
AGR_ARABLE	NOF	NOC	[M ha]	0.188
AGR_BEEF	NOF	FEED_MOD	[M animals]	0.083
AGR_BEEF	NOF	HAY_SIL	[M animals]	0.083
AGR_BEEF	NOF	NOC	[M animals]	0.092
AGR_COWS	NOF	FEED_MOD	[M animals]	0.078
AGR_COWS	NOF	HAY_SIL	[M animals]	0.078
AGR_COWS	NOF	NOC	[M animals]	0.087
AGR_OTANI	NOF	AGR1	[M animals]	0
AGR_OTANI	NOF	NOC	[M animals]	0
AGR_PIG	NOF	FEED_MOD	[M animals]	0.07
AGR_PIG	NOF	NOC	[M animals]	0.078
AGR_POULT	NOF	FEED_MOD	[M animals]	0.009
AGR_POULT	NOF	FREE	[M animals]	0.01
AGR_POULT	NOF	NOC	[M animals]	0.011
CONSTRUCT	NOF	NOC	[M m2]	0.009
CONSTRUCT	NOF	SPRAY	[M m2]	0.008
CON_COMB	GAS	NOC	[PJ]	0
CON_COMB	HF	GHIND	[PJ]	0.008

CON_COMB	HF	IN_HED	[PJ]	0
CON_COMB	HF	NOC	[PJ]	0.012
CON_COMB	LPG	NOC	[PJ]	0
CON_COMB1	HC1	IN_CYC	[PJ]	0.076
CON_COMB1	HC1	IN_ESP1	[PJ]	0.008
CON_COMB1	HC1	IN_ESP2	[PJ]	0.004
CON_COMB1	HC1	IN_HED	[PJ]	0.001
CON_COMB1	HC1	NOC	[PJ]	0.109
CON_COMB2	HC1	IN_ESP1	[PJ]	0.012
CON_COMB2	HC1	IN_ESP2	[PJ]	0.007
CON_COMB2	HC1	IN_HED	[PJ]	0.002
CON_COMB2	HC1	NOC	[PJ]	0.176
CON_COMB3	HC1	IN_ESP1	[PJ]	0.013
CON_COMB3	HC1	IN_ESP2	[PJ]	0.007
CON_COMB3	HC1	IN_HED	[PJ]	0.002
CON_COMB3	HC1	NOC	[PJ]	0.187
DOM	GAS	NOC	[PJ]	0
DOM	LPG	NOC	[PJ]	0
DOM	MD	GHDOM	[PJ]	0.001
DOM	MD	NOC	[PJ]	0.001
DOM_FPLACE	FWD	FP_IMP	[PJ]	0.286
DOM_FPLACE	FWD	FP_NEW	[PJ]	0.153
DOM_FPLACE	FWD	NOC	[PJ]	0.51
DOM_MB_A	FWD	MB_CYC	[PJ]	0.048
DOM_MB_A	FWD	MB_HED_F	[PJ]	0.001
DOM_MB_A	FWD	MB_PELL	[PJ]	0.008
DOM_MB_A	FWD	NOC	[PJ]	0.069
DOM_MB_M	FWD	MB_CYC	[PJ]	0.054
DOM_MB_M	FWD	MB_HED_F	[PJ]	0.001
DOM_MB_M	FWD	MB_PELL	[PJ]	0.008
DOM_MB_M	FWD	NOC	[PJ]	0.077
DOM_PIT	FWD	NOC	[PJ]	0.353
DOM_SHB_A	FWD	NOC	[PJ]	0.023
DOM_SHB_A	FWD	SHB_HED	[PJ]	0.002
DOM_SHB_M	FWD	NOC	[PJ]	0.233
DOM_SHB_M	FWD	SHB_IMP_B	[PJ]	0.093
DOM_SHB_M	FWD	SHB_NEW_B	[PJ]	0.046
DOM_SHB_M	FWD	SHB_PELL	[PJ]	0.023
DOM_SHB_M	FWD	SHB_PLESP	[PJ]	0.002
DOM_STOVE_C	FWD	NOC	[PJ]	0.511
DOM_STOVE_C	FWD	STV_FAN_B	[PJ]	0
DOM_STOVE_C	FWD	STV_IMP_B	[PJ]	0.189
DOM_STOVE_C	FWD	STV_LPG	[PJ]	0
DOM_STOVE_C	FWD	STV_NEW_B	[PJ]	0.102
DOM_STOVE_H	FWD	NOC	[PJ]	0.486

DOM_STOVE_H	FWD	STV_ESP_B	[PJ]	0.021
DOM_STOVE_H	FWD	STV_IMP_B	[PJ]	0.18
DOM_STOVE_H	FWD	STV_NEW_B	[PJ]	0.097
DOM_STOVE_H	FWD	STV_PELL	[PJ]	0.024
DOM_STOVE_H	FWD	STV_PLESP	[PJ]	0.004
IN_BO_CHEM	GAS	NOC	[PJ]	0
IN_BO_CHEM	HF	GHIND	[PJ]	0.007
IN_BO_CHEM	HF	IN_HED	[PJ]	0
IN_BO_CHEM	HF	NOC	[PJ]	0.01
IN_BO_CHEM	LPG	NOC	[PJ]	0
IN_BO_CON	GAS	NOC	[PJ]	0
IN_BO_CON	HF	GHIND	[PJ]	0.007
IN_BO_CON	HF	IN_HED	[PJ]	0
IN_BO_CON	HF	NOC	[PJ]	0.01
IN_BO_CON	LPG	NOC	[PJ]	0
IN_BO_OTH	GAS	NOC	[PJ]	0
IN_BO_OTH	HF	GHIND	[PJ]	0.007
IN_BO_OTH	HF	IN_HED	[PJ]	0
IN_BO_OTH	HF	NOC	[PJ]	0.01
IN_BO_OTH	OS1	IN_CYC	[PJ]	0.129
IN_BO_OTH	OS1	IN_ESP1	[PJ]	0.013
IN_BO_OTH	OS1	IN_ESP2	[PJ]	0.007
IN_BO_OTH	OS1	IN_HED	[PJ]	0.002
IN_BO_OTH	OS1	NOC	[PJ]	0.185
IN_BO_OTH_L	HC1	IN_ESP1	[PJ]	0.014
IN_BO_OTH_L	HC1	IN_ESP2	[PJ]	0.008
IN_BO_OTH_L	HC1	IN_HED	[PJ]	0.002
IN_BO_OTH_L	HC1	IN_HED_SN	[PJ]	0.002
IN_BO_OTH_L	HC1	IN_WSCRB	[PJ]	0.014
IN_BO_OTH_L	HC1	NOC	[PJ]	0.196
IN_BO_PAP	GAS	NOC	[PJ]	0
IN_BO_PAP	HF	GHIND	[PJ]	0.007
IN_BO_PAP	HF	IN_HED	[PJ]	0
IN_BO_PAP	HF	NOC	[PJ]	0.01
IN_OC	DC	IN_CYC	[PJ]	0.011
IN_OC	DC	IN_ESP1	[PJ]	0.001
IN_OC	DC	IN_ESP2	[PJ]	0.001
IN_OC	DC	IN_HED	[PJ]	0
IN_OC	DC	NOC	[PJ]	0.015
IN_OC	GAS	NOC	[PJ]	0
IN_OC	HF	GHIND	[PJ]	0.007
IN_OC	HF	IN_HED	[PJ]	0
IN_OC	HF	NOC	[PJ]	0.01
IN_OC	MD	GHIND	[PJ]	0
IN_OC	MD	NOC	[PJ]	0

IN_OC3	HC1	IN_ESP1	[PJ]	0.013
IN_OC3	HC1	IN_ESP2	[PJ]	0.007
IN_OC3	HC1	IN_HED	[PJ]	0.002
IN_OC3	HC1	NOC	[PJ]	0.187
LEAD_GASOL	LFL	NOC	[PJ]	0.012
MINE_OTH	NOF	MINE_GP	[Mt]	0.004
MINE_OTH	NOF	NOC	[Mt]	0.005
MSW_URB_FOOD	NOF	NOC	[Mt waste]	0
MSW_URB_FOOD	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_FOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_FOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_FOOD	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_FOOD	NOF	UNC_BURN	[Mt waste]	8.74
MSW_URB_GLA	NOF	NOC	[Mt waste]	0
MSW_URB_MET	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_OTH	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_OTH	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_OTH	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_OTH	NOF	UNC_BURN	[Mt waste]	8.74
MSW_URB_PAP	NOF	NOC	[Mt waste]	0
MSW_URB_PAP	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_PAP	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_PAP	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_PAP	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_PAP	NOF	UNC_BURN	[Mt waste]	8.74
MSW_URB_PLA	NOF	NOC	[Mt waste]	0
MSW_URB_PLA	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_PLA	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_PLA	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_PLA	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_PLA	NOF	UNC_BURN	[Mt waste]	8.74
MSW_URB_TEX	NOF	NOC	[Mt waste]	0
MSW_URB_TEX	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_TEX	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_TEX	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_TEX	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_TEX	NOF	UNC_BURN	[Mt waste]	8.74
MSW_URB_WOOD	NOF	NOC	[Mt waste]	0
MSW_URB_WOOD	NOF	SWD_FL A	[Mt waste]	0.045
MSW_URB_WOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.748
MSW_URB_WOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.874
MSW_URB_WOOD	NOF	TREAT_INC	[Mt waste]	0.045
MSW_URB_WOOD	NOF	UNC_BURN	[Mt waste]	8.74

NONEN	GAS	NOC	[PJ]	0
NONEN	GSL	NOC	[PJ]	0
NONEN	HF	NOC	[PJ]	0
OTHER_PM	NOF	NOC	[kt PM (TSP)]	1
OTHER_PM	NOF	RESP1	[kt PM (TSP)]	1
PP_ENG	GAS	LFEUI	[PJ]	0.001
PP_ENG	GAS	LFEUII	[PJ]	0.001
PP_ENG	GAS	LFEUIII	[PJ]	0
PP_ENG	GAS	LFEUIV	[PJ]	0
PP_ENG	GAS	LFEUV	[PJ]	0
PP_ENG	GAS	LFEUVI	[PJ]	0
PP_ENG	GAS	NOC	[PJ]	0.002
PP_ENG	MD	NOC	[PJ]	0.096
PP_ENG	MD	TIWEUI	[PJ]	0.064
PP_ENG	MD	TIWEUII	[PJ]	0.048
PP_ENG	MD	TIWEUIII	[PJ]	0.014
PP_ENG	MD	TIWEUIV	[PJ]	0.003
PP_ENG	MD	TIWEUV	[PJ]	0.003
PP_ENG	MD	TIWEUVI	[PJ]	0.002
PP_EX_OTH	GAS	NOC	[PJ]	0
PP_EX_OTH	HF	FF	[PJ]	0
PP_EX_OTH	HF	GHIND	[PJ]	0.007
PP_EX_OTH	HF	HED	[PJ]	0
PP_EX_OTH	HF	NOC	[PJ]	0.009
PP_EX_OTH	MD	GHIND	[PJ]	0.001
PP_EX_OTH	MD	NOC	[PJ]	0.001
PP_EX_OTH	OS2	CYC	[PJ]	0.026
PP_EX_OTH	OS2	ESP1	[PJ]	0.003
PP_EX_OTH	OS2	ESP2	[PJ]	0.001
PP_EX_OTH	OS2	FF	[PJ]	0
PP_EX_OTH	OS2	FFSINJ	[PJ]	0
PP_EX_OTH	OS2	HED	[PJ]	0
PP_EX_OTH	OS2	NOC	[PJ]	0.037
PP_MOD	GAS	NOC	[PJ]	0
PP_NEW	GAS	NOC	[PJ]	0
PP_NEW	GSL	NOC	[PJ]	0.001
PP_NEW	HF	FF	[PJ]	0
PP_NEW	HF	GHIND	[PJ]	0.007
PP_NEW	HF	HED	[PJ]	0
PP_NEW	HF	NOC	[PJ]	0.009
PP_NEW	OS1	CYC	[PJ]	0.129
PP_NEW	OS1	ESP1	[PJ]	0.013
PP_NEW	OS1	ESP2	[PJ]	0.007
PP_NEW	OS1	FF	[PJ]	0.001
PP_NEW	OS1	HED	[PJ]	0.001

PP_NEW	OS1	NOC	[PJ]	0.185
PP_NEW	OS2	CYC	[PJ]	0.026
PP_NEW	OS2	ESP1	[PJ]	0.003
PP_NEW	OS2	ESP2	[PJ]	0.001
PP_NEW	OS2	FF	[PJ]	0
PP_NEW	OS2	FFSINJ	[PJ]	0
PP_NEW	OS2	HED	[PJ]	0
PP_NEW	OS2	NOC	[PJ]	0.037
PROD_AGAS	CRU	NOC	[PJ]	0.003
PROD_AGAS	CRU	REC_USE	[PJ]	0
PROD_AGAS	CRU	REC_USE2	[PJ]	0
PR_ALSEC	NOF	NOC	[Mt]	5.195
PR_ALSEC	NOF	PR_CYC	[Mt]	3.637
PR_ALSEC	NOF	PR_HED	[Mt]	0.052
PR_BAOX	NOF	NOC	[Mt]	10.45
PR_BAOX	NOF	PR_CYC	[Mt]	7.315
PR_BAOX	NOF	PR_ESP1	[Mt]	0.732
PR_BAOX	NOF	PR_ESP2	[Mt]	0.418
PR_BAOX	NOF	PR_HED	[Mt]	0.104
PR_BRICK	NOF	DDK	[Mt]	0
PR_BRICK	NOF	FCBTK	[Mt]	0
PR_BRICK	NOF	HK	[Mt]	0
PR_BRICK	NOF	IDK_BTK	[Mt]	0.32
PR_BRICK	NOF	MCBTK	[Mt]	0
PR_BRICK	NOF	MK	[Mt]	0
PR_BRICK	NOF	NOC	[Mt]	0.4
PR_BRICK	NOF	TK_COAL	[Mt]	0
PR_BRICK	NOF	TK_EOF	[Mt]	0.004
PR_BRICK	NOF	TK_OTHER	[Mt]	0
PR_BRICK	NOF	VSBK	[Mt]	0.2
PR_BRICK	NOF	ZIG_ZAG	[Mt]	0
PR_CAST	NOF	NOC	[Mt]	10.685
PR_CAST	NOF	PR_CYC	[Mt]	7.48
PR_CAST	NOF	PR_ESP1	[Mt]	0.748
PR_CAST	NOF	PR_ESP2	[Mt]	0.427
PR_CAST	NOF	PR_HED	[Mt]	0.107
PR_CAST_F	NOF	NOC	[Mt]	1.38
PR_CAST_F	NOF	PRF_GP1	[Mt]	0.828
PR_CAST_F	NOF	PRF_GP2	[Mt]	0.276
PR_CBLACK	NOF	NOC	[Mt]	1.44
PR_CBLACK	NOF	PR_CYC	[Mt]	1.008
PR_CBLACK	NOF	PR_HED	[Mt]	0.014
PR_CEM	NOF	NOC	[Mt]	2.142
PR_CEM	NOF	PR_CYC	[Mt]	1.499
PR_CEM	NOF	PR_ESP1	[Mt]	0.15

PR_CEM	NOF	PR_ESP2	[Mt]	0.086
PR_CEM	NOF	PR_HED	[Mt]	0.021
PR_CEM	NOF	PR_WSCRBB	[Mt]	0.15
PR_COKE	NOF	NOC	[Mt]	1.997
PR_COKE	NOF	PR_CYC	[Mt]	1.398
PR_COKE	NOF	PR_ESP1	[Mt]	0.14
PR_COKE	NOF	PR_ESP2	[Mt]	0.08
PR_COKE	NOF	PR_HED	[Mt]	0.02
PR_EARC	NOF	NOC	[Mt]	7.546
PR_EARC	NOF	PR_CYC	[Mt]	5.283
PR_EARC	NOF	PR_HED	[Mt]	0.075
PR_FERT	NOF	NOC	[Mt]	9
PR_FERT	NOF	PR_CYC	[Mt]	6.3
PR_FERT	NOF	PR_HED	[Mt]	0.09
PR_GLASS	NOF	NOC	[Mt]	2.958
PR_GLASS	NOF	PR_CYC	[Mt]	2.07
PR_GLASS	NOF	PR_ESP1	[Mt]	0.207
PR_GLASS	NOF	PR_ESP2	[Mt]	0.118
PR_GLASS	NOF	PR_HED	[Mt]	0.03
PR_LIME	NOF	NOC	[Mt]	1.4
PR_LIME	NOF	PR_CYC	[Mt]	0.98
PR_LIME	NOF	PR_ESP1	[Mt]	0.098
PR_LIME	NOF	PR_ESP2	[Mt]	0.056
PR_LIME	NOF	PR_HED	[Mt]	0.014
PR_LIME	NOF	PR_WSCRBB	[Mt]	0.098
PR_OTHER	NOF	NOC	[Mt]	1.47
PR_OTHER	NOF	PR_CYC	[Mt]	1.029
PR_OTHER	NOF	PR_ESP1	[Mt]	0.103
PR_OTHER	NOF	PR_ESP2	[Mt]	0.059
PR_OTHER	NOF	PR_HED	[Mt]	0.015
PR_PIGI	NOF	NOC	[Mt]	0.148
PR_PIGI	NOF	PR_CYC	[Mt]	0.104
PR_PIGI	NOF	PR_ESP1	[Mt]	0.01
PR_PIGI	NOF	PR_ESP2	[Mt]	0.006
PR_PIGI	NOF	PR_HED	[Mt]	0.001
PR_PIGI_F	NOF	NOC	[Mt]	0.15
PR_PIGI_F	NOF	PRF_GP1	[Mt]	0.09
PR_PIGI_F	NOF	PRF_GP2	[Mt]	0.03
PR_REF	NOF	NOC	[Mt]	0.096
PR_REF	NOF	PR_CYC	[Mt]	0.067
PR_REF	NOF	PR_ESP1	[Mt]	0.007
PR_REF	NOF	PR_ESP2	[Mt]	0.004
PR_REF	NOF	PR_HED	[Mt]	0.001
PR_SINT	NOF	NOC	[Mt]	0.557
PR_SINT	NOF	PR_CYC	[Mt]	0.39

PR_SINT	NOF	PR_ESP1	[Mt]	0.039
PR_SINT	NOF	PR_ESP2	[Mt]	0.022
PR_SINT	NOF	PR_HED	[Mt]	0.006
PR_SINT	NOF	PR_WSCRB	[Mt]	0.039
PR_SINT_F	NOF	NOC	[Mt]	0.104
PR_SINT_F	NOF	PRF_GP1	[Mt]	0.062
PR_SINT_F	NOF	PRF_GP2	[Mt]	0.021
PR_SMIND_F	NOF	NOC	[M people]	0.034
PR_SMIND_F	NOF	PRF_GP1	[M people]	0.02
PR_SMIND_F	NOF	PRF_GP2	[M people]	0.007
RES_BBQ	NOF	FILTER	[M people]	0.032
RES_BBQ	NOF	NOC	[M people]	0.035
RES_CIGAR	NOF	NOC	[M people]	0.017
RES_CREM	NOF	ELEC	[M people]	0
RES_CREM	NOF	FF	[M people]	0
RES_CREM	NOF	FFSINJ	[M people]	0
RES_CREM	NOF	NOC	[M people]	0
RES_CREM	NOF	STH_GP	[M people]	0
RES_FIREW	NOF	BAN	[M people]	0
RES_FIREW	NOF	NOC	[M people]	0.015
STH_AGR	NOF	NOC	[Mt]	0.004
STH_AGR	NOF	STH_GP	[Mt]	0.004
STH_COAL	NOF	NOC	[Mt]	0.004
STH_COAL	NOF	STH_GP	[Mt]	0.004
STH_FEORE	NOF	NOC	[Mt]	0.008
STH_FEORE	NOF	STH_GP	[Mt]	0.007
STH_NPK	NOF	NOC	[Mt]	0.004
STH_NPK	NOF	STH_GP	[Mt]	0.004
STH_OTH_IN	NOF	NOC	[Mt]	0.001
STH_OTH_IN	NOF	STH_GP	[Mt]	0.001
TRA_OT_S_L	GAS	NOC	[PJ]	0.001
TRA_OT_S_L	HF	NOC	[PJ]	0.113
TRA_OT_S_L	HF	STLHCM	[PJ]	0.113
TRA_OT_S_L	HF	STLSCR	[PJ]	0.113
TRA_OT_S_L	MD	NOC	[PJ]	0.026
TRA_OT_S_L	MD	STLMCM	[PJ]	0.026
TRA_OT_S_L	MD	STLSCR	[PJ]	0.026
TRA_OT_S_M	GAS	NOC	[PJ]	0.001
TRA_OT_S_M	MD	NOC	[PJ]	0.026
TRA_OT_S_M	MD	STMCM	[PJ]	0.026
TRA_OT_AGR	GSL	LFEUI	[PJ]	0.015
TRA_OT_AGR	GSL	LFEUII	[PJ]	0.015
TRA_OT_AGR	GSL	LFEUIII	[PJ]	0.005
TRA_OT_AGR	GSL	LFEUIV	[PJ]	0.005
TRA_OT_AGR	GSL	LFEUV	[PJ]	0.005

TRA_OT_AGR	GSL	LFEUVI	[PJ]	0.004
TRA_OT_AGR	GSL	NOC	[PJ]	0.028
TRA_OT_AGR	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_AGR	MD	CAGEUI	[PJ]	0.06
TRA_OT_AGR	MD	CAGEUII	[PJ]	0.027
TRA_OT_AGR	MD	CAGEUIII	[PJ]	0.027
TRA_OT_AGR	MD	CAGEUIII_PF	[PJ]	0.003
TRA_OT_AGR	MD	CAGEUIV	[PJ]	0.006
TRA_OT_AGR	MD	CAGEUV	[PJ]	0.006
TRA_OT_AGR	MD	CAGEUVI	[PJ]	0.002
TRA_OT_AGR	MD	NOC	[PJ]	0.106
TRA_OT_AGR	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_AIR	GSL	NOC	[PJ]	0
TRA_OT_CNS	MD	CAGEUI	[PJ]	0.057
TRA_OT_CNS	MD	CAGEUII	[PJ]	0.026
TRA_OT_CNS	MD	CAGEUIII	[PJ]	0.026
TRA_OT_CNS	MD	CAGEUIII_PF	[PJ]	0.003
TRA_OT_CNS	MD	CAGEUIV	[PJ]	0.006
TRA_OT_CNS	MD	CAGEUV	[PJ]	0.006
TRA_OT_CNS	MD	CAGEUVI	[PJ]	0.002
TRA_OT_CNS	MD	NOC	[PJ]	0.101
TRA_OT_CNS	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_CNS	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_INW	GSL	LFEUI	[PJ]	0.015
TRA_OT_INW	GSL	LFEUII	[PJ]	0.015
TRA_OT_INW	GSL	LFEUIII	[PJ]	0.005
TRA_OT_INW	GSL	LFEUIV	[PJ]	0.005
TRA_OT_INW	GSL	LFEUV	[PJ]	0.005
TRA_OT_INW	GSL	LFEUVI	[PJ]	0.004
TRA_OT_INW	GSL	NOC	[PJ]	0.028
TRA_OT_INW	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	GAS	LFEUI	[PJ]	0.015
TRA_OT_LB	GAS	LFEUII	[PJ]	0.015
TRA_OT_LB	GAS	LFEUIII	[PJ]	0.005
TRA_OT_LB	GAS	LFEUIV	[PJ]	0.005
TRA_OT_LB	GAS	LFEUV	[PJ]	0.005
TRA_OT_LB	GAS	LFEUVI	[PJ]	0.004
TRA_OT_LB	GAS	NOC	[PJ]	0.028
TRA_OT_LB	GSL	LFEUI	[PJ]	0.015
TRA_OT_LB	GSL	LFEUII	[PJ]	0.015

TRA_OT_LB	GSL	LFEUIII	[PJ]	0.005
TRA_OT_LB	GSL	LFEUIV	[PJ]	0.005
TRA_OT_LB	GSL	LFEUV	[PJ]	0.005
TRA_OT_LB	GSL	LFEUVI	[PJ]	0.004
TRA_OT_LB	GSL	NOC	[PJ]	0.028
TRA_OT_LB	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LB	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	MD	HDEUI	[PJ]	0.069
TRA_OT_LB	MD	HDEUII	[PJ]	0.037
TRA_OT_LB	MD	HDEUIII	[PJ]	0.03
TRA_OT_LB	MD	HDEUIV	[PJ]	0.006
TRA_OT_LB	MD	HDEUV	[PJ]	0.006
TRA_OT_LB	MD	HDEUVI	[PJ]	0
TRA_OT_LB	MD	NOC	[PJ]	0.09
TRA_OT_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_RAI	H2	NOC	[PJ]	0
TRA_OT_RAI	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_RAI	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDB	ABRASION	NOC	[Gvkm]	0.021
TRA_RD_HDB	BRAKE	NOC	[Gvkm]	0.016
TRA_RD_HDB	GAS	HDSEI	[PJ]	0.001
TRA_RD_HDB	GAS	HDSEII	[PJ]	0
TRA_RD_HDB	GAS	HDSEIII	[PJ]	0
TRA_RD_HDB	GAS	NOC	[PJ]	0.002
TRA_RD_HDB	MD	HDEUI	[PJ]	0.044
TRA_RD_HDB	MD	HDEUII	[PJ]	0.024
TRA_RD_HDB	MD	HDEUIII	[PJ]	0.02
TRA_RD_HDB	MD	HDEUIV	[PJ]	0.004
TRA_RD_HDB	MD	HDEUV	[PJ]	0.004
TRA_RD_HDB	MD	HDEUVI	[PJ]	0
TRA_RD_HDB	MD	HDEUVII	[PJ]	0
TRA_RD_HDB	MD	NOC	[PJ]	0.058
TRA_RD_HDB	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDB	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDB	TYRE	NOC	[Gvkm]	0.004
TRA_RD_HDT	ABRASION	NOC	[Gvkm]	0.021
TRA_RD_HDT	BRAKE	NOC	[Gvkm]	0.016
TRA_RD_HDT	GAS	HDSEI	[PJ]	0.001
TRA_RD_HDT	GAS	HDSEII	[PJ]	0
TRA_RD_HDT	GAS	HDSEIII	[PJ]	0
TRA_RD_HDT	GAS	NOC	[PJ]	0.002
TRA_RD_HDT	MD	HDEUI	[PJ]	0.044
TRA_RD_HDT	MD	HDEUII	[PJ]	0.024
TRA_RD_HDT	MD	HDEUIII	[PJ]	0.02

TRA_RD_HDT	MD	HDEUIV	[PJ]	0.004
TRA_RD_HDT	MD	HDEUV	[PJ]	0.004
TRA_RD_HDT	MD	HDEUVI	[PJ]	0
TRA_RD_HDT	MD	HDEUVII	[PJ]	0
TRA_RD_HDT	MD	NOC	[PJ]	0.058
TRA_RD_HDT	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDT	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDT	TYRE	NOC	[Gvkm]	0.004
TRA_RD_LD2	ABRASION	NOC	[Gvkm]	0.002
TRA_RD_LD2	BRAKE	NOC	[Gvkm]	0
TRA_RD_LD2	GSL	MMO2I	[PJ]	0.06
TRA_RD_LD2	GSL	MMO2II	[PJ]	0.037
TRA_RD_LD2	GSL	MMO2III	[PJ]	0.015
TRA_RD_LD2	GSL	MMO2IV	[PJ]	0.015
TRA_RD_LD2	GSL	MMO2V	[PJ]	0.015
TRA_RD_LD2	GSL	MMO2VI	[PJ]	0.015
TRA_RD_LD2	GSL	NOC	[PJ]	0.186
TRA_RD_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD2	TYRE	NOC	[Gvkm]	0
TRA_RD_LD4C	ABRASION	NOC	[Gvkm]	0.004
TRA_RD_LD4C	BRAKE	NOC	[Gvkm]	0.003
TRA_RD_LD4C	GAS	LFEUI	[PJ]	0.001
TRA_RD_LD4C	GAS	LFEUII	[PJ]	0.001
TRA_RD_LD4C	GAS	LFEUIII	[PJ]	0
TRA_RD_LD4C	GAS	LFEUIV	[PJ]	0
TRA_RD_LD4C	GAS	LFEUV	[PJ]	0
TRA_RD_LD4C	GAS	LFEUVI	[PJ]	0
TRA_RD_LD4C	GAS	NOC	[PJ]	0.002
TRA_RD_LD4C	GSL	LFEUI	[PJ]	0.003
TRA_RD_LD4C	GSL	LFEUII	[PJ]	0.004
TRA_RD_LD4C	GSL	LFEUIII	[PJ]	0.001
TRA_RD_LD4C	GSL	LFEUIV	[PJ]	0.001
TRA_RD_LD4C	GSL	LFEUV	[PJ]	0.001
TRA_RD_LD4C	GSL	LFEUVI	[PJ]	0.001
TRA_RD_LD4C	GSL	NOC	[PJ]	0.004
TRA_RD_LD4C	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	LPG	LFEUI	[PJ]	0.001
TRA_RD_LD4C	LPG	LFEUII	[PJ]	0.001
TRA_RD_LD4C	LPG	LFEUIII	[PJ]	0
TRA_RD_LD4C	LPG	LFEUIV	[PJ]	0
TRA_RD_LD4C	LPG	LFEUV	[PJ]	0
TRA_RD_LD4C	LPG	LFEUVI	[PJ]	0
TRA_RD_LD4C	LPG	NOC	[PJ]	0.002

TRA_RD_LD4C	MD	MDEUI	[PJ]	0.03
TRA_RD_LD4C	MD	MDEUII	[PJ]	0.021
TRA_RD_LD4C	MD	MDEUIII	[PJ]	0.017
TRA_RD_LD4C	MD	MDEUIV	[PJ]	0.016
TRA_RD_LD4C	MD	MDEUV	[PJ]	0.001
TRA_RD_LD4C	MD	MDEUVI	[PJ]	0.001
TRA_RD_LD4C	MD	MDEUVII	[PJ]	0
TRA_RD_LD4C	MD	MDEUVip	[PJ]	0.001
TRA_RD_LD4C	MD	MDEUVit	[PJ]	0.001
TRA_RD_LD4C	MD	NOC	[PJ]	0.087
TRA_RD_LD4C	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	TYRE	NOC	[Gvkm]	0.001
TRA_RD_LD4T	ABRASION	NOC	[Gvkm]	0.004
TRA_RD_LD4T	BRAKE	NOC	[Gvkm]	0.003
TRA_RD_LD4T	GAS	LFEUI	[PJ]	0.001
TRA_RD_LD4T	GAS	LFEUII	[PJ]	0.001
TRA_RD_LD4T	GAS	LFEUIII	[PJ]	0
TRA_RD_LD4T	GAS	LFEUIV	[PJ]	0
TRA_RD_LD4T	GAS	LFEUV	[PJ]	0
TRA_RD_LD4T	GAS	LFEUVI	[PJ]	0
TRA_RD_LD4T	GAS	NOC	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUI	[PJ]	0.03
TRA_RD_LD4T	MD	MDEUII	[PJ]	0.021
TRA_RD_LD4T	MD	MDEUIII	[PJ]	0.017
TRA_RD_LD4T	MD	MDEUIV	[PJ]	0.016
TRA_RD_LD4T	MD	MDEUV	[PJ]	0.001
TRA_RD_LD4T	MD	MDEUVI	[PJ]	0.001
TRA_RD_LD4T	MD	MDEUVII	[PJ]	0
TRA_RD_LD4T	MD	MDEUVip	[PJ]	0.001
TRA_RD_LD4T	MD	MDEUVit	[PJ]	0.001
TRA_RD_LD4T	MD	NOC	[PJ]	0.087
TRA_RD_LD4T	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4T	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4T	TYRE	NOC	[Gvkm]	0.001
TRA_RD_M4	ABRASION	NOC	[Gvkm]	0.002
TRA_RD_M4	BRAKE	NOC	[Gvkm]	0
TRA_RD_M4	GSL	MOT4I	[PJ]	0.005
TRA_RD_M4	GSL	MOT4II	[PJ]	0.004
TRA_RD_M4	GSL	MOT4III	[PJ]	0.002
TRA_RD_M4	GSL	MOT4IV	[PJ]	0.002
TRA_RD_M4	GSL	MOT4V	[PJ]	0
TRA_RD_M4	GSL	MOT4VI	[PJ]	0
TRA_RD_M4	GSL	NOC	[PJ]	0.006
TRA_RD_M4	GSL_NV_HE	NOC	[thousand of vehicles]	0

TRA_RD_M4	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_M4	TYRE	NOC	[Gvkm]	0
WASTE_AGR	NOF	BAN	[Mt]	0
WASTE_AGR	NOF	NOC	[Mt]	9.264
WASTE_FLR	NOF	FLR_GP	[PJ]	0.002
WASTE_FLR	NOF	NOC	[PJ]	0.032

Table A9.7 – Emission factors for NMVOC emissions.

Sector	Fuel activity	Measure	Unit	FACTOR_ABTD [kt/unit]
AUTO_P_NEW	VEH	A_INC	[kveh]	0.003
AUTO_P_NEW	VEH	NOC	[kveh]	0.004
COIL	SC	INC	[mln m2]	0.004
COIL	SC	M_INC	[mln m2]	0.001
COIL	SC	NOC	[mln m2]	0.043
COIL	SC	POWDER	[mln m2]	0
CON_COMB	GAS	NOC	[PJ]	0.003
CON_COMB	HC1	NOC	[PJ]	0.015
CON_COMB	HF	NOC	[PJ]	0.005
DECO_P	PNT	BASE	[kt]	0.141
DECO_P	PNT	FSED	[kt]	0.062
DECO_P	PNT	NOC	[kt]	0.165
DECO_P	PNT	SED	[kt]	0.07
DEGR	SLV	ACA	[kt SLV]	0.142
DEGR	SLV	BEMT+ACA	[kt SLV]	0.106
DEGR	SLV	BEMT	[kt SLV]	0.532
DEGR	SLV	CLSD_A3+ACA	[kt SLV]	0.02
DEGR	SLV	CLSD_A3	[kt SLV]	0.025
DEGR	SLV	CLSD_CL	[kt SLV]	0.035
DEGR	SLV	CLSD_HF+ACA	[kt SLV]	0.02
DEGR	SLV	CLSD_HF	[kt SLV]	0.025
DEGR	SLV	COLD	[kt SLV]	0.08
DEGR	SLV	NOC	[kt SLV]	0.9
DEGR	SLV	WBD	[kt SLV]	0
DOM	GAS	NOC	[PJ]	0.002
DOM	LPG	NOC	[PJ]	0.003
DOM	MD	NOC	[PJ]	0.003
DOM_FPLACE	FWD	FP_IMP	[PJ]	0.518
DOM_FPLACE	FWD	FP_NEW	[PJ]	0.518
DOM_FPLACE	FWD	NOC	[PJ]	0.92
DOM_MB_A	FWD	MB_CYC	[PJ]	0.05
DOM_MB_A	FWD	MB_HED_F	[PJ]	0.015
DOM_MB_A	FWD	MB_PELL	[PJ]	0.015
DOM_MB_A	FWD	NOC	[PJ]	0.05

DOM_MB_M	FWD	MB_CYC	[PJ]	0.1
DOM_MB_M	FWD	MB_HED_F	[PJ]	0.015
DOM_MB_M	FWD	MB_PELL	[PJ]	0.015
DOM_MB_M	FWD	NOC	[PJ]	0.1
DOM_OS	POP	NOC	[M people]	2.183
DOM_OS	POP	REF	[M people]	1.4
DOM_OS	POP	REF1	[M people]	1.59
DOM_OS	POP	REF2	[M people]	1.27
DOM_OS	POP	REF3	[M people]	0.71
DOM_PIT	FWD	NOC	[PJ]	2
DOM_SHB_A	FWD	NOC	[PJ]	0.015
DOM_SHB_A	FWD	SHB_HED	[PJ]	0.015
DOM_SHB_M	FWD	NOC	[PJ]	0.4
DOM_SHB_M	FWD	SHB_IMP_B	[PJ]	0.14
DOM_SHB_M	FWD	SHB_NEW_B	[PJ]	0.012
DOM_SHB_M	FWD	SHB_PELL	[PJ]	0.002
DOM_SHB_M	FWD	SHB_PLESP	[PJ]	0.002
DOM_STOVE_C	FWD	NOC	[PJ]	1.6
DOM_STOVE_C	FWD	STV_FAN_B	[PJ]	0.08
DOM_STOVE_C	FWD	STV_IMP_B	[PJ]	0.24
DOM_STOVE_C	FWD	STV_LPG	[PJ]	0
DOM_STOVE_C	FWD	STV_NEW_B	[PJ]	0.08
DOM_STOVE_H	FWD	NOC	[PJ]	0.7
DOM_STOVE_H	FWD	STV_ESP_B	[PJ]	0.686
DOM_STOVE_H	FWD	STV_IMP_B	[PJ]	0.25
DOM_STOVE_H	FWD	STV_NEW_B	[PJ]	0.065
DOM_STOVE_H	FWD	STV_PELL	[PJ]	0.02
DOM_STOVE_H	FWD	STV_PLESP	[PJ]	0.02
DRY_NEW	TEX	ACA	[kt TEX]	0.016
DRY_NEW	TEX	HCM	[kt TEX]	0.009
DRY_NEW	TEX	NCCM	[kt TEX]	0.009
DRY_NEW	TEX	NOC	[kt TEX]	0.019
DRY_NEW	TEX	WCLEAN	[kt TEX]	0
D_GASST	GSL	NOC	[PJ]	0.06
D_GASST	GSL	ST(II+IB)	[PJ]	0.016
D_GASST	GSL	ST_IB	[PJ]	0.034
D_GASST	GSL	ST_II	[PJ]	0.041
D_REFDEP	GSL	IFC+ST_IAD	[PJ]	0.004
D_REFDEP	GSL	IFC+ST_IAS	[PJ]	0.005
D_REFDEP	GSL	IFC	[PJ]	0.031
D_REFDEP	GSL	NOC	[PJ]	0.047
D_REFDEP	GSL	ST_IAD	[PJ]	0.019
D_REFDEP	GSL	ST_IAS	[PJ]	0.02
D_REFDEP	MD	NOC	[PJ]	0.001
D_REFDEP_S	GSL	IFC+ST_IAD	[PJ]	0.004

D_REFDEP_S	GSL	IFC+ST_IAS	[PJ]	0.005
D_REFDEP_S	GSL	IFC	[PJ]	0.031
D_REFDEP_S	GSL	NOC	[PJ]	0.047
D_REFDEP_S	GSL	ST_IAD	[PJ]	0.019
D_REFDEP_S	GSL	ST_IAS	[PJ]	0.02
D_REFDEP_S	MD	NOC	[PJ]	0.001
EXD_GAS	EMI	NOC	[kt VOC]	1
EXD_LQ	EMI	F_IMP	[kt VOC]	1
EXD_LQ	EMI	NOC	[kt VOC]	1
EXD_LQ	EMI	VBAL+VF_ALT	[kt VOC]	1
EXD_LQ	EMI	VBAL+V_ALT	[kt VOC]	1
EXD_LQ	EMI	VBAL	[kt VOC]	1
EXD_LQ	EMI	V_ALT	[kt VOC]	1
FATOIL	SD	ACA	[kt]	0.005
FATOIL	SD	NOC	[kt]	0.018
FATOIL	SD	SHM+ACA	[kt]	0.004
FATOIL	SD	SHM+ACAN	[kt]	0.001
FOOD	POP	INC	[M people]	0.294
FOOD	POP	NOC	[M people]	0.46
GLUE_INH	ADH	ACA	[kt]	0.187
GLUE_INH	ADH	INC	[kt]	0.187
GLUE_INH	ADH	NOC	[kt]	0.78
GLUE_INT	ADH	ACA	[kt]	0.187
GLUE_INT	ADH	EMU	[kt]	0.014
GLUE_INT	ADH	HOTM	[kt]	0
GLUE_INT	ADH	INC	[kt]	0.187
GLUE_INT	ADH	NOC	[kt]	0.78
IND_OS	EMI	NAGR	[kt VOC]	1
IND_OS	EMI	NOC	[kt VOC]	1
IND_OS	EMI	PRM+NAGR	[kt VOC]	1
IND_OS	EMI	PRM	[kt VOC]	0.3
IND_OTH	EMI	BISUB	[kt VOC]	0.724
IND_OTH	EMI	HSE+BISUB	[kt VOC]	0.424
IND_OTH	EMI	HSE	[kt VOC]	0.7
IND_OTH	EMI	NOC	[kt VOC]	1
IND_P_CNT	PNT	INC	[kt]	0.166
IND_P_CNT	PNT	ISBP+INC	[kt]	0.104
IND_P_CNT	PNT	ISBP	[kt]	0.432
IND_P_CNT	PNT	NOC	[kt]	0.69
IND_P_CNT	PNT	POWDER	[kt]	0
IND_P_CNT	PNT	WBP	[kt]	0.033
IND_P_OT	PNT	CSBP+INC	[kt]	0.089
IND_P_OT	PNT	CSBP	[kt]	0.37
IND_P_OT	PNT	INC	[kt]	0.179
IND_P_OT	PNT	ISBP+INC	[kt]	0.046

IND_P_OT	PNT	ISBP	[kt]	0.19
IND_P_OT	PNT	NOC	[kt]	0.748
IND_P_OT	PNT	POWDER	[kt]	0
IND_P_OT	PNT	WBP	[kt]	0.031
IND_P_PL	PNT	CSBP+INC	[kt]	0.073
IND_P_PL	PNT	CSBP	[kt]	0.305
IND_P_PL	PNT	INC	[kt]	0.179
IND_P_PL	PNT	ISBP+INC	[kt]	0.06
IND_P_PL	PNT	ISBP	[kt]	0.249
IND_P_PL	PNT	NOC	[kt]	0.748
IND_P_PL	PNT	POWDER	[kt]	0
IND_P_PL	PNT	TSBP_IA+INC	[kt]	0.097
IND_P_PL	PNT	TSBP_IA	[kt]	0.403
IND_P_PL	PNT	WBP	[kt]	0.027
INORG	EMI	NOC	[kt VOC]	1
IN_BO_CHEM	GAS	NOC	[PJ]	0.002
IN_BO_CHEM	HF	NOC	[PJ]	0.005
IN_BO_CON	GAS	NOC	[PJ]	0.002
IN_BO_CON	HF	NOC	[PJ]	0.005
IN_BO_OTH	GAS	NOC	[PJ]	0.002
IN_BO_OTH	HF	NOC	[PJ]	0.005
IN_BO_OTH	OS1	NOC	[PJ]	0.048
IN_BO_OTH_L	HC1	NOC	[PJ]	0.01
IN_BO_PAP	GAS	NOC	[PJ]	0.002
IN_BO_PAP	HF	NOC	[PJ]	0.005
IN_OC	DC	NOC	[PJ]	0.01
IN_OC	GAS	NOC	[PJ]	0.002
IN_OC	HC1	NOC	[PJ]	0.01
IN_OC	HF	NOC	[PJ]	0.005
IN_OC	MD	NOC	[PJ]	0.004
LEATHER	CTG	BIO	[kt]	0.194
LEATHER	CTG	INC	[kt]	0.194
LEATHER	CTG	NOC	[kt]	1.02
LEATHER	CTG	WBC_L	[kt]	0.36
MSW_URB_FOOD	NOF	NOC	[Mt waste]	0
MSW_URB_FOOD	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_FOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.692
MSW_URB_FOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_FOOD	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_FOOD	NOF	UNC_BURN	[Mt waste]	8.46
MSW_URB_GLA	NOF	NOC	[Mt waste]	0
MSW_URB_MET	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	NOC	[Mt waste]	0
MSW_URB_OTH	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_OTH	NOF	SWD_UNM_HIGH	[Mt waste]	1.692

MSW_URB_OTH	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_OTH	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_OTH	NOF	UNC_BURN	[Mt waste]	8.46
MSW_URB_PAP	NOF	NOC	[Mt waste]	0
MSW_URB_PAP	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_PAP	NOF	SWD_UNM_HIGH	[Mt waste]	1.692
MSW_URB_PAP	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_PAP	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_PAP	NOF	UNC_BURN	[Mt waste]	8.46
MSW_URB_PLA	NOF	NOC	[Mt waste]	0
MSW_URB_PLA	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_PLA	NOF	SWD_UNM_HIGH	[Mt waste]	1.692
MSW_URB_PLA	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_PLA	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_PLA	NOF	UNC_BURN	[Mt waste]	8.46
MSW_URB_TEX	NOF	NOC	[Mt waste]	0
MSW_URB_TEX	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_TEX	NOF	SWD_UNM_HIGH	[Mt waste]	1.692
MSW_URB_TEX	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_TEX	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_TEX	NOF	UNC_BURN	[Mt waste]	8.46
MSW_URB_WOOD	NOF	NOC	[Mt waste]	0
MSW_URB_WOOD	NOF	SWD_FLA	[Mt waste]	0.002
MSW_URB_WOOD	NOF	SWD_UNM_HIGH	[Mt waste]	1.692
MSW_URB_WOOD	NOF	SWD_UNM_LOW	[Mt waste]	0.846
MSW_URB_WOOD	NOF	TREAT_INC	[Mt waste]	0.002
MSW_URB_WOOD	NOF	UNC_BURN	[Mt waste]	8.46
OTH_ORG_PR	EMI	LDAR_I	[kt VOC]	0.66
OTH_ORG_PR	EMI	LDAR_II	[kt VOC]	0.64
OTH_ORG_PR	EMI	LDAR_III	[kt VOC]	0.64
OTH_ORG_PR	EMI	LDAR_IV	[kt VOC]	0.53
OTH_ORG_PR	EMI	NOC	[kt VOC]	1
PHARMA	SLV	NOC	[kt SLV]	0.3
PHARMA	SLV	PRM1+LEOP	[kt SLV]	0.08
PHARMA	SLV	PRM2+HEOP	[kt SLV]	0.035
PIS	PG	BEMT	[kt PG]	0.007
PIS	PG	NOC	[kt PG]	0.014
PIS	PG	UPACA	[kt PG]	0.005
PLSTYR_PR	EPS	INC	[kt]	0.039
PLSTYR_PR	EPS	LPB	[kt]	0.04
PLSTYR_PR	EPS	NOC	[kt]	0.06
PLSTYR_PR	EPS	PB+REC+INC	[kt]	0.033
PLSTYR_PR	EPS	PB+REC	[kt]	0.051
PP_ENG	GAS	LFEUI	[PJ]	0.043
PP_ENG	GAS	LFEUII	[PJ]	0.019

PP_ENG	GAS	LFEUIII	[PJ]	0.012
PP_ENG	GAS	LFEUIV	[PJ]	0.006
PP_ENG	GAS	LFEUV	[PJ]	0.004
PP_ENG	GAS	LFEUVI	[PJ]	0.003
PP_ENG	GAS	NOC	[PJ]	0.15
PP_ENG	MD	NOC	[PJ]	0.06
PP_ENG	MD	TIWEUI	[PJ]	0.039
PP_ENG	MD	TIWEUII	[PJ]	0.032
PP_ENG	MD	TIWEUIII	[PJ]	0.024
PP_ENG	MD	TIWEUIV	[PJ]	0.024
PP_ENG	MD	TIWEUV	[PJ]	0.023
PP_ENG	MD	TIWEUVI	[PJ]	0.006
PP_EX_OTH	GAS	NOC	[PJ]	0.002
PP_EX_OTH	HF	NOC	[PJ]	0.005
PP_EX_OTH	MD	NOC	[PJ]	0.004
PP_EX_OTH	OS2	NOC	[PJ]	0.015
PP_MOD	GAS	NOC	[PJ]	0.001
PP_NEW	GAS	NOC	[PJ]	0.001
PP_NEW	GSL	NOC	[PJ]	0.002
PP_NEW	HF	NOC	[PJ]	0.002
PP_NEW	OS1	NOC	[PJ]	0.01
PP_NEW	OS2	NOC	[PJ]	0.01
PROD_AGAS	CRU	NOC	[PJ]	0
PROD_AGAS	CRU	REC_USE	[PJ]	0
PROD_AGAS	CRU	REC_USE2	[PJ]	0
PRT_OFFS_NEW	INK	INC	[kt INK]	0.154
PRT_OFFS_NEW	INK	NOC	[kt INK]	0.45
PRT_PACK_NEW	INK	INC	[kt INK]	0.155
PRT_PACK_NEW	INK	NOC	[kt INK]	0.4
PRT_PACK_NEW	INK	WBI+INC	[kt INK]	0.13
PRT_PACK_NEW	INK	WBI	[kt INK]	0.315
PRT_PUB_NEW	INK	NOC	[kt INK]	0.182
PRT_PUB_NEW	INK	WBI	[kt INK]	0.156
PRT_SCR_NEW	INK	ENC+INC	[kt INK]	0.198
PRT_SCR_NEW	INK	NOC	[kt INK]	0.36
PRT_SCR_NEW	INK	WBI+BIO	[kt INK]	0.259
PRT_SCR_NEW	INK	WBI+ENC+INC	[kt INK]	0.076
PRT_SCR_NEW	INK	WBI+ENC+INC_BIO	[kt INK]	0.06
PRT_SCR_NEW	INK	WBI	[kt INK]	0.117
PR_BRICK	NOF	DDK	[Mt]	0.15
PR_BRICK	NOF	FCBTK	[Mt]	0.1
PR_BRICK	NOF	HK	[Mt]	0.013
PR_BRICK	NOF	IDK_BTK	[Mt]	0.1
PR_BRICK	NOF	MCBTK	[Mt]	0.15
PR_BRICK	NOF	MK	[Mt]	0.04

PR_BRICK	NOF	NOC	[Mt]	0.15
PR_BRICK	NOF	TK_COAL	[Mt]	0.016
PR_BRICK	NOF	TK_EOF	[Mt]	0.016
PR_BRICK	NOF	TK_OTHER	[Mt]	0.01
PR_BRICK	NOF	VSBK	[Mt]	0.08
PR_BRICK	NOF	ZIG_ZAG	[Mt]	0.1
PR_REF	CRU	COWS	[Mt]	0.274
PR_REF	CRU	LDAR_I+COWS	[Mt]	0.149
PR_REF	CRU	LDAR_I	[Mt]	0.185
PR_REF	CRU	LDAR_II+COWS	[Mt]	0.124
PR_REF	CRU	LDAR_II	[Mt]	0.16
PR_REF	CRU	NOC	[Mt]	0.31
SHOE	SHO	BIO	[mIn pairs]	0.017
SHOE	SHO	INC	[mIn pairs]	0.017
SHOE	SHO	NOC	[mIn pairs]	0.06
SHOE	SHO	SPRM+BIO	[mIn pairs]	0.009
SHOE	SHO	SPRM+INC	[mIn pairs]	0.009
SHOE	SHO	SPRM	[mIn pairs]	0.031
SHOE	SHO	SPRMPLUS	[mIn pairs]	0.023
STCRACK_PR	EP	LDAR_I	[kt]	0.002
STCRACK_PR	EP	LDAR_II	[kt]	0.002
STCRACK_PR	EP	LDAR_III	[kt]	0.002
STCRACK_PR	EP	LDAR_IV	[kt]	0.002
STCRACK_PR	EP	NOC	[kt]	0.003
SYNTH_RUB	RUB	INC	[kt]	0.005
SYNTH_RUB	RUB	NOC	[kt]	0.021
SYNTH_RUB	RUB	SUB1_VT+INC	[kt]	0.002
SYNTH_RUB	RUB	SUB1_VT	[kt]	0.008
SYNTH_RUB	RUB	SUB2_VT	[kt]	0.005
SYNTH_RUB	RUB	SUB_V+INC	[kt]	0.003
SYNTH_RUB	RUB	SUB_V	[kt]	0.014
TRA_OT_S_L	GAS	NOC	[PJ]	0
TRA_OT_S_L	HF	NOC	[PJ]	0.03
TRA_OT_S_L	MD	NOC	[PJ]	0.03
TRA_OT_S_M	GAS	NOC	[PJ]	0
TRA_OT_S_M	MD	NOC	[PJ]	0.03
TRA_OT_AGR	GSL	LFEUI	[PJ]	0.218
TRA_OT_AGR	GSL	LFEUII	[PJ]	0.098
TRA_OT_AGR	GSL	LFEUIII	[PJ]	0.06
TRA_OT_AGR	GSL	LFEUIV	[PJ]	0.03
TRA_OT_AGR	GSL	LFEUV	[PJ]	0.023
TRA_OT_AGR	GSL	LFEUVI	[PJ]	0.015
TRA_OT_AGR	GSL	NOC	[PJ]	0.752
TRA_OT_AGR	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0

TRA_OT_AGR	MD	CAGEUI	[PJ]	0.13
TRA_OT_AGR	MD	CAGEUII	[PJ]	0.116
TRA_OT_AGR	MD	CAGEUIII	[PJ]	0.08
TRA_OT_AGR	MD	CAGEUIII_PF	[PJ]	0.08
TRA_OT_AGR	MD	CAGEUIV	[PJ]	0.08
TRA_OT_AGR	MD	CAGEUV	[PJ]	0.076
TRA_OT_AGR	MD	CAGEUVI	[PJ]	0.019
TRA_OT_AGR	MD	NOC	[PJ]	0.2
TRA_OT_AGR	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_AGR	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_AIR	GSL	NOC	[PJ]	0.005
TRA_OT_CNS	MD	CAGEUI	[PJ]	0.13
TRA_OT_CNS	MD	CAGEUII	[PJ]	0.116
TRA_OT_CNS	MD	CAGEUIII	[PJ]	0.08
TRA_OT_CNS	MD	CAGEUIII_PF	[PJ]	0.08
TRA_OT_CNS	MD	CAGEUIV	[PJ]	0.08
TRA_OT_CNS	MD	CAGEUV	[PJ]	0.076
TRA_OT_CNS	MD	CAGEUVI	[PJ]	0.019
TRA_OT_CNS	MD	NOC	[PJ]	0.2
TRA_OT_CNS	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_CNS	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_EV	GSL	NOC	[PJ]	0.154
TRA_OT_INW	GSL	LFEUI	[PJ]	2.9
TRA_OT_INW	GSL	LFEUII	[PJ]	1.3
TRA_OT_INW	GSL	LFEUIII	[PJ]	0.8
TRA_OT_INW	GSL	LFEUIV	[PJ]	0.4
TRA_OT_INW	GSL	LFEUV	[PJ]	0.3
TRA_OT_INW	GSL	LFEUVI	[PJ]	0.2
TRA_OT_INW	GSL	NOC	[PJ]	10
TRA_OT_INW	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_INW	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	GAS	LFEUI	[PJ]	0.009
TRA_OT_LB	GAS	LFEUII	[PJ]	0.004
TRA_OT_LB	GAS	LFEUIII	[PJ]	0.003
TRA_OT_LB	GAS	LFEUIV	[PJ]	0.001
TRA_OT_LB	GAS	LFEUV	[PJ]	0.001
TRA_OT_LB	GAS	LFEUVI	[PJ]	0.001
TRA_OT_LB	GAS	NOC	[PJ]	0.032
TRA_OT_LB	GSL	LFEUI	[PJ]	0.218
TRA_OT_LB	GSL	LFEUII	[PJ]	0.098
TRA_OT_LB	GSL	LFEUIII	[PJ]	0.06
TRA_OT_LB	GSL	LFEUIV	[PJ]	0.03
TRA_OT_LB	GSL	LFEUV	[PJ]	0.023

TRA_OT_LB	GSL	LFEUVI	[PJ]	0.015
TRA_OT_LB	GSL	NOC	[PJ]	0.752
TRA_OT_LB	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LB	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_LB	MD	HDEUI	[PJ]	0.14
TRA_OT_LB	MD	HDEUII	[PJ]	0.102
TRA_OT_LB	MD	HDEUIII	[PJ]	0.086
TRA_OT_LB	MD	HDEUIV	[PJ]	0.08
TRA_OT_LB	MD	HDEUV	[PJ]	0.07
TRA_OT_LB	MD	HDEUVI	[PJ]	0.021
TRA_OT_LB	MD	NOC	[PJ]	0.2
TRA_OT_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_OT_RAI	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_OT_RAI	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_EV	GSL	NOC	[PJ]	0
TRA_RD_HDB	GAS	HDSEI	[PJ]	0.357
TRA_RD_HDB	GAS	HDSEII	[PJ]	0.24
TRA_RD_HDB	GAS	HDSEIII	[PJ]	0.066
TRA_RD_HDB	GAS	NOC	[PJ]	0.53
TRA_RD_HDB	MD	HDEUI	[PJ]	0.059
TRA_RD_HDB	MD	HDEUII	[PJ]	0.04
TRA_RD_HDB	MD	HDEUIII	[PJ]	0.034
TRA_RD_HDB	MD	HDEUIV	[PJ]	0.002
TRA_RD_HDB	MD	HDEUV	[PJ]	0.002
TRA_RD_HDB	MD	HDEUVI	[PJ]	0.002
TRA_RD_HDB	MD	HDEUVII	[PJ]	0.002
TRA_RD_HDB	MD	NOC	[PJ]	0.072
TRA_RD_HDB	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDB	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_HDT	GAS	HDSEI	[PJ]	0.161
TRA_RD_HDT	GAS	HDSEII	[PJ]	0.079
TRA_RD_HDT	GAS	HDSEIII	[PJ]	0.041
TRA_RD_HDT	GAS	NOC	[PJ]	0.782
TRA_RD_HDT	MD	HDEUI	[PJ]	0.05
TRA_RD_HDT	MD	HDEUII	[PJ]	0.033
TRA_RD_HDT	MD	HDEUIII	[PJ]	0.029
TRA_RD_HDT	MD	HDEUIV	[PJ]	0.001
TRA_RD_HDT	MD	HDEUV	[PJ]	0.001
TRA_RD_HDT	MD	HDEUVI	[PJ]	0.001
TRA_RD_HDT	MD	HDEUVII	[PJ]	0.001
TRA_RD_HDT	MD	NOC	[PJ]	0.097
TRA_RD_HDT	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_HDT	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD2	GSL	MMO2I	[PJ]	3.95

TRA_RD_LD2	GSL	MMO2II	[PJ]	2.851
TRA_RD_LD2	GSL	MMO2III	[PJ]	2.818
TRA_RD_LD2	GSL	MMO2IV	[PJ]	0.73
TRA_RD_LD2	GSL	MMO2V	[PJ]	0.73
TRA_RD_LD2	GSL	MMO2VI	[PJ]	0.73
TRA_RD_LD2	GSL	NOC	[PJ]	6.864
TRA_RD_LD2	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD2	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	GAS	LFEUI	[PJ]	0.161
TRA_RD_LD4C	GAS	LFEUII	[PJ]	0.079
TRA_RD_LD4C	GAS	LFEUIII	[PJ]	0.041
TRA_RD_LD4C	GAS	LFEUIV	[PJ]	0.025
TRA_RD_LD4C	GAS	LFEUV	[PJ]	0.032
TRA_RD_LD4C	GAS	LFEUVI	[PJ]	0.031
TRA_RD_LD4C	GAS	NOC	[PJ]	0.782
TRA_RD_LD4C	GSL	LFEUI	[PJ]	0.206
TRA_RD_LD4C	GSL	LFEUII	[PJ]	0.101
TRA_RD_LD4C	GSL	LFEUIII	[PJ]	0.053
TRA_RD_LD4C	GSL	LFEUIV	[PJ]	0.032
TRA_RD_LD4C	GSL	LFEUV	[PJ]	0.041
TRA_RD_LD4C	GSL	LFEUVI	[PJ]	0.04
TRA_RD_LD4C	GSL	NOC	[PJ]	0.747
TRA_RD_LD4C	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C	LPG	LFEUI	[PJ]	0.395
TRA_RD_LD4C	LPG	LFEUII	[PJ]	0.193
TRA_RD_LD4C	LPG	LFEUIII	[PJ]	0.108
TRA_RD_LD4C	LPG	LFEUIV	[PJ]	0.053
TRA_RD_LD4C	LPG	LFEUV	[PJ]	0.044
TRA_RD_LD4C	LPG	LFEUVI	[PJ]	0.044
TRA_RD_LD4C	LPG	NOC	[PJ]	0.397
TRA_RD_LD4C	MD	MDEUI	[PJ]	0.02
TRA_RD_LD4C	MD	MDEUII	[PJ]	0.017
TRA_RD_LD4C	MD	MDEUIII	[PJ]	0.009
TRA_RD_LD4C	MD	MDEUIV	[PJ]	0.006
TRA_RD_LD4C	MD	MDEUV	[PJ]	0.007
TRA_RD_LD4C	MD	MDEUVI	[PJ]	0.007
TRA_RD_LD4C	MD	MDEUVII	[PJ]	0.007
TRA_RD_LD4C	MD	MDEUVIp	[PJ]	0.018
TRA_RD_LD4C	MD	MDEUVIt	[PJ]	0.007
TRA_RD_LD4C	MD	NOC	[PJ]	0.051
TRA_RD_LD4C	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4C	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_LD4C_EV	GSL	LFEUI	[PJ]	0.1
TRA_RD_LD4C_EV	GSL	LFEUII	[PJ]	0.1

TRA_RD_LD4C_EV	GSL	LFEUIII	[PJ]	0.1
TRA_RD_LD4C_EV	GSL	LFEUIV	[PJ]	0.1
TRA_RD_LD4C_EV	GSL	LFEUV	[PJ]	0.1
TRA_RD_LD4C_EV	GSL	LFEUVI	[PJ]	0.1
TRA_RD_LD4C_EV	GSL	NOC	[PJ]	0.2
TRA_RD_LD4T	GAS	LFEUI	[PJ]	0.101
TRA_RD_LD4T	GAS	LFEUII	[PJ]	0.044
TRA_RD_LD4T	GAS	LFEUIII	[PJ]	0.027
TRA_RD_LD4T	GAS	LFEUIV	[PJ]	0.016
TRA_RD_LD4T	GAS	LFEUV	[PJ]	0.02
TRA_RD_LD4T	GAS	LFEUVI	[PJ]	0.021
TRA_RD_LD4T	GAS	NOC	[PJ]	0.603
TRA_RD_LD4T	MD	MDEUI	[PJ]	0.043
TRA_RD_LD4T	MD	MDEUII	[PJ]	0.046
TRA_RD_LD4T	MD	MDEUIII	[PJ]	0.03
TRA_RD_LD4T	MD	MDEUIV	[PJ]	0.012
TRA_RD_LD4T	MD	MDEUV	[PJ]	0.012
TRA_RD_LD4T	MD	MDEUVI	[PJ]	0.012
TRA_RD_LD4T	MD	MDEUVII	[PJ]	0.012
TRA_RD_LD4T	MD	MDEUVI _p	[PJ]	0.012
TRA_RD_LD4T	MD	MDEUVI _t	[PJ]	0.012
TRA_RD_LD4T	MD	NOC	[PJ]	0.036
TRA_RD_LD4T	MD_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_LD4T	MD_NV_HE	VEH_IM	[thousand of vehicles]	0
TRA_RD_M4	GSL	MOT4I	[PJ]	0.8
TRA_RD_M4	GSL	MOT4II	[PJ]	0.58
TRA_RD_M4	GSL	MOT4III	[PJ]	0.33
TRA_RD_M4	GSL	MOT4IV	[PJ]	0.439
TRA_RD_M4	GSL	MOT4V	[PJ]	0.439
TRA_RD_M4	GSL	MOT4VI	[PJ]	0.439
TRA_RD_M4	GSL	NOC	[PJ]	1.6
TRA_RD_M4	GSL_NV_HE	NOC	[thousand of vehicles]	0
TRA_RD_M4	GSL_NV_HE	VEH_IM	[thousand of vehicles]	0
TYRES	TYR	INC	[kt]	0.003
TYRES	TYR	NOC	[kt]	0.01
TYRES	TYR	NPR	[kt]	0.003
TYRES	TYR	OPTPR	[kt]	0.007
VEHR_P	PNT	HAMP+SUB1	[kt]	0.28
VEHR_P	PNT	HAMP+SUB2	[kt]	0.197
VEHR_P	PNT	HAMP	[kt]	0.666
VEHR_P	PNT	NOC	[kt]	0.72
VEHTR	POP	NOC	[M people]	0.055
WASTE_AGR	NOF	BAN	[Mt]	0
WASTE_AGR	NOF	NOC	[Mt]	3.4
WASTE_FLR	NOF	FLR_GP	[PJ]	0

WASTE_FLR	NOF	NOC	[PJ]	0.002
WASTE_VOC	EMI	I_LAND	[kt VOC]	0.94
WASTE_VOC	EMI	NOC	[kt VOC]	1
WIRE	ENW	NOC	[kt]	0.017
WIRE	ENW	PRMPLUS	[kt]	0.004
WOOD_P	SC	HSS+PRM	[mln m2]	0.01
WOOD_P	SC	HSS	[mln m2]	0.022
WOOD_P	SC	INC	[mln m2]	0.083
WOOD_P	SC	LSS+PRM+INC	[mln m2]	0.039
WOOD_P	SC	LSS+PRM	[mln m2]	0.163
WOOD_P	SC	MSS+INC	[mln m2]	0.025
WOOD_P	SC	MSS+PRM	[mln m2]	0.046
WOOD_P	SC	MSS	[mln m2]	0.106
WOOD_P	SC	NOC	[mln m2]	0.346
WOOD_P	SC	VHSS+PRM	[mln m2]	0.002
WOOD_P	SC	VHSS	[mln m2]	0.005

Table A9.8 – Emission factors for NH₃ emissions.

Sector	Fuel activity	Measure	Unit	FACTOR_ABTD [kt/unit]
AGR_BEEF	OL	CS	[M animals]	16.883
AGR_BEEF	OL	CS_LNA	[M animals]	10.485
AGR_BEEF	OL	CS_high	[M animals]	16.883
AGR_BEEF	OL	CS_low	[M animals]	20.134
AGR_BEEF	OL	LNA	[M animals]	17.97
AGR_BEEF	OL	LNA_high	[M animals]	17.97
AGR_BEEF	OL	LNA_low	[M animals]	20.677
AGR_BEEF	OL	NOC	[M animals]	23.384
AGR_BEEF	OL	SA	[M animals]	15.502
AGR_BEEF	OL	SA_LNA	[M animals]	8.896
AGR_BEEF	OS	LNA_high	[M animals]	17.97
AGR_BEEF	OS	LNA_low	[M animals]	22.031
AGR_BEEF	OS	NOC	[M animals]	23.384
AGR_COWS	DL	CS	[M animals]	35.465
AGR_COWS	DL	CS_LNA	[M animals]	22.527
AGR_COWS	DL	CS_high	[M animals]	34.371
AGR_COWS	DL	CS_low	[M animals]	40.935
AGR_COWS	DL	LNA	[M animals]	36.448
AGR_COWS	DL	LNA_high	[M animals]	36.448
AGR_COWS	DL	LNA_low	[M animals]	41.973
AGR_COWS	DL	LNf	[M animals]	40.348
AGR_COWS	DL	LNf_CS	[M animals]	30.121
AGR_COWS	DL	LNf_CS_LNA	[M animals]	19.123
AGR_COWS	DL	LNf_LNA	[M animals]	30.956

AGR_COWS	DL	LNF_SA	[M animals]	26.873
AGR_COWS	DL	LNF_SA_LNA	[M animals]	15.366
AGR_COWS	DL	NOC	[M animals]	47.498
AGR_COWS	DL	SA	[M animals]	31.644
AGR_COWS	DL	SA_LNA	[M animals]	18.107
AGR_COWS	DS	LNA_high	[M animals]	30.435
AGR_COWS	DS	LNA_low	[M animals]	37.355
AGR_COWS	DS	LNF	[M animals]	33.691
AGR_COWS	DS	LNF_LNA_high	[M animals]	25.849
AGR_COWS	DS	LNF_LNA_low	[M animals]	31.731
AGR_COWS	DS	NOC	[M animals]	39.661
AGR_OTANI	BS	NOC	[M animals]	43.818
AGR_OTANI	CM	NOC	[M animals]	0.54
AGR_OTANI	FU	NOC	[M animals]	1.712
AGR_OTANI	HO	NOC	[M animals]	7.966
AGR_OTANI	SH	LNA	[M animals]	1.017
AGR_OTANI	SH	LNA_high	[M animals]	1.017
AGR_OTANI	SH	LNA_low	[M animals]	1.243
AGR_OTANI	SH	NOC	[M animals]	1.318
AGR_PIG	PL	BF	[M animals]	4.247
AGR_PIG	PL	BF_CS	[M animals]	3.609
AGR_PIG	PL	BF_CS_LNA	[M animals]	1.35
AGR_PIG	PL	BF_LNA	[M animals]	2.158
AGR_PIG	PL	CS	[M animals]	5.511
AGR_PIG	PL	CS_high	[M animals]	5.453
AGR_PIG	PL	CS_low	[M animals]	5.801
AGR_PIG	PL	LNA	[M animals]	4.06
AGR_PIG	PL	LNA_high	[M animals]	4.06
AGR_PIG	PL	LNA_low	[M animals]	5.105
AGR_PIG	PL	LNF	[M animals]	4.919
AGR_PIG	PL	LNF_BF	[M animals]	3.398
AGR_PIG	PL	LNF_BF_CS	[M animals]	2.887
AGR_PIG	PL	LNF_BF_CS_LNA	[M animals]	1.08
AGR_PIG	PL	LNF_BF_LNA	[M animals]	1.727
AGR_PIG	PL	LNF_CS	[M animals]	4.409
AGR_PIG	PL	LNF_CS_LNA	[M animals]	2.601
AGR_PIG	PL	LNF_LNA	[M animals]	3.248
AGR_PIG	PL	LNF_SA	[M animals]	3.803
AGR_PIG	PL	LNF_SA_LNA	[M animals]	1.834
AGR_PIG	PL	NOC	[M animals]	6.149
AGR_PIG	PL	SA	[M animals]	4.754
AGR_PIG	PL	SA_LNA	[M animals]	2.293
AGR_POULT	LH	BF	[M animals]	0.251
AGR_POULT	LH	BF_CS	[M animals]	0.193
AGR_POULT	LH	BF_CS_LNA	[M animals]	0.134

AGR_POULT	LH	BF_LNA	[M animals]	0.199
AGR_POULT	LH	CS	[M animals]	0.338
AGR_POULT	LH	CS_high	[M animals]	0.338
AGR_POULT	LH	CS_low	[M animals]	0.395
AGR_POULT	LH	LNA	[M animals]	0.343
AGR_POULT	LH	LNA_high	[M animals]	0.298
AGR_POULT	LH	LNA_low	[M animals]	0.371
AGR_POULT	LH	LNF	[M animals]	0.316
AGR_POULT	LH	LNF_BF	[M animals]	0.201
AGR_POULT	LH	LNF_BF_CS	[M animals]	0.155
AGR_POULT	LH	LNF_BF_CS_LNA	[M animals]	0.107
AGR_POULT	LH	LNF_BF_LNA	[M animals]	0.159
AGR_POULT	LH	LNF_CS	[M animals]	0.27
AGR_POULT	LH	LNF_CS_LNA	[M animals]	0.223
AGR_POULT	LH	LNF_LNA	[M animals]	0.275
AGR_POULT	LH	LNF_SA	[M animals]	0.2
AGR_POULT	LH	LNF_SA_LNA	[M animals]	0.144
AGR_POULT	LH	NOC	[M animals]	0.395
AGR_POULT	LH	SA	[M animals]	0.25
AGR_POULT	LH	SA_LNA	[M animals]	0.18
AGR_POULT	OP	BF	[M animals]	0.166
AGR_POULT	OP	BF_CS	[M animals]	0.117
AGR_POULT	OP	BF_CS_LNA	[M animals]	0.084
AGR_POULT	OP	BF_LNA	[M animals]	0.138
AGR_POULT	OP	CS	[M animals]	0.216
AGR_POULT	OP	CS_high	[M animals]	0.216
AGR_POULT	OP	CS_low	[M animals]	0.266
AGR_POULT	OP	LNA	[M animals]	0.237
AGR_POULT	OP	LNA_high	[M animals]	0.212
AGR_POULT	OP	LNA_low	[M animals]	0.252
AGR_POULT	OP	LNF	[M animals]	0.239
AGR_POULT	OP	LNF_BF	[M animals]	0.15
AGR_POULT	OP	LNF_BF_CS	[M animals]	0.105
AGR_POULT	OP	LNF_BF_CS_LNA	[M animals]	0.075
AGR_POULT	OP	LNF_BF_LNA	[M animals]	0.124
AGR_POULT	OP	LNF_CS	[M animals]	0.194
AGR_POULT	OP	LNF_CS_LNA	[M animals]	0.165
AGR_POULT	OP	LNF_LNA	[M animals]	0.213
AGR_POULT	OP	LNF_SA	[M animals]	0.117
AGR_POULT	OP	LNF_SA_LNA	[M animals]	0.081
AGR_POULT	OP	NOC	[M animals]	0.266
AGR_POULT	OP	PM_INC	[M animals]	0.198
AGR_POULT	OP	SA	[M animals]	0.13
AGR_POULT	OP	SA_LNA	[M animals]	0.09
CON_COMB	GAS	IOGCM	[PJ]	0

CON_COMB	GAS	IOGCSC	[PJ]	0.001
CON_COMB	GAS	IOGCSN	[PJ]	0.002
CON_COMB	GAS	NOC	[PJ]	0
CON_COMB	HC1	ISFCM	[PJ]	0
CON_COMB	HC1	ISFCSC	[PJ]	0.002
CON_COMB	HC1	ISFCSN	[PJ]	0.004
CON_COMB	HC1	NOC	[PJ]	0
CON_COMB	HF	IOGCM	[PJ]	0
CON_COMB	HF	IOGCSC	[PJ]	0.002
CON_COMB	HF	IOGCSN	[PJ]	0.003
CON_COMB	HF	NOC	[PJ]	0.001
CON_COMB	LPG	IOGCM	[PJ]	0
CON_COMB	LPG	NOC	[PJ]	0
COWS_3000_MILK	DL	CS	[kt milk]	0.002
COWS_3000_MILK	DL	CS_LNA	[kt milk]	0.001
COWS_3000_MILK	DL	CS_high	[kt milk]	0.002
COWS_3000_MILK	DL	CS_low	[kt milk]	0.002
COWS_3000_MILK	DL	LNA	[kt milk]	0.002
COWS_3000_MILK	DL	LNA_high	[kt milk]	0.002
COWS_3000_MILK	DL	LNA_low	[kt milk]	0.002
COWS_3000_MILK	DL	LNf	[kt milk]	0.002
COWS_3000_MILK	DL	LNf_CS	[kt milk]	0.002
COWS_3000_MILK	DL	LNf_CS_LNA	[kt milk]	0.001
COWS_3000_MILK	DL	LNf_LNA	[kt milk]	0.002
COWS_3000_MILK	DL	LNf_SA	[kt milk]	0.001
COWS_3000_MILK	DL	LNf_SA_LNA	[kt milk]	0.001
COWS_3000_MILK	DL	NOC	[kt milk]	0.002
COWS_3000_MILK	DL	SA	[kt milk]	0.002
COWS_3000_MILK	DL	SA_LNA	[kt milk]	0.001
COWS_3000_MILK	DS	LNA_high	[kt milk]	0.002
COWS_3000_MILK	DS	LNA_low	[kt milk]	0.002
COWS_3000_MILK	DS	LNf	[kt milk]	0.002
COWS_3000_MILK	DS	LNf_LNA_high	[kt milk]	0.001
COWS_3000_MILK	DS	LNf_LNA_low	[kt milk]	0.002
COWS_3000_MILK	DS	NOC	[kt milk]	0.002
DOM	GAS	DGCCOM	[PJ]	0
DOM	GAS	DGCCR	[PJ]	0
DOM	GAS	NOC	[PJ]	0
DOM	LPG	DMDCCO	[PJ]	0
DOM	LPG	DMDCCR	[PJ]	0
DOM	LPG	NOC	[PJ]	0
DOM	MD	DMDCCO	[PJ]	0.001
DOM	MD	DMDCCR	[PJ]	0
DOM	MD	NOC	[PJ]	0.001
DOM	OS1	NOC	[PJ]	0.012

FCON_OTHN	NOF	NOC	[kt N]	0.039
FCON_UREA	NOF	NOC	[kt N]	0.131
FCON_UREA	NOF	SUB_U	[kt N]	0.02
FERTPRO	NOF	NOC	[kt N]	0.003
FERTPRO	NOF	STRIP	[kt N]	0
IN_BO_CHEM	GAS	IOGCM	[PJ]	0
IN_BO_CHEM	GAS	IOGCSC	[PJ]	0.001
IN_BO_CHEM	GAS	IOGCSN	[PJ]	0.002
IN_BO_CHEM	GAS	NOC	[PJ]	0
IN_BO_CHEM	HF	IOGCM	[PJ]	0
IN_BO_CHEM	HF	IOGCSC	[PJ]	0.002
IN_BO_CHEM	HF	ISFCSN	[PJ]	0.003
IN_BO_CHEM	HF	NOC	[PJ]	0.001
IN_BO_CHEM	LPG	IOGCM	[PJ]	0
IN_BO_CHEM	LPG	NOC	[PJ]	0
IN_BO_CON	GAS	IOGCM	[PJ]	0
IN_BO_CON	GAS	IOGCSC	[PJ]	0.001
IN_BO_CON	GAS	IOGCSN	[PJ]	0.002
IN_BO_CON	GAS	NOC	[PJ]	0
IN_BO_CON	HF	IOGCM	[PJ]	0
IN_BO_CON	HF	IOGCSC	[PJ]	0.002
IN_BO_CON	HF	ISFCSN	[PJ]	0.003
IN_BO_CON	HF	NOC	[PJ]	0.001
IN_BO_CON	LPG	IOGCM	[PJ]	0
IN_BO_CON	LPG	NOC	[PJ]	0
IN_BO_OTH	GAS	IOGCM	[PJ]	0
IN_BO_OTH	GAS	IOGCSC	[PJ]	0.001
IN_BO_OTH	GAS	IOGCSN	[PJ]	0.002
IN_BO_OTH	GAS	NOC	[PJ]	0
IN_BO_OTH	HF	IOGCM	[PJ]	0
IN_BO_OTH	HF	IOGCSC	[PJ]	0.002
IN_BO_OTH	HF	ISFCSN	[PJ]	0.003
IN_BO_OTH	HF	NOC	[PJ]	0.001
IN_BO_OTH	OS1	ISFCM	[PJ]	0.003
IN_BO_OTH	OS1	NOC	[PJ]	0.005
IN_BO_OTH_L	HC1	ISFCM	[PJ]	0
IN_BO_OTH_L	HC1	ISFCSC	[PJ]	0.002
IN_BO_OTH_L	HC1	ISFCSN	[PJ]	0.004
IN_BO_OTH_L	HC1	NOC	[PJ]	0
IN_BO_PAP	GAS	IOGCM	[PJ]	0
IN_BO_PAP	GAS	IOGCSC	[PJ]	0.001
IN_BO_PAP	GAS	IOGCSN	[PJ]	0.002
IN_BO_PAP	GAS	NOC	[PJ]	0
IN_BO_PAP	HF	IOGCM	[PJ]	0
IN_BO_PAP	HF	IOGCSC	[PJ]	0.002

IN_BO_PAP	HF	ISFCSN	[PJ]	0.003
IN_BO_PAP	HF	NOC	[PJ]	0.001
IN_OC	DC	NOC	[PJ]	0
IN_OC	GAS	IOGCM	[PJ]	0
IN_OC	GAS	IOGCSC	[PJ]	0.001
IN_OC	GAS	IOGCSN	[PJ]	0.002
IN_OC	GAS	NOC	[PJ]	0
IN_OC	HC1	ISFCM	[PJ]	0
IN_OC	HC1	ISFCSC	[PJ]	0.002
IN_OC	HC1	ISFCSN	[PJ]	0.004
IN_OC	HC1	NOC	[PJ]	0
IN_OC	HF	IOGCM	[PJ]	0
IN_OC	HF	IOGCSC	[PJ]	0.002
IN_OC	HF	ISFCSN	[PJ]	0.003
IN_OC	HF	NOC	[PJ]	0.001
IN_OC	MD	IOGCM	[PJ]	0
IN_OC	MD	NOC	[PJ]	0
IO_NH3_EMISS	NOF	NOC	[kt NH3]	1
OTH_NH3_EMISS	NOF	NOC	[kt NH3]	1
PP_ENG	GAS	LFEUI	[PJ]	0.001
PP_ENG	GAS	LFEUII	[PJ]	0.001
PP_ENG	GAS	LFEUIII	[PJ]	0.001
PP_ENG	GAS	LFEUIV	[PJ]	0.001
PP_ENG	GAS	LFEUV	[PJ]	0.001
PP_ENG	GAS	LFEUVI	[PJ]	0.001
PP_ENG	GAS	NOC	[PJ]	0.001
PP_ENG	MD	NOC	[PJ]	0
PP_ENG	MD	TIWEUI	[PJ]	0
PP_ENG	MD	TIWEUII	[PJ]	0
PP_ENG	MD	TIWEUIII	[PJ]	0
PP_ENG	MD	TIWEUIV	[PJ]	0.001
PP_ENG	MD	TIWEUV	[PJ]	0.001
PP_ENG	MD	TIWEUVI	[PJ]	0.001
PP_EX_OTH	GAS	NOC	[PJ]	0
PP_EX_OTH	GAS	POGCM	[PJ]	0
PP_EX_OTH	GAS	POGCSC	[PJ]	0.001
PP_EX_OTH	HF	NOC	[PJ]	0.001
PP_EX_OTH	HF	POGCM	[PJ]	0
PP_EX_OTH	HF	POGCSC	[PJ]	0.001
PP_EX_OTH	MD	NOC	[PJ]	0
PP_EX_OTH	MD	POGCM	[PJ]	0
PP_EX_OTH	OS2	NOC	[PJ]	0.002
PP_EX_OTH	OS2	PHCCM	[PJ]	0.001
PP_MOD	GAS	NOC	[PJ]	0
PP_MOD	GAS	POGSCR	[PJ]	0.001

PP_NEW	GAS	NOC	[PJ]	0
PP_NEW	GAS	POGSCR	[PJ]	0.001
PP_NEW	GSL	NOC	[PJ]	0
PP_NEW	HF	NOC	[PJ]	0
PP_NEW	HF	POGSCR	[PJ]	0.001
PP_NEW	OS1	NOC	[PJ]	0.003
PP_NEW	OS2	NOC	[PJ]	0.001
PP_NEW	OS2	PHCSCR	[PJ]	0.001
PROD_AGAS	CRU	NOC	[PJ]	0
PROD_AGAS	CRU	REC_USE	[PJ]	0
PROD_AGAS	CRU	REC_USE2	[PJ]	0
TRA_OT_S_L	HF	NOC	[PJ]	0.001
TRA_OT_S_L	HF	STLHCM	[PJ]	0.001
TRA_OT_S_L	HF	STLSCR	[PJ]	0.002
TRA_OT_S_L	MD	NOC	[PJ]	0
TRA_OT_S_L	MD	STLMCM	[PJ]	0
TRA_OT_S_L	MD	STLSCR	[PJ]	0.001
TRA_OT_S_M	MD	NOC	[PJ]	0
TRA_OT_S_M	MD	STMCM	[PJ]	0
TRA_OT_AGR	GSL	LFEUI	[PJ]	0.03
TRA_OT_AGR	GSL	LFEUII	[PJ]	0.012
TRA_OT_AGR	GSL	LFEUIII	[PJ]	0.012
TRA_OT_AGR	GSL	LFEUIV	[PJ]	0.002
TRA_OT_AGR	GSL	LFEUV	[PJ]	0.002
TRA_OT_AGR	GSL	LFEUVI	[PJ]	0.002
TRA_OT_AGR	GSL	NOC	[PJ]	0.001
TRA_OT_AGR	MD	CAGEUI	[PJ]	0
TRA_OT_AGR	MD	CAGEUII	[PJ]	0
TRA_OT_AGR	MD	CAGEUIII	[PJ]	0
TRA_OT_AGR	MD	CAGEUIII_PF	[PJ]	0
TRA_OT_AGR	MD	CAGEUIV	[PJ]	0.001
TRA_OT_AGR	MD	CAGEUV	[PJ]	0.001
TRA_OT_AGR	MD	CAGEUVI	[PJ]	0.001
TRA_OT_AGR	MD	NOC	[PJ]	0
TRA_OT_CNS	MD	CAGEUI	[PJ]	0
TRA_OT_CNS	MD	CAGEUII	[PJ]	0
TRA_OT_CNS	MD	CAGEUIII	[PJ]	0
TRA_OT_CNS	MD	CAGEUIII_PF	[PJ]	0
TRA_OT_CNS	MD	CAGEUIV	[PJ]	0.001
TRA_OT_CNS	MD	CAGEUV	[PJ]	0.001
TRA_OT_CNS	MD	CAGEUVI	[PJ]	0.001
TRA_OT_CNS	MD	NOC	[PJ]	0
TRA_OT_INW	GSL	LFEUI	[PJ]	0.03
TRA_OT_INW	GSL	LFEUII	[PJ]	0.012
TRA_OT_INW	GSL	LFEUIII	[PJ]	0.012

TRA_OT_INW	GSL	LFEUIV	[PJ]	0.002
TRA_OT_INW	GSL	LFEUV	[PJ]	0.002
TRA_OT_INW	GSL	LFEUVI	[PJ]	0.002
TRA_OT_INW	GSL	NOC	[PJ]	0.001
TRA_OT_LB	GAS	LFEUI	[PJ]	0.03
TRA_OT_LB	GAS	LFEUII	[PJ]	0.012
TRA_OT_LB	GAS	LFEUIII	[PJ]	0.012
TRA_OT_LB	GAS	LFEUIV	[PJ]	0.002
TRA_OT_LB	GAS	LFEUV	[PJ]	0.002
TRA_OT_LB	GAS	LFEUVI	[PJ]	0.002
TRA_OT_LB	GAS	NOC	[PJ]	0.001
TRA_OT_LB	GSL	LFEUI	[PJ]	0.03
TRA_OT_LB	GSL	LFEUII	[PJ]	0.012
TRA_OT_LB	GSL	LFEUIII	[PJ]	0.012
TRA_OT_LB	GSL	LFEUIV	[PJ]	0.002
TRA_OT_LB	GSL	LFEUV	[PJ]	0.002
TRA_OT_LB	GSL	LFEUVI	[PJ]	0.002
TRA_OT_LB	GSL	NOC	[PJ]	0.001
TRA_OT_LB	MD	HDEUI	[PJ]	0
TRA_OT_LB	MD	HDEUII	[PJ]	0
TRA_OT_LB	MD	HDEUIII	[PJ]	0
TRA_OT_LB	MD	HDEUIV	[PJ]	0.001
TRA_OT_LB	MD	HDEUV	[PJ]	0.001
TRA_OT_LB	MD	HDEUVI	[PJ]	0.001
TRA_OT_LB	MD	NOC	[PJ]	0
TRA_RD_HDB	GAS	HDSEI	[PJ]	0.003
TRA_RD_HDB	GAS	HDSEII	[PJ]	0.002
TRA_RD_HDB	GAS	HDSEIII	[PJ]	0.001
TRA_RD_HDB	GAS	NOC	[PJ]	0.001
TRA_RD_HDB	MD	HDEUI	[PJ]	0
TRA_RD_HDB	MD	HDEUII	[PJ]	0
TRA_RD_HDB	MD	HDEUIII	[PJ]	0
TRA_RD_HDB	MD	HDEUIV	[PJ]	0
TRA_RD_HDB	MD	HDEUV	[PJ]	0
TRA_RD_HDB	MD	HDEUVI	[PJ]	0
TRA_RD_HDB	MD	HDEUVII	[PJ]	0
TRA_RD_HDB	MD	NOC	[PJ]	0
TRA_RD_HDT	GAS	HDSEI	[PJ]	0.032
TRA_RD_HDT	GAS	HDSEII	[PJ]	0.043
TRA_RD_HDT	GAS	HDSEIII	[PJ]	0.01
TRA_RD_HDT	GAS	NOC	[PJ]	0.001
TRA_RD_HDT	MD	HDEUI	[PJ]	0
TRA_RD_HDT	MD	HDEUII	[PJ]	0
TRA_RD_HDT	MD	HDEUIII	[PJ]	0
TRA_RD_HDT	MD	HDEUIV	[PJ]	0

TRA_RD_HDT	MD	HDEUV	[PJ]	0
TRA_RD_HDT	MD	HDEUVI	[PJ]	0
TRA_RD_HDT	MD	HDEUVII	[PJ]	0
TRA_RD_HDT	MD	NOC	[PJ]	0
TRA_RD_LD2	GSL	MMO2I	[PJ]	0.001
TRA_RD_LD2	GSL	MMO2II	[PJ]	0.002
TRA_RD_LD2	GSL	MMO2III	[PJ]	0.002
TRA_RD_LD2	GSL	MMO2IV	[PJ]	0
TRA_RD_LD2	GSL	MMO2V	[PJ]	0
TRA_RD_LD2	GSL	MMO2VI	[PJ]	0
TRA_RD_LD2	GSL	NOC	[PJ]	0.001
TRA_RD_LD4C	GAS	LFEUI	[PJ]	0.032
TRA_RD_LD4C	GAS	LFEUII	[PJ]	0.043
TRA_RD_LD4C	GAS	LFEUIII	[PJ]	0.01
TRA_RD_LD4C	GAS	LFEUIV	[PJ]	0.01
TRA_RD_LD4C	GAS	LFEUV	[PJ]	0.011
TRA_RD_LD4C	GAS	LFEUVI	[PJ]	0.012
TRA_RD_LD4C	GAS	NOC	[PJ]	0.001
TRA_RD_LD4C	GSL	LFEUI	[PJ]	0.042
TRA_RD_LD4C	GSL	LFEUII	[PJ]	0.055
TRA_RD_LD4C	GSL	LFEUIII	[PJ]	0.013
TRA_RD_LD4C	GSL	LFEUIV	[PJ]	0.013
TRA_RD_LD4C	GSL	LFEUV	[PJ]	0.014
TRA_RD_LD4C	GSL	LFEUVI	[PJ]	0.015
TRA_RD_LD4C	GSL	NOC	[PJ]	0.002
TRA_RD_LD4C	LPG	LFEUI	[PJ]	0.005
TRA_RD_LD4C	LPG	LFEUII	[PJ]	0.003
TRA_RD_LD4C	LPG	LFEUIII	[PJ]	0.001
TRA_RD_LD4C	LPG	LFEUIV	[PJ]	0.001
TRA_RD_LD4C	LPG	LFEUV	[PJ]	0.001
TRA_RD_LD4C	LPG	LFEUVI	[PJ]	0.001
TRA_RD_LD4C	LPG	NOC	[PJ]	0.001
TRA_RD_LD4C	MD	MDEUI	[PJ]	0
TRA_RD_LD4C	MD	MDEUII	[PJ]	0.001
TRA_RD_LD4C	MD	MDEUIII	[PJ]	0.002
TRA_RD_LD4C	MD	MDEUIV	[PJ]	0.002
TRA_RD_LD4C	MD	MDEUV	[PJ]	0.003
TRA_RD_LD4C	MD	MDEUVI	[PJ]	0.003
TRA_RD_LD4C	MD	MDEUVII	[PJ]	0.003
TRA_RD_LD4C	MD	MDEUVIt	[PJ]	0
TRA_RD_LD4C	MD	NOC	[PJ]	0
TRA_RD_LD4T	GAS	LFEUI	[PJ]	0.021
TRA_RD_LD4T	GAS	LFEUII	[PJ]	0.029
TRA_RD_LD4T	GAS	LFEUIII	[PJ]	0.006
TRA_RD_LD4T	GAS	LFEUIV	[PJ]	0.006

TRA_RD_LD4T	GAS	LFEUV	[PJ]	0.007
TRA_RD_LD4T	GAS	LFEUVI	[PJ]	0.007
TRA_RD_LD4T	GAS	NOC	[PJ]	0.001
TRA_RD_LD4T	MD	MDEUI	[PJ]	0
TRA_RD_LD4T	MD	MDEUII	[PJ]	0.001
TRA_RD_LD4T	MD	MDEUIII	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUIV	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUV	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUVI	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUVII	[PJ]	0.002
TRA_RD_LD4T	MD	MDEUVIt	[PJ]	0
TRA_RD_LD4T	MD	NOC	[PJ]	0
TRA_RD_M4	GSL	MOT4I	[PJ]	0.002
TRA_RD_M4	GSL	MOT4II	[PJ]	0.002
TRA_RD_M4	GSL	MOT4III	[PJ]	0.002
TRA_RD_M4	GSL	MOT4IV	[PJ]	0
TRA_RD_M4	GSL	MOT4V	[PJ]	0
TRA_RD_M4	GSL	MOT4VI	[PJ]	0
TRA_RD_M4	GSL	NOC	[PJ]	0.001
WASTE_AGR	NOF	BAN	[Mt]	0
WASTE_AGR	NOF	NOC	[Mt]	1.3
WASTE_FLR	NOF	FLR_GP	[PJ]	0
WASTE_FLR	NOF	NOC	[PJ]	0
WT_NH3_EMISS	NOF	NOC	[kt NH3]	1

